ISSN 0378–4347

# CUMULATIVE AUTHOR AND SUBJECT INDEXES VOLUMES 221 —230

Supplied free of charge to the 1982 subscribers of the Journal of Chromatography, Biomedical Applications

# JOURNAL OF CHROMATOGRAPHY BIOMEDICAL APPLICATIONS

EDITOR, K. Macek (Prague)

CONSULTING EDITOR, M. Lederer (Switzerland)

EDITORIAL BOARD
R. F. Adams (North Ryde)
B. G. Belenkii (Leningrad)
L. D. Bergelson (Moscow)
A. A. Boulton (Saskatoon)
C. J. W. Brooks (Glasgow)
P. R. Brown (Kingston, RI)
W. L. Chiou (Chicago, IL)
H. Ch. Curtius (Zürich)
J. A. F. de Silva (Nutley, NJ)
Z. Deyl (Prague)
R. A. de Zeeuw (Groningen)
J. W. Drysdale (Boston, MA)
F. M. Everaerts (Eindhoven)
M. G. Horning (Houston, TX)
E. Jellum (Oslo)
A. Kuksis (Toronto)
H. M. Liebich (Tübingen)
M. Novotný (Bloomington, IN)
P. Padieu (Dijon)
J. Roboz (New York, NY)
N. Seiler (Strasbourg)
J. Sjövall (Stockholm)
L. R. Snyder (Yorktown Heights, NY)
W. J. A. VandenHeuvel (Rahway, NJ)
J. Vessman (Mölndal)
J. Wagner (Leipzig)

EDITOR, NEWS AND BOOK REVIEW SECTIONS
Z. Deyl (Prague)

UNIVERSITY OF HAWAII
LIBRARIES

FEB 08 1983

LIBRARY USE ONLY

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM

PERIODICAL
QD
271
.J6
Index
221-230
U.H. LIBRARY

JOURNAL OF CHROMATOGRAPHY

**Scope.** The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications,* which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews,* reviews on all aspects of chromatography, electrophoresis and related methods are published.

**Submission of Papers.** Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications.* An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

**Subscription Orders.** Subscription orders should be sent to: Elsevier Scientific Publishing Company, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

**Publication.** The *Journal of Chromatography* (incl. *Biomedical Applications, Chromatographic Reviews* and *Cumulative Author and Subject Indexes, Vols. 221–230, 231–240* and *241–250*) has 25 volumes in 1982. The subscription prices for 1982 are:

*J. Chromatogr.* (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 221–230, 231–240* and *241–250*) + *Biomed. Appl.* (Vols. 227–251):
Dfl. 3625.00 plus Dfl. 500.00 (postage) (total *ca.* US$ 1650.00)

*J. Chromatogr.* (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 231–240* and *241–250*) only (Vols. 234–251):
Dfl. 2826.00 plus Dfl. 360.00 (postage) (total *ca.* US$ 1274.50)

*Biomed. Appl.* (incl. *Cum. Indexes Vols. 221–230*) only (Vols 227–233):
Dfl. 1050.00 plus Dfl. 140.00 (postage) (total *ca.* US$ 476.00).

Journals are automatically sent by airmail to the U.S.A. and Canada at no extra costs, and to Japan, Australia and New Zealand, with a small additional postal charge. Back volumes of the *Journal of Chromatography* (Vols. 1 through 226) are available at Dfl. 173.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. For customers in the U.S.A. and Canada wishing additional bibliographic information on this and other Elsevier journals, please contact Elsevier Science Publishing Company Inc., Journal Information Center, 52 Vanderbilt Avenue, New York, NY 10017. Tel: (212) 867-9040.

**Abstracts/Contents Lists** published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Index Medicus, Mass Spectrometry Bulletin, and Science Citation Index.

**See page 3 of cover** for Publication Schedule, Information for Authors, and information on the News Section and Advertisements.

---

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY — 1982

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

**Special regulations for readers in the U.S.A.** This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

**Special regulations for authors in the U.S.A.** Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

For advertising information please contact our advertising representatives

## U.S.A./CANADA

**Leni Ellinger**

3133 Connecticut Ave, NW
Suite 712
WASHINGTON, DC 20008
Tel.: (202) 232-3366

## GREAT BRITAIN

**T.G. Scott & Son Ltd.**

Mr. M.L. White
30-32 Southampton Street
LONDON, WC2E 7HE
Tel.: (01) 379-7264

**OR**

General Advertising Department

**Elsevier/ Excerpta Medica/ North-Holland**

Ms W. van Cattenburch
P.O. Box 211
1000 AE AMSTERDAM
The Netherlands
Tel.: (020) 5803.714/715
Telex: 18582 ESP NL

631EVe

SPECAC

**SWITCHING AND INJECTION VALVES FOR HPLC**

NEW RANGE includes:

* 4 port injector for microbore work - available with 0.2 to 5µl sample loops.
* High performance centre injection valve for use up to 7000 psi.

All valves feature:

* Low torque for ease of switching
* Excellent reproducibility
* All tube connections on front face
* Bracket or panel mounting facility

**ANALYTICAL ACCESSORIES LIMITED**
Unit 3, Lagoon Road, St. Mary Cray,
Orpington, Kent, BR5 3QX

002

*Please mention this journal when answering advertisements*

★

015E

# Pyrolysis Mass Spectrometry of Recent and Fossil Biomaterials

## Compendium and Atlas

*by* H. L. C. MEUZELAAR, *Salt Lake City, UT, U.S.A.*, J. HAVERKAMP, *Amsterdam, The Netherlands and* F. D. HILEMAN, *Dayton, OH, U.S.A.*
**Techniques and Instrumentation in Analytical Chemistry, Vol. 3**

Since the 3rd International Symposium on Analytical Pyrolysis in 1976, the importance of pyrolysis mass spectrometry (Py-MS) as an analytical method has increased considerably. Specially designed Py-MS systems using galvanically heated filament or direct probe pyrolysis have been quite successful in structural investigations and kinetic studies involving synthetic polymers and model compounds. The Curie-point Py-MS has, however, made a lasting impression on the field, being unique in those applications that require maximum reproducibility. This book presents an in-depth discussion of basic principles (including sample requirements, short- and long-term reproducibility problems and data processing approaches) of Py-MS techniques.

The Compendium has been put together by authors who have themselves been deeply involved in an extensively used fully-automated Py-MS system. It is thought that this system, particularly useful in characterising bacteria, could revolutionise clinical methods which have not changed since Pasteur.

The advent of commercially available Curie-point Py-MS systems has prompted the authors to include a small Atlas of over 150 reference spectra of carefully selected biomaterials which should help users of these systems to "tune" their instruments to the existing systems and to evaluate unknown spectra.

**An asset to any chemistry library and particularly useful to those scientists considering the application of Py-MS techniques to their own specific problems in the analysis and data processing of biomaterials.**

**Contents: Part I. Compendium of Basic Principles and Applications.**
1. Origins and Development of Pyrolysis Mass Spectrometry of Biomaterials. 2. From Fingerprinting to Structural Investigation. 3. Pyrolysis Mechanisms in Biomaterials. 4. The Technique of Curie-Point Py-MS. 5. Reproducibility in Curie-Point Py-MS. 6. Data Analysis Procedures. 7. Selected Applications to Biomaterials. **Part II. Atlas of Selected Pyrolysis Mass Spectra.**

1982 xiv + 294 pages
US $67.50/Dfl. 145.00
ISBN 0-444-42099-1

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
P.O. Box 211
1000 AE Amsterdam, The Netherlands.

ELSEVIER SCIENCE PUBLISHING CO. INC.
52 Vanderbilt Avenue
New York, NY 10017, U.S.A.

7246

# ELECTROPHORESIS

## A Survey of Techniques and Applications

*edited by* Z. DEYL, *Czechoslovak Academy of Sciences, Prague*

JOURNAL OF CHROMATOGRAPHY LIBRARY, 18

## PART A: TECHNIQUES

Z. DEYL *(editor)*
F. M. EVERAERTS, Z. PRUSÍK *and*
P. J. SVENDSEN *(co-editors)*

*"... provides a sound, state-of-the-art survey of its subject".*

– Chemistry in Britain

*"... the editors have set out to bring everything together into a coherent whole... they have succeeded remarkably well... the book is bound to be well liked and appreciated by readers".*

– Journal of Chromatography

This first part deals with the principles, theory and instrumentation of modern electromigration methods. Both standard procedures and newer developments are discussed and hints are included to help the reader overcome difficulties frequently arising from the lack of suitable equipment. Adequate theoretical background of the individual techniques is given and a theoretical approach to the deteriorative processes is presented to facilitate further development of a particular technique and its application to a special problem. In each chapter practical realisations of different techniques are described and examples are presented to demonstrate the limits of each method.

CONTENTS:
Introduction. Chapters: 1. Theory of electromigration processes *(J. Vacík)*. 2. Classification of electromigration methods *(J. Vacík)*. 3. Evaluation of the results of electrophoretic separations *(J. Vacík)*. 4. Molecular size and shape in electrophoresis *(Z. Deyl)*. 5. Zone electrophoresis (except gel-type techniques and immunoelectrophoresis) *(W. Ostrowski)*. 6. Gel-type techniques *(Z. Hrkal)*. 7. Quantitative immunoelectrophoresis *(P.J. Svendsen)*. 8. Moving boundary electrophoresis in narrow-bore tubes *(F.M. Everaerts and J.L. Beckers)*. 9. Isoelectric focusing *(N. Catsimpoolas)*. 10. Analytical isotachophoresis *(J. Vacík and F.M. Everaerts)*. 11. Continuous flow-through electrophoresis *(Z. Prusík)*. 12. Continuous flow deviation electrophoresis *(A. Kolin)*. 13. Preparative electrophoresis in gel media *(Z. Hrkal)*. 14. Preparative electrophoresis in columns *(P.J. Svendsen)*. 15. Preparative isoelectric focusing *(P. Blanický)*. 16. Preparative isotachophoresis *(P.J. Svendsen)*. 17. Preparative isotachophoresis on the micro scale *(L. Arlinger)*. List of frequently occurring symbols. Subject Index.

1979 xvi + 390 pp. US $72.25/Dfl. 170.00
ISBN 0-444-41721-4

## PART B: APPLICATIONS

Z. DEYL *(editor)*
A. CHRAMBACH, F.M. EVERAERTS *and*
Z. PRUSÍK *(co-editors)*

Part B is an exhaustive survey of the present status of the application of electrophoretic techniques to many diverse compounds. Those categories of compounds most suited to these separations, such as proteins and peptides, are dealt with in detail, while the perspectives of the applications of these techniques to other categories of compounds less commonly electrophoresed are given. Special attention is paid to naturally occurring mixtures of compounds and their treatment. This is the first attempt to cover the field on such a broad scale and the book will be valuable to separation chemists, pharmacologists, organic chemists and those involved in biomedical research.

CONTENTS: 1. Alcohols and phenolic compounds *(Z. Deyl)*. 2. Aldehydes and ketones *(Z. Deyl)*. 3. Carbohydrates *(Z. Deyl)*. 4. Carboxylic acids *(F.M. Everaerts)*. 5. Steroids and steroid conjugates *(Z. Deyl)*. 6. Amines *(Z. Deyl)*. 7. Amino acids and their derivatives *(Z. Deyl)*. 8. Peptides and structural analysis of proteins *(Z. Prusík)*. 9. Gel electrophoresis and electrofocusing of proteins *(edited by A. Chrambach)*. Usefulness of second-generation gel electrophoretic tools in protein fractionation *(A. Chrambach)*. Membrane proteins, native *(L.M. Hjelmeland)*. Membrane proteins, denatured *(H. Baumann, D. Doyle)*. Protein membrane receptors *(U. Lang)*. Steroid receptors *(S. Ben-Or)*. Cell surface antigens *(R.A. Reisfeld, M.A. Pellegrino)*. Lysosomal glycosidases and sulphatases *(A.L. Fluharty)*. Heamocyanins *(M. Brenowitz et al.)*. Human haemoglobins *(A.B. Schneider, A.N. Schechter)*. Isoelectric focusing of immunoglobulins *(M.H. Freedman)*. Contractile and cytoskeletal proteins *(P. Rubenstein)*. Proteins of connective tissue *(Z. Deyl, M. Horáková)*. Microtubular proteins *(K.F. Sullivan, L. Wilson)*. Protein hormones *(A.D. Rogol)*. Electrophoresis of plasma proteins: a contemporary clinical approach *(M. Engliš)*. Allergens *(H. Baer, M.C. Anderson)*. 10. Glycoproteins and glycopeptides (affinity electrophoresis) *(T.C. Bøg-Hansen, J. Hau,)*. 11. Lipoproteins *(H. Peeters)*. 12. Lipopolysaccharides *(P.F. Coleman, O. Gabriel)*. 13. Electrophoretic examination of enzymes *(W. Ostrowski)*. 14. Nucleotides, nucleosides, nitrogenous constituents of nucleic acids *(S. Zadražil)*. 15. Nucleic acids *(S. Zadražil)*. 16. Alkaloids *(Z. Deyl)*. 17. Vitamins *(Z. Deyl)*. 18. Antibiotics *(V. Betina)*. 19. Dyes and pigments *(Z. Deyl)*. 20. Inorganic compounds *(F.M. Everaerts, Th. P.E.M. Verheggen)*. Contents of "Electrophoresis, Part A: Techniques". Subject Index. Index of compounds separated.

1982 xiii + 462 pp. US $104.75/Dfl. 225.00
ISBN 0-444-42114-9



ELSEVIER
P.O. Box 211, Amsterdam
The Netherlands

52 Vanderbilt Avenue
New York, NY 10017, U.S.A.

7255

# STATISTICAL TREATMENT OF EXPERIMENTAL DATA

*By* J.R. GREEN, *Lecturer in Computational and Statistical Science, University of Liverpool, U.K. and* D. MARGERISON, *Senior Lecturer in Inorganic, Physical and Industrial Chemistry, University of Liverpool, U.K.*

**PHYSICAL SCIENCES DATA 2**

This book first appeared in 1977. In 1978 a revised reprint was published and in response to demand, further reprints appeared in 1979 and 1980. Intended for researchers wishing to analyse experimental data, this work will also be useful to students of statistics. Statistical methods and concepts are explained and the ideas and reasoning behind statistical methodology clarified. Noteworthy features of the text are numerical worked examples to illustrate formal results, and the treatment of many practical topics which are often omitted from standard texts, for example testing for outliers, stabilization of variances and polynomial regression.

**What the reviewers had to say:**

*"The index is detailed; the format is good; the presentation is clear; and no mathematics beyond calculus is assumed".*
*–CHOICE*

*"A lot of thought has gone into this book and I like it very much. It deserves a place on every laboratory bookshelf".*
*–CHEMISTRY IN BRITAIN*

**1977. Reprinted 1978, 1979, 1980.**
**xiv + 382 pages**
**US $39.25/Dfl.90.00**
**ISBN: 0-444-41725-7**

**ELSEVIER**

P.O. Box 211, 1000 AE Amsterdam, The Netherlands.
52 Vanderbilt Ave., New York, NY 10017.

*The Dutch guilder price is definitive. US$ prices are subject to exchange rate fluctuations.*

7/84

# JOURNAL
*of*
# CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY,
ELECTROPHORESIS AND RELATED METHODS

# BIOMEDICAL APPLICATIONS

*Cumulative Author and Subject Indexes*
*Volumes 221—230 (1981—1982)*

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM
1982

Electronic data processing:
Büro für Satztechnik W. Meyer KG
Weissensberg, G.F.R.

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY — 1982

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

**Special regulations for readers in the U.S.A.** This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

**Special regulations for authors in the U.S.A.** Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

# AUTHOR INDEX

## VOLUMES 221—230

*The volume number is placed in parentheses before the page number*

# A


Abang, A., see Swezey, S.F. (227) 510
Abbott, F.S., Burton, R., Orr, J., Wladichuk, D., Ferguson, S. and Sun, T.-H.
Valproic acid analysis in saliva and serum using selected ion monitoring (electron ionization) of the *tert.* -butyldimethylsilyl derivatives (227) 433
Abdel-Monem, M.M. and Merdink, J.L.
Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363
Abe, K., see Shimada, K. (227) 445
Abernethy, D.R., Greenblatt, D.J. and Zumbo, A.M.
Antipyrine determination in human plasma by GLC using nitrogen-phosphorus detection (223) 432
Abernethy, D.R., see Ameer, B. (226) 224
Abeywardene, S.A., see Glencross, R.G. (223) 193
Abreu, R.A. de, see De Abreu, R.A. (227) 45, (227) 526, (229) 67
Abshagen, U., see Endele, R. (227) 187
Ackerly, C.C., Newman, R.A., Myers, C. and McCormack, J.J.
LC separation and quantitation of 2-amino-1,3,4-thiadiazole (NSC-4728) from human and murine serum (230) 175
Adachi, M., see Katogi, Y. (228) 404
Adachi, Y., see Matsumura, K. (222) 53
Adam, H.K., Douglas, E.J., Plummer, G.F. and Cosgrove, M.B.
Estimation of ICI 35,868 (Diprivan) in blood by HPLC, following coupling with Gibbs' reagent (223) 232
Adam, M., see Deyl, Z. (230) 409
Adams, J.S., Clemens, T.L. and Holick, M.F.
Silica Sep-Pak preparative chromatography for vitamin D and its metabolites (226) 198
Aerde, P. van, see Van Aerde, P. (222) 467
Agris, P.F., see Gehrke, C.W. (230) 297
Aherne, G.W., see Lawson, G.J. (223) 225
Ahmed, A.E. and Hsu, T.-F.
Quantitative analysis of melphalan and its major hydrolysate in patients and animals by reversed-phase HPLC (222) 453
Ahokas, J.T., see Norris, R.L.G. (230) 433
Aizawa, Y., see Yamada, K. (223) 176
Akagi, H., see Matsushima, E. (230) 188
Akedo, H. and Shinkai, K.
Isotachophoretic determination of adriamycin and adriamycinol in human plasma (227) 262
Al-Noaemi, M.C., see McDermott, J.R. (222) 371
Albaigés, J., see Caturla, M.C. (227) 219
Albani, F., Riva, R. and Baruzzi, A.
Simple and rapid determination of propranolol and its active metabolite, 4-hydroxypropranolol, in human plasma by LC with fluorescence detection (228) 362
Albani, F., see Riva, R. (221) 75, (222) 491, (225) 219
Albert, J., see Pahuja, S.L. (225) 37
Albessard, F., see Moulin, M.A. (226) 250
Alderweireldt, F., see Marescau, B. (224) 185
Alexander, N.M. and Neeley, W.E.
Chromatofocusing of human hemoglobins (230) 137
Ali, H.M. and Beckett, A.H.
Rapid method for the determination of chlorpheniramine in urine (223) 208
Allain, P., see Turcant, A. (229) 222
Allen, B.A., Newman, R.A., Griffin, J.P. and McCormack, J.J.
LC determination of an antineoplastic aziridinylbenzoquinone in human and murine serum (222) 146
Allenmark, S., Berg, A.A., Hammar, M. and Lindström, E.
Combined PC and reversed-phase HPLC method for the study of pregnenolone and progesterone metabolites (224) 399
Allenmark, S. and Qureshi, G.A.
Studies on dopamine-converting enzymes in human plasma (223) 188
Alric, R., Cociglio, M., Blayac, J.P. and Puech, R.
Performance evaluation of a reversed-phase, HPLC assay of valproic acid involving a "solvent demixing" extraction procedure and precolumn derivatisation (224) 289

Althaus, Z.R., Rowland, J.M., Freeman, J.P. and Slikker, W. Jr.
Separation of some natural and synthetic corticosteroids in biological fluids and tissues by HPLC (227) 11
Alton, K.B.
Determination of the antifungal agent, ketoconazole, in human plasma by HPLC (221) 337
Alván, G., Ekman, L. and Lindström, B.
Determination of chloroquine and its desethyl metabolite in plasma, red blood cells in urine by LC (229) 241
Alvinerie, M., Galtier, P. and Escoula, G.
Ion-pair HPLC assay of levamisole in biological fluids (223) 445
Alvinerie, M. and Toutain, P.L.
Determination of xylazine in plasma using HPLC (222) 308
Ambrose, R.T., see Lauff, J.J. (226) 391
Ameer, B., Greenblatt, D.J., Divoll, M., Abernethy, D.R. and Shargel, L.
HPLC determination of acetaminophen in plasma: single-dose pharmacokinetic studies (226) 224
Anastassiades, T., Puzic, R. and Puzic, O.
Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309
Andermann, G. and Dietz, M.
Simultaneous determination of cyclandelate and its metabolite in human plasma by capillary column GLC (223) 365
Anderson, G.M., Young, J.G., Batter, D.K., Young, S.N., Cohen, D.J. and Shaywitz, B.A.
Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315
Anderson, G.M., Young, J.G., Cohen, D.J., Shaywitz, B.A. and Batter, D.K.
Amperometric determination of 3-methoxy-4-hydroxyphenylethyleneglycol in human cerebrospinal fluid (222) 112
Anderson, G.M., Young, J.G., Cohen, D.J. and Young, S.N.
Determination of indoles in human and rat pineal (228) 155
Anderson, R.D., Ilett, K.F., Dusci, L.J. and Hackett, L.P.
HPLC analysis of pentazocine in blood and plasma (227) 239
Andresen, B.D., see Ng, K.J. (228) 43
Andrews, P.A., Egorin, M.J., May, M.E. and Bachur, N.R.
Reversed-phase HPLC analysis of 6-thioguanine applicable to pharmacologic studies in humans (227) 83
Anjaneyulu, B., see Rajagopalan, T.G. (224) 265
Anna, M. de, see De Anna, M. (222) 491
Annesley, T.M., see Hay, I.D. (226) 383
Arai, M., see Koh, S. (226) 461
Aravind, M.K., Miceli, J.N. and Kauffman, R.E.
Determination of moxalactam by HPLC (228) 418
Aravind, M.K., Miceli, J.N., Kauffman, R.E., Strebel, L.E. and Done, A.K.
Simultaneous measurement of chloramphenicol and chloramphenicol succinate by HPLC (221) 176
Arimori, S., see Hiramatsu, K. (227) 423, (229) 455
Arita, T., see Miyazaki, K. (223) 449
Arranz Peña, M.I.
Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486
Arranz Peña, M.I.
Rapid GC determination of valproic acid in serum (225) 459
Asada, H., see Niwa, T. (230) 1
Asada, S., see Takada, K. (222) 156
Ascalone, V.
Assay of trimethoprim, sulfadiazine and its $N^4$-acetyl metabolite in biological fluids by normal-phase HPLC (224) 59
Aso, Y., see Noda, A. (230) 345
Astier, A. and Deutsch, A.M.
Quantitative HPLC determination of antispasmodic trimebutine in human plasma. Pharmacokinetic studies after intravenous administration in humans (224) 149
Au, J.L.-S., Wientjes, M.G., Luccioni, C.M. and Rustum, Y.M.
Reversed-phase ion-pair HPLC assay of 5-fluorouracil, 5'-deoxy-5-fluorouridine, their nucleosides, mono-, di-, and triphosphate nucleotides with a mixture of quaternary ammonium ions (228) 245
Axelson, J.E., see Pillai, G.K. (229) 103
Axelson, M., Sahlberg, B.-L. and Sjövall, J.
Analysis of profiles of conjugated steroids in urine by ion-exchange separation and GC-MS (224) 355

# B

Baal, J.M. van, see Van Baal, J.M. (227) 45, (227) 526, (229) 67
Baars, A.M., see Vree, T.B. (224) 519
Bachur, N.R., see Andrews, P.A. (227) 83
Badcock, N.R. and Pollard, A.C.
Micro-determination of clonazepam in plasma or serum by electron-capture GLC (230) 353
Bahr, C. von, see Von Bahr, C. (221) 109, (227) 113

Bailey, E., Della Corte, L., Farmer, P.B. and Gray, A.J.
Determination of the antioxidant 3-*tert.*-butyl-4-hydroxyanisole in rat plasma using high-resolution GC-MS (225) 83
Baker, G.B., see LeGatt, D.F. (225) 301
Baker, J.K., McChesney, J.D., Hufford, C.D. and Clark, A.M.
HPLC analysis of the metabolism of primaquine and the identification of a new mammalian metabolite (230) 69
Bakkeren, J.A.J.M., see De Abreu, R.A. (227) 45, (229) 67
Balali-Mood, M., King, I.S. and Prescott, L.F.
Rapid estimation of diflunisal in plasma and urine by HPLC and a comparison with a fluorometric method (229) 234
Balish, E., see Witter, J.P. (229) 450
Bannier, A., Brazier, J.L., Soubeyrand, J. and Comet, F.
Determination of a new anti-inflammatory agent, 1-isobutyl-3,4-diphenylpyrazole-5-acetic acid, by HPLC (227) 213
Bannier, A., see Brazier, J.L. (224) 439
Bannon, M.J., see Michaud, R.L. (225) 335
Barazi, S., see Bonini, M. (224) 332
Barchowsky, A., see Shand, D.G. (224) 343
Bardócz, S. and Karsai, T.
Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198
Bareggi, S.R., see Zecca, L. (230) 168
Barends, D.M., Zwaan, C.L. and Hulshoff, A.
Improved microdetermination of gentamicin and sisomicin in serum by HPLC with ultraviolet detection (222) 316
Barends, DM., Zwaan, C.L. and Hulshoff, A.
Micro-determination of tobramycin in serum by HPLC with UV detection (225) 417
Barfod, N.M.
HPLC purification of endogenous tumour cell growth inhibitory peptides (230) 289
Bargnoux, H., see Lartigue-Mattei, C. (229) 211
Barlogie, B., see Plunkett, W. (221) 425
Barlow, E.C., see Hundt, H.K.L. (223) 165
Barnicoat, A.J., Van 't Hoff, W.G., Morrison, P.J. and Bradbrook, I.D.
Determination of salicylhydroxamic acid, a trypanocidal agent, by reversed-phase HPLC (225) 236
Barrett, E., see Jacob, R. (229) 188
Baruzzi, A., see Albani, F. (228) 362
Baruzzi, A., see Riva, R. (221) 75, (222) 491, (225) 219
Basset, P., Braconnier, F. and Rosa, J.
An update on electrophoretic and chromatographic methods in the diagnosis of hemoglobinopathies (Review) (227) 267
Batter, D.K., see Anderson, G.M. (222) 112, (223) 315
Bauer, L.A., see Watts, M.T. (230) 79
Baughman, R.A. Jr., see Mico, B.A. (230) 203
Baumann, P., see Tinguely, D. (229) 319
Baune, A., Bromet, N., Courte, S. and Voisin, C.
Trace determination of almitrine in plasma by GLC using a nitrogen-phosphorus detector (223) 219
Bautista-Cerqueira, S., see Krstulović, A.M. (227) 379
Bayer, E., see Frank, H. (224) 177
Bayley, A.C., see Lovelace, C.E.A. (227) 256
Beales, D., Finch, R., McLean, A.E.M., Smith, M. and Wilson, I.D.
Determination of penicillamine and other thiols by combined HPLC and post-column reaction with Ellman's reagent: application to human urine (226) 498
Beaudoin, N., see Ong, H. (229) 433
Bechgaard, E. and Nielsen, A.
Determination of bromhexine in human plasma and urine by HPLC (228) 392
Beckett, A.H., see Ali, H.M. (223) 208
Bedford, J.A., see Jones, A.B. (226) 99
Bélanger, P.M. and Grech-Bélanger, O.
GLC determination of plasma and urinary levels of amantadine in man (228) 327
Belkien, L., see Schöneshöfer, M. (227) 492
Bellon, G., Bisker, A., Maquart, F.X., Thoanes, H. and Borel, J.P.
Improved method for the evaluation of 3-hydroxyproline and 4-hydroxyproline in the urine (230) 420
Benassi, E., see Ranise, A. (222) 120
Benedict, J.J., see Chester, T.L. (225) 17
Benet, L.Z., see Buskin, J.N. (230) 454
Benet, L.Z., see Mico, B.A. (230) 203
Benjamin, R.S., see Benvenuto, J.A. (222) 518
Benowitz, N.L., see Jacob, P. III (222) 61
Benschop, H.P., Konings, K.A.G., Kossen, S.P. and Ligtenstein, D.A.
Determination of some pyridinium aldoxime compounds by means of ion-pair reversed-phase HPLC: application in biological material (225) 107
Benvenuto, J.A., Stewart, D.J., Benjamin, R.S., Smith, R.G. and Loo, T.L.
HPLC analysis of pentamethylmelamine and its metabolites in biological fluids (222) 518
Benvenuto, J.A., see Miller, A.A. (228) 165
Berg, A.A., see Allenmark, S. (224) 399
Berg, M. van den, see Van den Berg, M. (226) 259
Berger, J.A., see Chabard, J.L. (221) 9
Berger, J.A., see Lartigue-Mattei, C. (229) 211
Bergqvist, Y. and Eckerbom, S.
Simultaneous determination of chloroquine and its desethyl metabolites in human plasma by GC (226) 91
Bergqvist, Y. and Frisk-Holmberg, M.
Sensitive method for the determination of chloroquine and its metabolite desethyl-chloroquine in human plasma and urine by HPLC (221) 119
Bergstrom, R.F., Kay, D.R. and Wagner, J.G.
HPLC determination of penicillamine in whole blood, plasma, and urine (222) 445

Bernard, N., Cuisinaud, G. and Sassard, J.
Determination of penbutolol and its hydroxylated metabolite in biological fluids by reversed-phase HPLC (228) 355
Bernstein, M.S. and Evans, M.A.
HPLC-fluorescence analysis for indomethacin and metabolites in biological fluids (229) 179
Berry, G., see Yandrasitz, J.R. (225) 319
Bertani-Dziedzic, L., see Krstulovic, A.M. (223) 305
Bertani-Dziedzic, L., see Krstulović, A.M. (227) 379
Bertrand, O., see Kroviarski, Y. (228) 298
Beschke, K., see Roth, W. (222) 13
Besio, G., see Ranise, A. (222) 120
Besio-Moreno, M., see Bizzozero, O. (227) 33
Bevill, R.F., see McCarthy, T.P. (225) 9
Bianchine, J.R., see Ng, K.J. (228) 43
Biddlecome, C.E., see Park, G.B. (227) 534
Bieder, A., see Le Roux, Y. (230) 401
Biggins, J.A., see McDermott, J.R. (222) 371
Bilodeau, G., see Losito, R. (226) 61
Binz, K., see Mueller, H.W. (228) 75
Biou, D., Queyrel, N., Visseaux, M.N., Collignon, I. and Pays, M.
Separation and identification of dansylated human serum and urinary amino acids by two-dimensional TLC. Application to aminoacidopathies (226) 477
Birkett, D.J., see Chivers, D.A. (225) 261
Bishop-Freudling, G.B. and Vergin, H.
Determination of 2,4-diamino-5-benzylpyrimidines in combination with sulphadiazine in biological fluids using HPLC (224) 301
Bisker, A., see Bellon, G. (230) 420
Bizzozero, O., Besio-Moreno, M., Pasquini, J.M., Soto, E.F. and Gómez, C.J.
Rapid purification of proteolipids from rat brain subcellular fractions by chromatography on a lipophilic dextran gel (227) 33
Blackman, G.L., see Harman, P.J. (225) 131
Blanchard, J.
Evaluation of the relative efficacy of various techniques for deproteinizing plasma samples prior to HPLC (226) 455
Blanford, M.F., see Wu, C.C. (228) 333
Blaschke, T.F., see Swezey, S.F. (227) 510
Blass, K.G., see Briand, R.L. (223) 277
Blayac, J.P., see Alric, R. (224) 289
Blendis, L.M., see Goldberg, E.M. (226) 291
Boček, P., see Dolník, V. (225) 455
Bockbrader, H.N., see Guay, D.R.P. (228) 398
Böcker, R.
Rapid analysis of hexobarbital and its main metabolites in mice by HPLC (228) 350
Boivin, P., see Kroviarski, Y. (228) 298
Boksay, I., see Hui, K.-S. (222) 512
Bols, N.C. and Mosser, D.D.
Separation by TLC of nucleotides from bases and nucleosides in trichloroacetic acid extracts of cells (229) 460
Bondy, P.K. and Canellakis, Z.N.
HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371
Bonini, M., Mokofio, F. and Barazi, S.
Contribution to the determination of four antimalaria drugs, chloroquine, quinine, pyrimethamine and sulfadoxine, alone or in mixtures in biological fluids. II. GC (in French) (224) 332
Boppana, V.K., see Swanson, B.N. (225) 123
Borel, J.P., see Bellon, G. (230) 420
Borsa, M., see Crolla, T. (222) 257
Borstrok, J.J.M., see Moleman, P. (227) 391
Boschi, S. and Marchesini, B.
HPLC method for the simultaneous determination of iothalamate and *o*-iodohippurate (224) 139
Bosin, T.R., see Mais, D.E. (225) 27
Boujet, C., Douet, C., Decoux, G. and Favier, A.
Study of the extraction of organic acids from urine. Preliminary step to their GC separation (in French) (228) 67
Boulton, A.A., see Davis, B.A. (222) 161, (222) 271, (230) 219
Boux, L.J., Ireland, C.M., Wright, D.J., Holder, G.M. and Ryan, A.J.
TLC and HPLC separation of metabolites of the weak carcinogen, 7-methylbenz[*c*]acridine (227) 149
Bovill, J.G., see Van Rooy, H.H. (223) 85
Bowen, D.V., see Ching, N.P.H. (222) 171
Bowman, D.B. and Kauffman, R.E.
Reversed-phase HPLC method to determine vidarabine and hypoxanthine arabinoside in biological fluids (229) 487
Boxtel, C.J. van, see Van Boxtel, C.J. (226) 259
Braconnier, F., see Basset, P. (227) 267
Bradbrook, I.D., see Barnicoat, A.J. (225) 236
Braeckman, P., see Van Aerde, P. (222) 467
Braithwaite, R., see Broquaire, M. (224) 43
Braithwaite, R.A., see Guinebault, P.R. (223) 103
Brandt, A.E., see Thomas, D.W. (225) 433
Brass, C., see Swezey, S.F. (227) 510
Braughler, J.M., see McGinley, P.A. (230) 29
Brazier, J.L., Delaye, D., Desage, M. and Bannier, A.
Simultaneous microdetermination of theophylline, caffeine and phenobarbital in blood collected on paper (224) 439
Brazier, J.L., see Bannier, A. (227) 213
Breccia, A., see Gattavecchia, E. (224) 465
Breckenridge, W.C., see Kuksis, A. (224) 1, (230) 231
Bredenkamp, B.L.F. and Joubert, S.M.
The effect of ampholytes on ferritin isoelectric focussing patterns (230) 327
Bree, F., see Houin, G. (223) 351
Breese, G.R., see Kilts, C.D. (225) 347
Breimer, D.D., see Heusler, H. (226) 403
Breimer, D.D., see Jochemsen, R. (223) 438, (227) 199

Breithaupt, H. and Goebel, G.
Determination of allopurinol and oxipurinol in biological fluids by HPLC (226) 237
Breithaupt, H. and Schick, J.
Determination of cytarabine and uracil arabinoside in human plasma and cerebrospinal fluid by HPLC (225) 99
Breithaupt, H. and Wilfling, M.
Determination of mexiletine in biological fluids by HPLC (230) 97
Brenner, R., see Tschöpe, W. (222) 41
Briand, R.L., Harold, S. and Blass, K.G.
HPLC determination of the lecithin/sphingomyelin ratio in amniotic fluid (223) 277
Brien, J.F., see Loomis, C.W. (222) 421
Brisson, A.M. and Fourtillan, J.B.
Determination of cephalosporins in biological material by reversed-phase LCC (223) 393
Brockerhoff, P., see Höckel, M. (221) 205
Brodie, B., see McCarthy, T.P. (225) 9
Brodie, R.R. and Chasseaud, L.F.
Determination of 11-bromovincamine in human plasma by HPLC (228) 413
Brodie, R.R., Chasseaud, L.F. and Walmsley, L.M.
Determination of the diuretic agent metolazone in plasma by HPLC (226) 526
Bromet, N., see Baune, A. (223) 219
Bromet, N., see Courte, S. (224) 162
Brooks, M.A., Strojny, N., Hackman, M.R. and De Silva, J.A.F.
Determination of bromo-lasalocid in plasma by HPLC with fluorimetric detection (229) 167
Brooks, M.A., see Hackman, M.R. (222) 179
Brooks, M.A., see Lee, T.L. (224) 275, (227) 137
Broquaire, M., Rovei, V. and Braithwaite, R.
Quantitative determination of naproxen in plasma by a simple HPLC method (224) 43
Broquaire, M., see Guinebault, P.R. (223) 103
Brosemer, R.W., see Gurusiddaiah, S. (223) 179
Brown, A.R., see DeAngelis, R.L. (221) 353
Brown, J.E., Wilkinson, P.A. and Brown, J.R.
Rapid HPLC assay for the anthracyclines daunorubicin and 7-con-O-methylnogarol in plasma (226) 521
Brown, J.R., see Brown, J.E. (226) 521
Brown, L.W., see Hundt, H.K.L. (225) 482
Brown, M.J., see Jenner, D.A. (224) 507
Brown, N.D., Poon, B.T. and Chulay, J.D.
Determination of chloroquine and its de-ethylated metabolites in human plasma by ion-pair HPLC (229) 248
Brown, P.R., see Zakaria, M. (226) 267, (230) 381
Brown, R.D., see Robert, T.A. (224) 116
Bruckner, G., see Mai, J. (230) 15
Bruinvis, L., see Kamerling, J.P. (222) 276
Brunton, J., see Rubin, P.C. (221) 193
Bruyn, C. de, see De Bruyn, C. (225) 369
Bruyn, C.H.M.M. de, see De Bruyn, C.H.M.M. (227) 45, (227) 526, (229) 67
Buchwald, W., see Cone, E.J. (223) 331
Buchwald, W.F., see Cone, E.J. (230) 57
Budna, K.W., see Lankmayr, E.P. (222) 249
Buggé, C.J.L., see Vane, F.M. (227) 471
Burford, S.R., see Vose, C.W. (222) 311
Burkman, A.M., see Wu, C.C. (228) 333
Burrows, G.D., see Maguire, K.P. (222) 399
Burton, R., see Abbott, F.S. (227) 433
Buskin, J.N., Upton, R.A., Jones, R.M. and Williams, R.L.
HPLC assay of acebutolol and two of its metabolites in plasma and urine (230) 438
Buskin, J.N., Upton, R.A., Sörgel, F., Williams, R.L., Lang, E. and Benet, L.Z.
Specific and sensitive assay of celiprolol in blood, plasma and urine using HPLC (230) 454
Buskin, J.N., Upton, R.A. and Williams, R.L.
Improved acetaminophen assay sensitivity by modification of a HPLC technique (230) 443
Bye, A., see Dean, K. (221) 408
Bye, A., see Land, G. (222) 135, (224) 51
Bye, A., see Lo, L.Y. (222) 297

# C

Cabana, B.E., see Rapaka, R.S. (227) 463
Cacciapuoti, G., see Della Ragione, F. (226) 243
Cailleux, A., see Turcant, A. (229) 222
Calatroni, A., see Pietta, P. (228) 377, (229) 445
Calder, I.C., see Hart, S.J. (225) 387
Callaghan, O.H., see Mosnaim, A.D. (224) 481
Calvey, T.N., see Chan, K. (223) 213
Cam, G.R., see Kreutzmann, D.J. (228) 95
Campos-Flor, S., see Kiang, C.-H. (222) 81
Camsonne, R., see Moulin, M.A. (226) 250
Canellakis, Z.N., see Bondy, P.K. (224) 371
Cani, D., see Cotellessa, L. (221) 149
Cani, D., see Mussini, E. (224) 94
Cannell, G.R., Mortimer, R.H. and Thomas, M.J.
HPLC estimation of cyproterone acetate in human plasma (226) 492
Cano, J.P., Sumirtapura, Y.C., Cautreels, W. and Sales, Y.
Analysis of the metabolites of ethyl loflazepate by GC with electron-capture detection (226) 413
Cano, J.P., see Sommadossi, J.P. (225) 516, (228) 205
Cano, J.P., see Viala, A. (224) 503
Canoy, D.W., see Morris, J.E. (224) 407
Cantoni, L., Ruggieri, R., Dal Fiume, D. and Rizzardini, M.
The determination of uroporphyrinogen decarboxylase in tissues by HPLC coupled to fluorescence detection (229) 311

Capogrossi, M.C., Holdiness, M.R. and Israili, Z.H.
Determination of adenosine in normal human plasma and serum by HPLC (227) 168
Carenzi, A., see Longo, A. (224) 257
Carey, P.F., Martin, L.E. and Owen, P.E.
Determination of ranitidine and its metabolites in human urine by reversed-phase ion-pair HPLC (225) 161
Carlyle, R.F., see Chauhan, J. (227) 305
Carnis, G., see Godbillon, J. (222) 461
Caron, J.C., Martin, B. and Shroot, B.
HPLC determination of urocanic acid isomers in biological samples (230) 125
Carruthers, M.E., see Causon, R.C. (229) 301
Cartier, P., see Rocchiccioli, F. (226) 325
Cassara, G., Gianazza, E., Righetti, P.G., Poma, S., Vicentini, L. and Scortecci, V.
Isoelectric focusing studies of human pancreatic secretion (221) 279
Castagnoli, N. Jr., see Shinohara, Y. (230) 363
Castellucci, M.G., see Ghebregzabher, M. (222) 191
Caturla, M.C. and Albaigés, J.
Rapid determination of procetofenic acid in plasma by HPLC (227) 219
Caudal, F., see Sioufi, A. (221) 419
Caude, M., see Guyon, F. (229) 475
Caudill, W.L., Houck, G.P. and Wightman, R.M.
Determination of $\gamma$-aminobutyric acid by LC with electrochemical detection (227) 331
Causon, R.C. and Carruthers, M.E.
Measurement of catecholamines in biological fluids by HPLC. A comparison of fluorimetric with electrochemical detection (229) 301
Causon, R.C., Collins, S.L. and Fry, D.E.
Determination of urinary dehydroepiandrosterone sulphate by combined HPLC and radioimmunoassay (227) 485
Cautreels, W., see Cano, J.P. (226) 413
Chabard, J.L., Lartigue-Mattei, C., Vedrine, F., Petit, J. and Berger, J.A.
Mass fragmentographic determination of xanthine and hypoxanthine in biological fluids (221) 9
Chabard, J.L., see Lartigue-Mattei, C. (229) 211
Chamberlain, J., see Dell, D. (226) 431
Chambers, D.M., White, M.H. and Kostenbauder, H.B.
Efficient extraction and reversed-phase HPLC-UV quantitation of acetazolamide in serum (225) 231
Chan, J.Y.Y., see Lim, C.K. (228) 305
Chan, K., Murray, G.R., Rostron, C., Calvey, T.N. and Williams, N.E.
Quantitative GLC method for the determination of phenoperidine in human plasma (223) 213
Chan, W.-Y., see Griesmann, G.E. (230) 121
Chang, S.F., Hansen, C.S., Fox, J.M. and Ober, R.E.
Quantitative determination of nefopam in human plasma, saliva and cerebrospinal fluid by GLC using a nitrogen-selective detector (226) 79
Charette, C., McGilveray, I.J. and Midha, K.K.
GLC procedure with alkali flame ionization detection for the determination of maprotiline in plasma (224) 128
Charles, B.G., Cole, J.J. and Ravenscroft, P.J.
Method for rapid determination of urinary tetracycline by HPLC (222) 152
Chasseaud, L.F., see Brodie, R.R. (226) 526, (228) 413
Chasseaud, L.F., see Walmsley, L.M. (226) 155, (226) 441
Chatterji, D.C., see Narang, P.K. (230) 373
Chauhan, J., Darbre, A. and Carlyle, R.F.
Determination of urinary amino acids by means of glass capillary GLC with alkali-flame ionisation detection and flame ionisation detection (227) 305
Chauhan, M.S. and Dakshinamurti, K.
GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323
Chen, M.-L. and Chiou, W.L.
Sensitive and rapid HPLC method for the simultaneous determination of methotrexate and its metabolites in plasma, saliva and urine (226) 125
Chen, S.S.-H. and Kou, A.Y.
Improved procedure for the separation of phospholipids by HPLC (227) 25
Cheng, K.-P., see Hui, K.-S. (222) 512
Chester, T.L., Lewis, E.C., Benedict, J.J., Sunberg, R.J. and Tettenhorst, W.C.
Determination of (dichloromethylene)diphosphonate in physiological fluids by ion-exchange chromatography with phosphorus-selective detection (225) 17
Cheung, L.K., see Smith, R.G. (229) 464
Chew, P., see Reeve, J. Jr. (229) 57
Ching, N.P.H., Jham, G.N., Subbarayan, C., Bowen, D.V., Smit, A.L.C. Jr., Grossi, C.E., Hicks, R.G., Field, F.H. and Nealon, T.F. Jr.
GC-MS detection of circulating plasticizers in surgical patients (222) 171
Ching, N.P.H., Jham, G.N., Subbarayan, C., Grossi, C., Hicks, R. and Nealon, T.F. Jr.
GC quantitation of two plasticizers contaminating intravenous fluids stored in plastic containers (225) 196
Chiou, W.L., see Chen, M.-L. (226) 125
Chiu, A., see Warsh, J.J. (228) 131
Chivers, D.A., Birkett, D.J. and Miners, J.O.
Simultaneous determination of pentoxifylline and its hydroxy metabolite in plasma by HPLC (225) 261
Choi, R.L., Rosenberg, M., Grebow, P.E. and Huntley, T.E.
HPLC analysis of indapamide (RHC 2555) in urine, plasma and blood (230) 181
Christman, N.T. and Hamilton, L.H.
A new chromatographic instrument for

measuring trace concentrations of breath-hydrogen (229) 259
Chubb, S., see Plunkett, W. (221) 425
Chulay, J.D., see Brown, N.D. (229) 248
Churchill, F.C. II, see Staiger, M.A. (225) 139
Cisse, H., Farinotti, R., Kirkiacharian, S. and Dauphin, A.
LC determination of valproic acid in plasma with spectrofluorimetric detection (in French) (225) 509
Clark, A.M., see Baker, J.K. (230) 69
Clemens, T.L., see Adams, J.S. (226) 198
Cloppet, H., see Francina, A. (222) 116, (228) 177
Coates, P.E., see Walker, S.E. (223) 131
Cochet, S., see Krovíarski, Y. (228) 298
Cociglio, M., see Alric, R. (224) 289
Cocks, D.A., Dyer, T.F. and Edgar, K.
Simple and rapid GLC method for estimating carbamazepine in serum (222) 496
Coco, S.J., see Dokladalova, J. (224) 33
Cohen, D.J., see Anderson, G.M. (222) 112, (223) 315, (228) 155
Cole, D.E.C. and Scriver, C.R.
Microassay of inorganic sulfate in biological fluids by controlled flow anion chromatography (225) 359
Cole, J.J., see Charles, B.G. (222) 152
Cole, K.D. and Little, G.H.
Isocratic HPLC of bile pigments (227) 503
Collignon, I., see Biou, D. (226) 477
Collins, S.L., see Causon, R.C. (227) 485
Colombo, L., see Cotellessa, L. (221) 149
Colombo, L., see Mussini, E. (224) 94
Colvin, M., see Strife, R.J. (224) 168
Comet, F., see Bannier, A. (227) 213
Cone, E.J., Buchwald, W. and Yousefnejad, D.
Simultaneous determination of phencyclidine and monohydroxylated metabolites in urine of man by GC-MF with methane chemical ionization (223) 331
Cone, E.J., Gorodetzky, C.W., Yeh, S.Y., Darwin, W.D. and Buchwald, W.F.
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57
Cooper, J.D.H. and Turnell, D.C.
Fluorescence detection of cystine by *o*-phthalaldehyde derivatisation and its separation using HPLC (227) 158
Cooper, S.F. and Lapierre, Y.D.
GLC determination of pipotiazine in plasma of psychiatric patients (222) 291
Cooper, T.B., see Jindal, S.P. (221) 301
Cooper, T.B., see Suckow, R.F. (230) 391
Coppin, F., see Dell, D. (226) 431
Corcia, A. di, see Di Corcia, A. (229) 365
Corney, S.J., see Glencross, R.G. (223) 193
Corran, P.H., see Zanelli, J.M. (223) 59
Corte, L. della, see Della Corte, L. (225) 83
Cosgrove, M.B., see Adam, H.K. (223) 232
Cotellessa, L., Marcucci, F., Cani, D., Sfondrini, P., Colombo, L., Mussini, E. and Poy, F.
Glass capillary GC determination of $N^{\tau}$-methylhistidine in urine (221) 149
Cotellessa, L., see Mussini, E. (224) 94
Cotler, S., Puglisi, C.V. and Gustafson, J.H.
Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95
Čoupek, J., see Macek, K. (222) 284
Courte, S. and Bromet, N.
Trace determination of trimetazidine in plasma by HPLC using fluorescence detection (224) 162
Courte, S., see Baune, A. (223) 219
Coutts, R.T., see LeGatt, D.F. (225) 301
Cozamanis, I., see Reece, P.A. (225) 151
Crolla, T., Borsa, M. and Pifferi, G.
HPLC determination of trithiozine and its neutral metabolites in human plasma and urine (222) 257
Crowne, H., Lim, C.K. and Samson, D.
Determination of 5-aminolaevulinic acid dehydrase activity in erythrocytes by HPLC (223) 421
Cuisinaud, G., see Bernard, N. (228) 355
Curry, S.H., see Whelpton, R. (228) 321

# D

Daigneault, E.A., see Robert, T.A. (224) 116
Dakshinamurti, K., see Chauhan, M.S. (227) 323
Dal Fiume, D., see Cantoni, L. (229) 311
Danielson, N.D. and Huth, J.A.
Direct assay for creatine kinase by reversed-phase HPLC (221) 39
Dankers, J., see Leferink, J.G. (229) 217
Danzin, C., see Wagner, J. (227) 349
Darbre, A., see Chauhan, J. (227) 305
D'Arconte, L., see Strojny, N. (223) 111
Darwin, W.D., see Cone, E.J. (230) 57
Dauphin, A., see Cisse, H. (225) 509
Davis, B.A. and Boulton, A.A.
Longitudinal urinary excretion of some "trace" acids in a human male (222) 161
Davis, B.A. and Boulton, A.A.
Excretion of *m*-hydroxymandelic acid in human urine (222) 271
Davis, B.A., Durden, D.A. and Boulton, A.A.
Plasma concentrations of *p*- and *m*-hydroxyphenylacetic acid and phenylacetic acid in humans. GC-high-resolution MS analysis (230) 219
Davis, C.M. and Fenimore, D.C.
Rapid microanalysis of anticonvulsants by HPTLC (222) 265
Davis, K.L., see Hsieh, J.Y.-K. (225) 521

Davis, T.P., Gehrke, C.W. Jr., Williams, C.H., Gehrke, C.W. and Gerhardt, K.O.
Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113
Dayton, D.A., see Mohammed, H.Y. (226) 471
De Abreu, R.A., Van Baal, J.M., Bakkeren, J.A.J.M., De Bruyn, C.H.M.M. and Schretlen, E.D.A.M.
HPLC assay for identification and quantitation of nucleotides in lymphocytes and malignant lymphoblasts (227) 45
De Abreu, R.A., Van Baal, J.M., De Bruyn, C.H.M.M., Bakkeren, J.A.J.M. and Schretlen, E.D.A.M.
HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67
De Abreu, R.A., Van Baal, J.M., Schouten, T.J., Schretlen, E.D.A.M. and De Bruyn, C.H.M.M.
HPLC determination of plasma 6-mercaptopurine in clinically relevant concentrations (227) 526
De Anna, M., see Riva, R. (222) 491
De Bruyn, C., see Oerlemans, F. (225) 369
De Bruyn, C.H.M.M., see De Abreu, R.A. (227) 45, (227) 526, (229) 67
De Jong, J.W., Hegge, J.A.J., Harmsen, E. and De Tombe, P.P.
Fluorometric LC assay of the antiarrhythmic agent flecainide in blood plasma (229) 498
De Jong, J.W., see Harmsen, E. (230) 131
De Neef, J.H., Porsius, A.J. and Van Rooy, H.H.
Quantitative analysis of the cholinesterase inhibitor paraoxon in brain tissue using HPLC (224) 133
De Silva, J.A.F., see Brooks, M.A. (229) 167
De Silva, J.A.F., see Lee, T.L. (224) 275
De Silva, J.A.F., see Pao, J. (221) 97
De Silva, J.A.F., see Puglisi, C.V. (226) 135
De Silva, J.A.F., see Strojny, N. (223) 111
De Tombe, P.P., see De Jong, J.W. (229) 498
De Tombe, P.P., see Harmsen, E. (230) 131
De Vries, J.X., Günthert, W. and Ding, R.
Determination of nicotinamide in human plasma and urine by ion-pair reversed-phase HPLC (221) 161
De Wildt, D.J., Porsius, A.J. and Van Rooy, H.H.
Quantitative determination of the cholinesterase inhibitor physostigmine in brain tissue samples using reversed-phase HPLC (225) 381
Dean, K., Land, G. and Bye, A.
Analysis of procyclidine in human plasma and urine by GLC (221) 408
Dean, K., see Land, G. (222) 135
DeAngelis, R.L., Robinson, M.M., Brown, A.R., Johnson, T.E. and Welch, R.M.
Quantitation of the anticonvulsant cinromide (3-bromo-N-ethylcinnamamide) and its major plasma metabolites by TLC (221) 353
Decoux, G., see Boujet, C. (228) 67
Dege, P.H. and Ervik, M.
Determination of prenalterol in plasma and in urine by GLC (222) 437
Deheny, T., see Vijayakumar, R. (225) 47
Delaunay, J., see Francina, A. (228) 177
Delaye, D., see Brazier, J.L. (224) 439
Dell, D., Chamberlain, J. and Coppin, F.
Determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC (226) 431
Della Corte, L., see Bailey, E. (225) 83
Della Ragione, F., Varteni-Farina, M., Porcelli, M., Cacciapuoti, G. and Zappia, V.
HPLC analysis of 5'-methylthioadenosine in rat tissues (226) 243
Desage, M., see Brazier, J.L. (224) 439
Deturmeny, E., see Viala, A. (224) 503
Deutsch, A.M., see Astier, A. (224) 149
Deutsch, E., see Zodda, J.P. (227) 249
Deyl, Z., Macek, K., Adam, M. and Horáková, M.
High-performance gel permeation chromatography of collagens (230) 409
Deyl, Z., see Macek, K. (222) 284
Dhar, A.K. and Kutt, H.
Improved GC procedure for the determination of clonazepam levels in plasma using a nitrogen-sensitive detector (222) 203
Di Corcia, A., Ripani, L. and Samperi, R.
Chromatographic micro-procedure for trace determination of phenobarbital in blood serum (229) 365
Di Toro, M., see Longo, A. (224) 257
Dickens, J.P., see Vose, C.W. (225) 488
Dietz, M., see Andermann, G. (223) 365
Dijkstra, D., see Feenstra, M.G.P. (230) 271
Dimaline, R., see Reeve, J. Jr. (229) 57
D'Incalci, M., see Farina, P. (222) 141
Ding, R., see De Vries, J.X. (221) 161
Divoll, M. and Greenblatt, D.J.
Plasma concentrations of temazepam, a 3-hydroxy benzodiazepine, determined by electron-capture GLC (222) 125
Divoll, M., see Ameer, B. (226) 224
Divoll, M., see Greenblatt, D.J. (225) 202
Dixon, P.F., see Lawson, G.J. (223) 225
Dokladalova, J., Coco, S.J., Lemke, P.R., Quercia, G.T. and Korst, J.J.
Determination of polythiazide and prazosin in human plasma by HPLC (224) 33
Dolník, V. and Boček, P.
Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455
Done, A.K., see Aravind, M.K. (221) 176
Dooley, T. and Maddocks, J.L.
Assay of an active metabolite of 6-thioguanine, 6-thioguanosine 5'-monophosphate, in human red blood cells (229) 121

Dorleac, E., see Francina, A. (222) 116, (228) 177
Døssing, M., see Hansen, S.H. (229) 141
Douet, C., see Boujet, C. (228) 67
Douglas, E.J., see Adam, H.K. (223) 232
Drehsen, G. and Rohdewald, P.
Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479
Drehsen, G., see Rohdewald, P. (225) 427
Drenthe-Schonk, A., see Linssen, P. (223) 371
Dring, L.G., see Gillet, G. (230) 154
Driskell, J.A., see Tangney, C.C. (224) 389
Duchoň, J., see Pavel, S. (222) 329
Dulce, H.J., see Schöneshöfer, M. (222) 478, (227) 492
Dünges, W., see Höckel, M. (221) 205
Dunn, G.L., see Gupta, R.N. (224) 496
Duran, M., see Kamerling, J.P. (222) 276
Durand, A., see Viala, A. (224) 503
Durden, D.A., see Davis, B.A. (230) 219
Dusci, L.J., see Anderson, R.D. (227) 239
Duvaldestin, P., see Meulemans, A. (226) 255
Dyer, T.F., see Cocks, D.A. (222) 496
Dziedzic, S.W., see Krstulovic, A.M. (223) 305

# E

Eadie, M.J., see Hooper, W.D. (223) 426
Eckerbom, S., see Bergqvist, Y. (226) 91
Edelson, J., see Park, G.B. (222) 213, (227) 534
Edenharder, R. and Slemr, J.
GC and MS analysis of bile acids as trifluoroacetyl-hexafluoroisopropyl and heptafluorobutyryl derivatives (222) 1
Edgar, K., see Cocks, D.A. (222) 496
Edlund, P.O.
Determination of ergot alkaloids in plasma by HPLC and fluorescence detection (226) 107
Edwardson, J.A., see McDermott, J.R. (222) 371
Egan, C.M., Jones, C.R. and McCluskey, M.
Method for the measurement of melphalan in biological samples by HPLC (224) 338
Egorin, M.J., see Andrews, P.A. (227) 83
Eichelman, B., see Hegstrand, L.R. (222) 107
Eichhorst, O. and Hinderling, P.H.
Specific assay for radiolabelled digoxin and its known apolar metabolites in biological fluids. I (224) 67
Eimiller, A., see Frank, H. (224) 177
Ekman, L., see Alván, G. (229) 241
El-Sharaky, A.S., see Reisch, J. (222) 475
Ellard, G.A., see Jenner, P.J. (225) 245
Ellin, R.I., Zvirblis, P. and Wilson, M.R.
Method for isolation and determination of pyridostigmine and metabolites in urine and blood (228) 235
Elsohly, M.A., see Jones, A.B. (226) 99
Emanuelsson, B.-M., see Hartvig, P. (228) 340
Endele, R., Senn, M. and Abshagen, U.
Determination of deacetylmetipranolol in body fluids by GC-chemical-ionization MS (227) 187
Eng Tho Gan, I., see Schier, G.M. (225) 208
Epping, J., see Heusler, H. (226) 403
Erb, R., see Williams, C. II (225) 225
Erdtmansky, P., see Park, G.B. (222) 213
Eriksson, B.-M. and Persson, B.-A.
Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143
Eriksson, B.-M., Tekenbergs, L., Magnusson, J.-O. and Molin, L.
Determination of tritiated digoxin and metabolites in urine by LC (223) 401
Ervik, M., Kylberg-Hanssen, K. and Lagerström, P.-O.
Assay of prenalterol in plasma and urine by GC-MS (229) 87
Ervik, M., see Dege, P.H. (222) 437
Erwin, V.G., see Towell, J.F. (223) 295
Escoula, G., see Alvinerie, M. (223) 445
Esmans, E., see Marescau, B. (224) 185
Esposito, C., see Porta, R. (226) 208
Estadieu, M., see Viala, A. (224) 503
Evans, G.L., see Vose, C.W. (222) 311
Evans, M.A., see Bernstein, M.S. (229) 179
Everaerts, F., see Oerlemans, F. (225) 369

# F

Faber, D.B., see Kok, R.M. (222) 389
Fagerlund, C., see Hartvig, P. (228) 340
Fandrich, S.E. and Pittman, K.A.
Analysis of carminomycin in human serum by fluorometric HPLC (223) 155
Farina, P., Marzillo, G. and D'Incalci, M.
HPLC determination of 4'-demethylepipodophyllotoxin-9-(4,6-0-ethylidene β-D-glucopyranoside) (VP 16-213) in human plasma (222) 141
Farinotti, R., see Cisse, H. (225) 509
Farmer, P.B., see Bailey, E. (225) 83
Favier, A., see Boujet, C. (228) 67
Febvre, N., see Lecaillon, J.B. (228) 257
Feenstra, M.G.P., Homan, J.W., Dijkstra, D., Mulder, T.B.A., Rollema, H., Westerink, B.H.C. and Horn, A.S.
Reversed-phase LC with amperometric detection of lipophilic dopamine analogues and determination of brain and serum concentrations after sample clean-up on small Sephadex G-10 columns (230) 271
Felten, D.L., see Rhodes, G. (228) 33
Felten, D.L., see Smith, S.L. (221) 19

Fencik, M., see Hui, K.-S. (222) 512
Fenimore, D.C., see Davis, C.M. (222) 265
Fenner, A., see Schöneshöfer, M. (224) 229, (224) 472
Ferguson, S., see Abbott, F.S. (227) 433
Fernstrom, J., see Spindel, E. (222) 381
Field, F.H., see Ching, N.P.H. (222) 171
Finch, R., see Beales, D. (226) 498
Fischer, C., Maier, K. and Klotz, U.
Simplified HPLC method for 5-aminosalicylic acid in plasma and urine (225) 498
Fisher, L., see Spindel, E. (222) 381
Fiume, D. dal, see Dal Fiume, D. (229) 311
Flanagan, R.J., see Gettings, S.D. (225) 469
Fleitman, J., Torosian, G. and Perrin, J.H.
Improved HPLC assay for cimetidine using ranitidine as an internal standard (229) 255
Fleitman, J.S., Schulman, S.G. and Perrin, J.H.
HPLC assay for fenbufen and two serum metabolites (228) 372
Floberg, S., Hartvig, P., Lindström, B., Lönnerholm, G. and Odlind, B.
Extractive alkylation of 6-mercaptopurine and determination in plasma by GC-MS (225) 73
Floberg, S., Lindström, B. and Lönnerholm, G.
Simultaneous determination of theophylline and caffeine after extractive alkylation in small volumes of plasma by GC-MS (221) 166
Fornstedt, N.
HPLC determination of homovanillic acid in urine using Sephadex G-10 for isolation (225) 440
Fornstedt, N.
A method for HPLC screening of UV-positive components in urine eluate from Sephadex G-10 and modifications for determination of urinary salicylic, salicyluric and gentisic acids (230) 253
Forte, A.J., see Wilson, J.M. (227) 453
Fournier, P.E., see Tournet, M.-C. (224) 348
Fourtillan, J.B., see Brisson, A.M. (223) 393
Fourtillan, J.B., see Lefebvre, M.A. (230) 199
Fox, J.M., see Chang, S.F. (226) 79
Fozard, J.R., see Grove, J. (223) 409
Fraisse-André, J., see Gillet, G. (230) 154
Francina, A., Dorleac, E. and Cloppet, H.
Carrier-ampholyte displacement chromatography (chromatofocusing) on ion-exchange papers. Application to the separation of haemoglobin variants (222) 116
Francina, A., Dorleac, E., Cloppet, H. and Delaunay, J.
Chromatofocusing of human hemoglobins: application to the quantitation of hemoglobin $A_2$ (228) 177
Frank, H., Eimiller, A., Kornhuber, H.H. and Bayer, E.
GC determination of glutamine in the presence of glutamic acid in biological fluids (224) 177
Freeman, J.P., see Althaus, Z.R. (227) 11
Fremouw-Ottevangers, D.C., see Muskiet, F.A.J. (230) 142
Frenkel, E.P., see Strauss, J.F. (221) 139
Frey, B.M. and Frey, F.J.
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283
Frey, F.J., see Frey, B.M. (229) 283
Friedman, Z., see Hancock, W.S. (228) 273
Frisk-Holmberg, M., see Bergqvist, Y. (221) 119
Fry, D.E., see Causon, R.C. (227) 485
Fuchs, D., see Hausen, A. (227) 61
Fukunishi, T., see Takada, K. (222) 156
Fuller, R.K., see Neiderhiser, D.H. (229) 470
Furukawa, H., see Niwa, T. (230) 1

# G

Gabryś, J. and Konecki, J.
GC analysis of free amino acids in the hyaloplasm of the hypophysis, pineal gland, thyroid gland, spinal cord, thymus and lymph nodes of the cow (222) 345
Gaillot, J., see Le Roux, Y. (230) 401
Gaillot, J., see Sommadossi, J.P. (228) 205
Galloway, D.B., see Pickup, M.E. (225) 493
Galtier, P., see Alvinerie, M. (223) 445
Gans, E.H., see Yeung, D. (224) 513
Gasa, S., see Kaji, H. (221) 145
Gates, S.C., see Rehman, A. (228) 103
Gatley, S.J., see Witter, J.P. (229) 450
Gattavecchia, E., Tonelli, D. and Breccia, A.
Determination of metronidazole, misonidazole and its metabolite in serum and urine on RP-18 HPTLC plates (224) 465
Gattiker, H., see Losito, R. (226) 61
Gauron, S., see Godbillon, J. (227) 516
Gauron, S., see Sioufi, A. (222) 429
Geher, K., see Kuksis, A. (224) 1, (230) 231
Gehrke, C.W., Kuo, K.C., McCune, R.A., Gerhardt, K.O. and Agris, P.F.
Quantitative enzymatic hydrolysis of tRNAs. Reversed-phase HPLC of tRNA nucleosides (230) 297
Gehrke, C.W., see Davis, T.P. (228) 113
Gehrke, C.W. Jr., see Davis, T.P. (228) 113
Gelijkens, C.F., Smith, D.L. and McCloskey, J.A.
Capillary GC of pyrimidines and purines: N,O-peralkyl and trifluoroacetyl-N,O-alkyl derivatives (225) 291
Gerhardt, K.O., see Davis, T.P. (228) 113
Gerhardt, K.O., see Gehrke, C.W. (230) 297
Gerwig, G.J., see Kamerling, J.P. (222) 276
Gesztesi, A., see Sajgó, M. (222) 303

Gettings, S.D., Flanagan, R.J. and Holt, D.W.
Simple method for the measurement of tocainide and lignocaine in blood plasma or serum using GLC with flame ionisation detection (225) 469
Ghebregzabher, M., Rufini, S., Castellucci, M.G. and Lato, M.
Analysis of some tryptophan and phenylalanine metabolites in urine by a straight-phase HPLC technique (222) 191
Giacomini, K.M., see Swezey, S.F. (227) 510
Gianazza, E., see Cassara, G. (221) 279
Gidoh, M., Tsutsumi, S. and Takitani, S.
Determination of three main antileprosy drugs and their main metabolites in serum by HPLC (223) 379
Gilbert, T.W., see Zodda, J.P. (227) 249
Gill, R., Lopes, A.A.T. and Moffat, A.C.
Analysis of barbiturates in blood by HPLC (226) 117
Gillespie, T.J. and Sipes, I.G.
Sensitive GC determination of trifluoperazine in human plasma (223) 95
Gillet, G., Fraisse-André, J., Lee, C.R., Dring, L.G. and Morselli, P.L.
GC method for the determination of progabide (SL 76.002) in biological fluids (230) 154
Ginneken, C.A.M. van, see Van Ginneken, C.A.M. (229) 129
Girre, C., see Tournet, M.-C. (224) 348
Gitlow, S.E., see Krstulovic, A.M. (223) 305
Gitlow, S.E., see Krstulović, A.M. (227) 379
Gjessing, L.R., see Undrum, T. (227) 53
Glencross, R.G., Abeywardene, S.A., Corney, S.J. and Morris, H.S.
The use of oestradiol-17β antiserum covalently coupled to Sepharose to extract oestradiol-17β from biological fluids (223) 193
Gochin, R., Kanfer, I. and Haigh, J.M.
Simultaneous determination of trimethoprim, sulphamethoxazole and $N^4$-acetylsulphamethoxazole in serum and urine by HPLC (223) 139
Godbillon, J. and Carnis, G.
Determination of the major metabolite of phentolamine in human plasma and urine by HPLC (222) 461
Godbillon, J., Gauron, S. and Gosset, G.
HPLC analysis of the sulfide metabolite of sulfinpyrazone in plasma (227) 516
Godbillon, J., see Sioufi, A. (221) 419
Godse, D.D., see Warsh, J.J. (228) 131
Goebel, G., see Breithaupt, H. (226) 237
Goehl, T.J., see Rapaka, R.S. (227) 463
Goldberg, E.M., Blendis, L.M. and Sandler, S.
A GC-MS study of profiles of volatile metabolites in hepatic encephalopathy (226) 291
Gómez, C.J., see Bizzozero, O. (227) 33
Gonzalez, M.A., see Williams, C. II (225) 225
Gordon, G.R., see Peters, J.H. (229) 503
Gorman, C.A., see Hay, I.D. (226) 383
Gorodetzky, C.W., see Cone, E.J. (230) 57
Gorsen, R.M., Weiss, A.J. and Manthei, R.W.
Analysis of procarbazine and metabolites by GC-MS (221) 309
Gosset, G., see Godbillon, J. (227) 516
Goswami, S.K., see Mai, J. (230) 15
Goto, J., Kato, H., Saruta, Y. and Nambara, T.
Studies on steroids. CLXX. Separation and determination of bile acid 3-sulfates in human bile by HPLC (226) 13
Goto, M., Nakamura, T. and Ishii, D.
Micro HPLC system with micro precolumn and dual electrochemical detector for direct injection analysis of catecholamines in body fluids (226) 33
Grandjean, E.M., see Simona, M.G. (224) 532
Granneman, G.R. and Sennello, L.T.
A very precise HPLC procedure for the determination of cefmenoxime, a new cephalosporin antibiotic, in plasma (229) 149
Grassl, M., see Haegele, E.-O. (223) 69
Gray, A.J., see Bailey, E. (225) 83
Gray, M.J., see Silver, H.K.B. (224) 381
Grebow, P.E., see Choi, R.L. (230) 181
Grech-Bélanger, O., see Bélanger, P.M. (228) 327
Greenberg, R., see Haughey, D.B. (229) 387
Greenblatt, D.J., Divoll, M., Moschitto, L.J. and Shader, R.I.
Electron-capture GC analysis of the triazolobenzodiazepines alprazolam and triazolam (225) 202
Greenblatt, D.J., see Abernethy, D.R. (223) 432
Greenblatt, D.J., see Ameer, B. (226) 224
Greenblatt, D.J., see Divoll, M. (222) 125
Grewal, R.S., see Rajagopalan, T.G. (224) 265
Gribnau, F.W.J., see Kerremans, A.L.M. (229) 129
Griesmann, G.E., Chan, W.-Y. and Rennert, O.M.
Determination of γ-aminobutyric acid by reversed-phase HPLC and pre-column labeling for fluorescence detection (230) 121
Griffin, C.A., see Henderson, R.J. Jr. (226) 202
Griffin, J.P., see Allen, B.A. (222) 146
Gröningsson, K., see Vessman, J. (229) 227
Grossi, C., see Ching, N.P.H. (225) 196
Grossi, C.E., see Ching, N.P.H. (222) 171
Grove, J., Fozard, J.R. and Mamont, P.S.
Assay of α-difluoromethylornithine in body fluids and tissues by automatic amino-acid analysis (223) 409
Grupe, A. and Spiteller, G.
New polar acid metabolites in human urine (226) 301
Grupe, A. and Spiteller, G.
Unexpected metabolites produced from clomethiazole (230) 335
Guadagni, L., see Zecca, L. (230) 168
Guay, D.R.P., Bockbrader, H.N. and Matzke, G.R.
HPLC analysis of cimetidine in serum and urine (228) 398
Guerret, M.
Determination of pindolol in biological

fluids by an electron-capture GLC method on a wall-coated open tubular column (221) 387
Guinebault, P.R., Broquaire, M., Sanjuan, M., Rovei, V. and Braithwaite, R.A.
Determination of antrafenine and its main acid metabolite, 2-{[7-(trifluoromethyl)-4-quinolinyl]amino}-benzoic acid, in biological fluids using HPLC with large volume automatic injection and GLC with derivative formation (223) 103
Gundert-Remy, U., see Hildebrandt, R. (228) 409
Günthert, W., see De Vries, J.X. (221) 161
Gupta, R.N., Dunn, G.L., Morison, D.H. and Kumana, C.R.
Determination of minaxolone (CCI 12923) by GC with nitrogen-sensitive detection (224) 496
Gurusiddaiah, S. and Brosemer, R.W.
Separation of taurine and glycerophosphoryl ethanolamine on amino acid analyzers (223) 179
Gustafson, J.H., see Cotler, S. (222) 95
Guyon, F., Tambute, A., Caude, M. and Rosset, R.
Determination of N-methylpyridinium 2-aldoxime methylsulfate (Contrathion) in rat plasma and urine by copper(II)-silica ligand-exchange HPLC (229) 475

# H

Haanen, C., see Linssen, P. (223) 371
Hackett, L.P., see Anderson, R.D. (227) 239
Hackman, M.R. and Brooks, M.A.
Differential pulse amperometric detection of drugs in plasma using a dropping mercury electrode as a HPLC detector (222) 179
Hackman, M.R., see Brooks, M.A. (229) 167
Haefelfinger, P.
Determination of bufuralol and its major metabolites in plasma by HPLC (221) 327
Haeffner, E.W. and Hoffmann, C.J.K.
Direct quantitation of free cholesterol in total serum lipid extracts by computer-assisted GLC (228) 268
Haegele, E.-O., Schaich, E., Rauscher, E., Lehmann, P. and Grassl, M.
Action pattern of human pancreatic α-amylase on maltoheptaose, a substrate for determining α-amylase in serum (223) 69
Hagardorn, A.N., see Robert, T.A. (224) 116
Haigh, J.M., see Gochin, R. (223) 139
Hall, E.D., see McGinley, P.A. (230) 29
Halpern, B., see Sims, P. (222) 337
Hamaki, M., see Hayakawa, K. (229) 159
Hamilton, L.H., see Christman, N.T. (229) 259
Hammar, M., see Allenmark, S. (224) 399
Hamme, K.J., see Peters, J.H. (229) 503
Hammill, S.C., see Shand, D.G. (224) 343
Hancock, W.S., Harding, D.R.K. and Friedman, Z.
Rapid analysis of 3-methylhistidine in urine, plasma, muscle and amniotic fluid with a single HPLC system but with different ion-pairing reagents (228) 273
Hansen, C.S., see Chang, S.F. (226) 79
Hansen, S.H.
Assay of 5-aminosalicylate and its acetylated metabolite in biological fluids by HPLC on dynamically modified silica (226) 504
Hansen, S.H. and Døssing, M.
Determination of urinary hippuric acid and *o*-cresol, as indices of toluene exposure, by LC on dynamically modified silica (229) 141
Hansen, S.H., see Schultz, B. (228) 279
Hara, I., see Okazaki, M. (221) 257, (223) 285
Hara, S., see Yamada, K. (223) 176
Haraguchi, S.-I., Terasawa, M., Toshima, H., Matsumoto, I., Kuhara, T. and Shinka, T.
Analysis of organic acids in the hearts of patients with idiopathic cardiomyopathy by GC-MS (230) 7
Haraguchi, S.-I., Toshima, H., Matsumoto, I., Kuhara, T. and Shinka, T.
Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1
Harapat, S.R. and Kates, R.E.
HPLC analysis of propafenone in human plasma samples (230) 448
Harding, D.R.K., see Hancock, W.S. (228) 273
Harkness, R.A., see Simmonds, R.J. (226) 369
Harman, P.J., Blackman, G.L. and Phillipou, G.
HPLC determination of clomiphene using post-column on-line photolysis and fluorescence detection (225) 131
Harmsen, E., De Tombe, P.P. and De Jong, J.W.
Simultaneous determination of myocardial adenine nucleotides and creatine phosphate by HPLC (230) 131
Harmsen, E., see De Jong, J.W. (229) 498
Harold, S., see Briand, R.L. (223) 277
Hart, S.J., Tontodonati, R. and Calder, I.C.
Reversed-phase chromatography of urinary metabolites of paracetamol using ion suppression and ion pairing (225) 387
Hartman, C.A., Kucharczyk, N., Sofia, R.D. and Perhach, J.L. Jr.
Determination of methyclothiazide in human plasma by HPLC (226) 510
Hartman, C.A., see Valia, K.H. (221) 170
Hartvig, P., Fagerlund, C. and Emanuelsson, B.-M.
Extractive alkylation of probenecid in plasma and cerebrospinal fluid and determination by electron-capture GC (228) 340

Hartvig, P., see Floberg, S. (225) 73
Hashimoto, M., see Maeda, T. (223) 357
Hashino, M., see Iwai, M. (225) 275
Haughey, D.B., Greenberg, R., Schaal, S.F. and Lima, J.J.
LC determination of caffeine in biologic fluids (229) 387
Haugland, T., see Salvesen, B. (225) 463
Hausen, A., Fuchs, D., König, K. and Wachter, H.
Determination of neopterine in human urine by reversed-phase HPLC (227) 61
Hawgood, S., see Ward, L.C. (223) 417
Hay, I.D., Annesley, T.M., Jiang, N.S. and Gorman, C.A.
Simultaneous determination of D- and L-thyroxine in human serum by LC with electrochemical detection (226) 383
Hayakawa, K., Kitada, K., Hamaki, M. and Miyazaki, M.
HPLC determination of clioquinol and its conjugates in biological materials (229) 159
Hayashi, C., see Yamaguchi, Y. (224) 111, (230) 263
Hayashi, T., Todoriki, H. and Naruse, H.
HPLC determination of $\alpha$-keto acids (224) 197
Hedegaard, E. and Jensen, B.
Nano-scale densitometric quantitation of phospholipids (225) 450
Hegge, J.A.J., see De Jong, J.W. (229) 498
Hegstrand, L.R. and Eichelman, B.
Determination of rat brain tissue catecholamines using LC with electrochemical detection (222) 107
Heineman, W.R., see Zodda, J.P. (227) 249
Hekster, Y.A., see Vree, T.B. (224) 519
Helge, H., see Nau, H. (226) 69
Henderson, R.J. Jr. and Griffin, C.A.
Analysis of adenosine, inosine and hypoxanthine in suspensions of cardiac myocytes by HPLC (226) 202
Henning, H. von, see Von Henning, H. (223) 1
Henzel, D., see Meulemans, A. (226) 255
Hermansson, J. and Von Bahr, C.
Simultaneous determination of *d*- and *l*-propranolol in human plasma by HPLC (221) 109
Hermansson, J. and Von Bahr, C.
Determination of (*R*)- and (*S*)-alprenolol and (*R*)- and (*S*)-metoprolol as their diastereomeric derivatives in human plasma by reversed-phase LC (227) 113
Hermsdorf, D., see Riedel, E. (229) 417
Herzfeld, L., see Timm, U. (229) 111
Heusler, H., Epping, J., Heusler, S., Richter, E., Vermeulen, N.P.E. and Breimer, D.D.
Simultaneous determination of blood concentrations of methohexital and its hydroxy metabolite by GC and identification of 4'-hydroxymethohexital by combined GLC-MS (226) 403
Heusler, S., see Heusler, H. (226) 403
Heykants, J., see Woestenborghs, R. (224) 25, (224) 122
Hicks, R., see Ching, N.P.H. (225) 196
Hicks, R.G., see Ching, N.P.H. (222) 171
Higuchi, S. and Kawamura, S.
Specific determination of plasma nicardipine hydrochloride levels by GC-MS (223) 341
Higuchi, S., see Isomura, H. (224) 423
Higuchi, S., see Kamimura, H. (229) 337
Hijikata, J., see Murakami, K. (227) 103
Hikida, K., see Sadanaga, T. (223) 243
Hildebrandt, R. and Gundert-Remy, U.
Improved procedure for the determination of the ureidopenicillins azlocillin and mezlocillin in plasma by HPLC (228) 409
Hinderling, P.H., see Eichhorst, O. (224) 67
Hiraga, Y. and Kinoshita, T.
Post-column derivatization of guanidino compounds in HPLC using ninhydrin (226) 43
Hirai, K., see Konishi, M. (229) 355
Hiramatsu, K. and Arimori, S.
Rapid determination of lipids in healthy human lymphocytes (227) 423
Hiramatsu, K. and Arimori, S.
Lipids in human neutrophils determined by a microanalytical method (229) 455
Hirano, N., see Yuki, H. (221) 271
Hirata, M., see Noda, A. (230) 345
Hirata, T., Kai, M., Kohashi, K. and Ohkura, Y.
Determination of phenylpyruvic acid in urine and serum by HPLC with fluorescence detection (226) 25
Hirobe, M., see Matsumura, K. (222) 53, (230) 148
Hirose, S., see Murakami, K. (225) 329
Höckel, M., Dünges, W., Holzer, A., Brockerhoff, P. and Rathgen, G.H.
A microliter method for the GC determination of long-chain non-esterified fatty acids in human serum or plasma (221) 205
Hoff, W.G. van 't, see Van 't Hoff, W.G. (225) 236
Hoffmann, C.J.K., see Haeffner, E.W. (228) 268
Holder, G.M., see Boux, L.J. (227) 149
Holdiness, M.R., Israili, Z.H. and Justice, J.B.
GC-MS determination of ethambutol in human plasma (224) 415
Holdiness, M.R., Justice, J.B., Salamone, J.D. and Neill, D.B.
GC-MS determination of glutamic acid decarboxylase activity in subregions of rat brain (225) 283
Holdiness, M.R., see Capogrossi, M.C. (227) 168
Holdiness, M.R., see Saraswat, L.D. (222) 353
Holick, M.F., see Adams, J.S. (226) 198
Holland, M.L., see Rhodes, G. (228) 33
Holliday, S.B., see Hoskins, J.A. (230) 162
Holmén, A., see Kinberger, B. (224) 449, (229) 492
Holt, D.W., see Gettings, S.D. (225) 469
Holzer, A., see Höckel, M. (221) 205

Holzer, P., see Skofitsch, G. (226) 53
Homan, J.W., see Feenstra, M.G.P. (230) 271
Honda, A., see Honda, S. (226) 341
Honda, S., Suzuki, S., Kakehi, K., Honda, A. and Takai, T.
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341
Hoogewijs, G., Michotte, Y., Lambrecht, J. and Massart, D.L.
HPLC determination of papaverine in whole blood (226) 423
Hooper, W.D., Kunze, H.E. and Eadie, M.J.
Simultaneous assay of methylphenobarbital and phenobarbital in plasma using GC-MS with selected ion monitoring (223) 426
Horáková, M., see Deyl, Z. (230) 409
Horikawa, R., see Uchikura, K. (223) 41
Horino, M., see Kakita, K. (222) 33
Horn, A.S., see Feenstra, M.G.P. (230) 271
Hoskins, J.A. and Holliday, S.B.
Determination of α-methyldopa and methyldopate in human breast milk and plasma by ion-exchange chromatography using electrochemical detection (230) 162
Houck, G.P., see Caudill, W.L. (227) 331
Houin, G., Bree, F. and Tillement, J.P.
Quantitative determination of niflumic acid and its β-morpholinoethyl ester in human plasma by GLC (223) 351
Houston, J.B., see Taylor, G. (230) 194
Howell, S.K. and Wang, Y.-M.
Quantitation of physiological α-tocopherol, metabolites, and related compounds by reversed-phase HPLC (227) 174
Hsieh, J.Y.-K. and Davis, K.L.
Rapid LC determination of probenecid in human cerebrospinal fluid (225) 521
Hsu, K.-Y., see Noda, A. (230) 345
Hsu, T.-F., see Ahmed, A.E. (222) 453
Huang, C.S., see Williams, C. II (225) 225
Huber, G.A., see Wermeille, M.M. (228) 187
Hufford, C.D., see Baker, J.K. (230) 69
Huggins, G.R., see Kostelc, J.G. (226) 315
Hui, K.-S., Hui, M., Cheng, K.-P., Lajtha, A., Boksay, I. and Fencik, M.
Determination of the melanotropin-inhibiting factor analogue pareptide in urine by HPLC (222) 512
Hui, M., see Hui, K.-S. (222) 512
Hulshoff, A., see Barends, D.M. (222) 316
Hulshoff, A., see Barends, DM. (225) 417
Hulshoff, A., see Sorel, R.H.A. (221) 129, (222) 241
Hundt, H.K.L. and Barlow, E.C.
TLC method for the quantitative analysis of nalidixic acid in human plasma (223) 165
Hundt, H.K.L. and Brown, L.W.
Determination of 6,11-dihydro-11-dibenz[*b,e*]oxepin-2-acetic acid (isoxepac) in plasma by HPLC (225) 482
Huntley, T.E., see Choi, R.L. (230) 181
Huth, J.A., see Danielson, N.D. (221) 39
Hwang, P.T.R., see Meyer, M.C. (223) 466

# I

Iams, J.D., see Ng, K.J. (228) 43
Ichikawa, Y., see Takada, K. (222) 156
Idzu, G., Ishibashi, M., Miyazaki, H. and Yamamoto, K.
Determination of glyceryl trinitrate in human plasma by GC-negative ion chemical ionization-selected ion monitoring (229) 327
Iguchi, S., see Noda, A. (230) 345
Ikenaka, T., see Omichi, K. (230) 415
Ilett, K.F., see Anderson, R.D. (227) 239
Imai, Y., see Shimada, K. (227) 445
Imamura, T., see Sugihara, J. (229) 193
Imamura, Y., see Nakajima, O. (225) 91
Imoto, T., see Sugihara, J. (229) 193
Inturrisi, C.E., see Kiang, C.-H. (222) 81
Ireland, C.M., see Boux, L.J. (227) 149
Ishibashi, M., see Idzu, G. (229) 327
Ishida, K., see Nimura, N. (221) 249
Ishiguro, M., see Okamoto, M. (223) 473
Ishii, D., see Goto, M. (226) 33
Ishikawa, K. and McGaugh, J.L.
Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35
Ishimoto, M., see Kaji, H. (221) 145
Isoda, T., see Nakajima, O. (225) 91
Isomura, H., Higuchi, S. and Kawamura, S.
GC-MS determination of plasma 5-fluorouracil after administration of 1-hexylcarbamoyl-5-fluorouracil to dogs and humans (224) 423
Israili, Z.H., see Capogrossi, M.C. (227) 168
Israili, Z.H., see Holdiness, M.R. (224) 415
Iwai, M., Kanno, H., Hashino, M., Suzuki, J., Yanaihara, T., Nakayama, T. and Mori, H.
Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

# J

Jack, D.B. and Willis, J.V.
Comments to the article "Improved GC method of determining diclofenac in plasma" (223) 484
Jacob, P. III, Wilson, M. and Benowitz, N.L.
Improved GC method for the determination of nicotine and cotinine in biologic fluids (222) 61

Jacob, R. and Barrett, E.
Chromatographic analysis of glutamine in plasma (229) 188
Jähnchen, E., see Wiegand, U.-W. (223) 238
Jakobs, C., see Luley, C. (224) 500
Jakobs, C., see Nau, H. (226) 69
Jakobs, U., see Krause, W. (230) 37
Jakobsen, P. and Pedersen, A.K.
Simultaneous determination of sulfinpyrazone and four of its metabolites by HPLC (223) 460
Jannuzzo, M.G., see Moro, E. (230) 207
Jardine, I., see Strife, R.J. (224) 168
Jarreau, C., see Leroyer, R. (228) 366
Jatlow, P.I., Miller, R. and Swigar, M.
Measurement of haloperidol in human plasma using reversed-phase HPLC (227) 233
Jauch, R., see Roth, W. (222) 13
Jenner, D.A., Brown, M.J. and Lhoste, F.J.M.
Determination of α-methyldopa, α-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507
Jenner, P.J. and Ellard, G.A.
HPLC determination of ethionamide and prothionamide in body fluids (225) 245
Jensen, B., see Hedegaard, E. (225) 450
Jham, G.N., see Ching, N.P.H. (222) 171, (225) 196
Jiang, N.S., see Hay, I.D. (226) 383
Jindal, S.P., Lutz, T. and Cooper, T.B.
Determination of phenelzine in human plasma with GC-MS using an isotope labeled internal standard (221) 301
Jochemsen, R. and Breimer, D.D.
Assay of triazolam in plasma by capillary GC (223) 438
Jochemsen, R. and Breimer, D.D.
Assay of flunitrazepam, temazepam and desalkylflurazepam in plasma by capillary GC with electron-capture detection (227) 199
Johansson, B. and Nilsson, B.
Separation and quantitation of urinary porphyrins by HPLC (229) 439
Johansson, M., see Vessman, J. (229) 227
Johnson, D.A., see Kunitani, M.G. (224) 156
Johnson, E.A.
Gel chromatography of heparin (230) 216
Johnson, T.E., see DeAngelis, R.L. (221) 353
Jones, A.B., Elsohly, M.A., Bedford, J.A. and Turner, C.E.
Determination of cannabidiol in plasma by electron-capture GC (226) 99
Jones, C.R., Ryle, P.R. and Weatherley, B.C.
Measurement of pyrimethamine in human plasma by GLC (224) 492
Jones, C.R., see Egan, C.M. (224) 338
Jones, D.P.
Determination of pyridine dinucleotides in cell extracts by HPLC (225) 446
Jones, G.
Ternary solvent mixtures for improved resolution of hydroxylated metabolites of vitamin $D_2$ and vitamin $D_3$ during HPLC (221) 27
Jones, G.J.L., see Kuksis, A. (224) 1
Jones, R.M., see Buskin, J.N. (230) 438
Jones, S.P., see Lea, E.J.A. (221) 367
Jong, J.W. de, see De Jong, J.W. (229) 498, (230) 131
Jonkman, J.H.G., see Muir, K.T. (221) 85
Joseph, M.H., Kadam, B.V. and Risby, D.
Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361
Joubert, S.M., see Bredenkamp, B.L.F. (230) 327
Jozefczak, C., Ktorza, N. and Uzan, A.
HPLC determination of indalpine, a new non-tricyclic antidepressant, in human plasma. Identification and simultaneous measurement of its major plasma metabolite (230) 87
Juge, F., see Lecaillon, J.B. (228) 257
Julian, B., see Lefebvre, M.A. (230) 199
Jusko, W.J., see Pautler, D.B. (228) 215
Jusko, W.J., see Rocci, M.L. Jr. (224) 221
Justice, J.B., see Holdiness, M.R. (224) 415, (225) 283
Justice, J.B., see Saraswat, L.D. (222) 353
Justová, V., Wildtová, Z. and Stárka, L.
Separation of nanogram quantities of hydroxy metabolites of vitamin $D_3$ in plasma by TLC on silica gel (230) 319

# K

Kaa, E.
Rapid GC method for emergency determination of paracetamol in human serum (221) 414
Kabasawa, Y., see Uchikura, K. (223) 41
Kadam, B.V., see Joseph, M.H. (226) 361
Kågedal, B. and Källberg, M.
Reversed-phase ion-pair HPLC of mercaptoacetate and N-acetylcysteine after derivatization with N-(1-pyrene)maleimide and N-(7-dimethylamino-4-methyl-3-coumarinyl)maleimide (229) 409
Kai, M., see Hirata, T. (226) 25
Kaji, H., Saito, N., Murao, M., Ishimoto, M., Kondo, H., Gasa, S. and Saito, K.
GC and GC-MS studies on α-keto-γ-methylthiobutyric acid in urine following ingestion of optical isomers of methionine (221) 145

Kakehi, K., see Honda, S. (226) 341
Kakita, K., Horino, M., Tenku, A., Nishida, S., Matsumura, S., Matsuki, M. and Kakita, S.
Gel chromatographic separation of human C-peptide and proinsulin (222) 33
Kakita, S., see Kakita, K. (222) 33
Källberg, M., see Kågedal, B. (229) 409
Kamerling, J.P., Duran, M., Gerwig, G.J., Ketting, D., Bruinvis, L., Vliegenthart, J.F.G. and Wadman, S.K.
Determination of the absolute configuration of some biologically important urinary 2-hydroxydicarboxylic acids by capillary GLC (222) 276
Kamimura, H., Sasaki, H., Higuchi, S. and Shiobara, Y.
Quantitative determination of the β-adrenoceptor stimulant formoterol in urine by GC-MS (229) 337
Kamimura, H., Sasaki, H. and Kawamura, S.
Determination of the α,β-adrenoceptor blocker YM-09538 in plasma by HPLC with fluorescence detection (225) 115
Kanfer, I., see Gochin, R. (223) 139
Kang-Lee, Y.A., see McKee, R.W. (230) 309
Kanno, H., see Iwai, M. (225) 275
Karim, K.A., see Silver, H.K.B. (224) 381
Kärkkäinen, S. and Seppälä, E.
GC analysis of therapeutic concentrations of maprotiline in serum, using flame-ionization detection (221) 319
Karmen, A., see Smith, S.L. (223) 173
Karsai, T., see Bardócz, S. (223) 198
Kasper, M.E., see Lauff, J.J. (226) 391
Kasper, W., see Wiegand, U.-W. (223) 238
Kates, R.E., see Harapat, S.R. (230) 448
Kates, R.E., see Yee, Y.-G. (223) 454
Kato, H., see Goto, J. (226) 13
Kato, H., see Matsumura, K. (222) 53, (230) 148
Kato, T., see Rahman, M.K. (221) 265
Katogi, Y., Tamaki, N., Adachi, M., Terao, J. and Mitomi, M.
Simultaneous determination of dantrolene and its metabolite, 5-hydroxydantrolene, in human plasma by HPLC (228) 404
Kauffman, R.E., see Aravind, M.K. (221) 176, (228) 418
Kauffman, R.E., see Bowman, D.B. (229) 487
Kawaguchi, S., see Maeda, K. (221) 199
Kawahara, K., Matsumura, M. and Kimura, K.
Determination of flurbiprofen in human plasma using GC-MS with selected ion monitoring (223) 202
Kawai, S., see Koh, S. (226) 461
Kawai, S., see Uji, A. (221) 155
Kawamura, M., see Suzuki, M. (221) 361
Kawamura, S., see Higuchi, S. (223) 341
Kawamura, S., see Isomura, H. (224) 423
Kawamura, S., see Kamimura, H. (225) 115
Kawasaki, H., see Yuki, H. (221) 271
Kawasaki, T., Maeda, M. and Tsuji, A.
Determination of 17-oxosteroids in serum and urine by fluorescence HPLC using dansyl hydrazine as a pre-labeling reagent (226) 1
Kay, D.R., see Bergstrom, R.F. (222) 445
Kayama, E., see Yamada, K. (223) 176
Kearns, G.L. and Wilson, J.T.
Determination of ibuprofen in serum by HPLC and application to ibuprofen disposition (226) 183
Kegel, H., see Luley, C. (224) 500
Kerremans, A.L.M., Tan, Y., Van Ginneken, C.A.M. and Gribnau, F.W.J.
Specimen handling and HPLC determination of furosemide (229) 129
Kersten, H., see Mach, M. (223) 51
Kersten, W., see Mach, M. (223) 51
Ketelaars, H.C.J. and Peters, J.G.P.
Determination of guaiphenesin and its metabolite, β-(2-methoxyphenoxy)lactic acid, in plasma by HPLC (224) 144
Ketting, D., see Kamerling, J.P. (222) 276
Kiang, C.-H., Campos-Flor, S. and Inturrisi, C.E.
Determination of acetylmethadol and metabolites by use of HPLC (222) 81
Kientz, C.E., see Verweij, A. (226) 165
Kilts, C.D., Breese, G.R. and Mailman, R.B.
Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347
Kimura, K., see Kawahara, K. (223) 202
Kimura, Y., see Konaka, R. (225) 169
Kinberger, B. and Holmén, A.
Simultaneous determination of acetaminophen, theophylline and salicylate in serum by HPLC (229) 492
Kinberger, B., Holmén, A. and Wahrgren, P.
Determination of barbiturates and some neutral drugs in serum using quartz glass capillary GC (224) 449
King, I.S., see Balali-Mood, M. (229) 234
King, T.B., see Sternson, L.A. (227) 223
Kinney, C.D.
Determination of metoprolol in plasma and urine by GLC with electron-capture detection (225) 213
Kinoshita, T., see Hiraga, Y. (226) 43
Kinoshita, T., see Kubo, H. (227) 244
Kinoshita, T., see Nimura, N. (221) 249
Kinsella, J.E., see Mai, J. (230) 15
Kirkiacharian, S., see Cisse, H. (225) 509
Kitada, K., see Hayakawa, K. (229) 159
Kitchens, R.L., see Strauss, J.F. (221) 139
Klebovich, I. and Vereczkey, L.
GC method for the determination of flumecinol in biological fluids (221) 403
Kleijn, E. van der, see Van der Kleijn, E. (224) 519
Klotz, U.
Determination of bromazepam by GLC and its application for pharmacokinetic studies in man (222) 501
Klotz, U., see Fischer, C. (225) 498
Knapp, D.R., see Kuhnert, B.R. (224) 488
Knapp, G., see Lankmayr, E.P. (224) 239
Knödgen, B., see Seiler, N. (221) 227

Kobayashi, K., see Maeda, K. (221) 199
Kobayashi, Y., see Kubo, H. (227) 244
Koblantz, C., see Park, G.B. (227) 534
Koh, S., Arai, M., Kawai, S. and Okamoto, M.
Assay of catechol O-methyltransferase activity by HPLC with electrochemical detection (226) 461
Kohashi, K., see Hirata, T. (226) 25
Kohashi, K., see Tsuruta, Y. (224) 105
Kok, R.M. and Faber, D.B.
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389
Konaka, R., Kuruma, K., Nishimura, R., Kimura, Y. and Yoshida, T.
HPLC analysis of a new β-lactam antibiotic, 6059-S (moxalactam) (225) 169
Kondo, H., see Kaji, H. (221) 145
Konecki, J., see Gabryś, J. (222) 345
König, K., see Hausen, A. (227) 61
Konings, K.A.G., see Benschop, H.P. (225) 107
Konishi, M., Mori, Y. and Hirai, K.
Simultaneous determination of triazolobenzophenone [2',5-dichloro-2-(3-glycylaminomethyl-5-methyl-4H-1,2,4-triazol-4-yl)-benzophenone] and its major blood metabolite, triazolam, in monkey plasma by electron-capture GLC (229) 355
Kornhuber, H.H., see Frank, H. (224) 177
Korst, J.J., see Dokladalova, J. (224) 33
Koss, F.W., see Roth, W. (222) 13
Koss, F.-W., see Schmid, J. (227) 71
Kossen, S.P., see Benschop, H.P. (225) 107
Kostelc, J.G., Preti, G., Zelson, P.R., Tonzetich, J. and Huggins, G.R.
Volatiles of exogenous origin from the human oral cavity (226) 315
Kostenbauder, H.B., see Chambers, D.M. (225) 231
Kou, A.Y., see Chen, S.S.-H. (227) 25
Koyama, T., see Miyazaki, K. (223) 449
Koyama, Y., see Nakajima, O. (225) 91, (229) 481
Kramer, P.A., see Masso, P.D. (224) 457
Kramer, P.A., see Wong, S.H.Y. (226) 147
Krause, W.
Determination of plasma and urine levels of a new anti-inflammatory agent, 4,5-bis-(4-methoxyphenyl)-2-(2-hydroxymethylsulfinyl)-imidazole, by HPLC and electrochemical or UV detection (222) 71
Krause, W. and Jakobs, U.
Determination of plasma levels of spironerone, a new aldosterone antagonist, and one of its metabolites by HPLC (230) 37
Kreutz, G., see Riedel, E. (229) 417
Kreutzmann, D.J., Silink, M. and Cam, G.R.
Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17α-hydroxyprogesterone from small plasma or tissue samples (228) 95
Kroviarski, Y., Cochet, S., Boivin, P. and Bertrand, O.
Rapid separation of human globin chains by high flow-rate chromatography under low pressure using CM-Trisacryl M (228) 298
Krstulović, A.M.
Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1
Krstulović, A.M., Bertani-Dziedzic, L., Bautista-Cerqueira, S. and Gitlow, S.E.
Simultaneous determination of 4-hydroxy-3-methoxy-phenylacetic (homovanillic) acid and other monoamine metabolites in human lumbar cerebrospinal fluid. An improved HPLC study with electrochemical detection (227) 379
Krstulovic, A.M., Bertani-Dziedzic, L., Dziedzic, S.W. and Gitlow, S.E.
Quantitative determination of 3-methoxy-4-hydroxyphenylethyleneglycol and its sulfate conjugate in human lumbar cerebrospinal fluid using LC with aperometric detection (223) 305
Ktorza, N., see Jozefczak, C. (230) 87
Kubo, H., Kinoshita, T., Kobayashi, Y. and Tokunaga, K.
Micro determination of gentamicin in serum by HPLC (227) 244
Kubo, O., see Matsumura, K. (222) 53, (230) 148
Kubota, T., see Nakajima, O. (229) 481
Kucharczyk, N., see Hartman, C.A. (226) 510
Kucharczyk, N., see Valia, K.H. (221) 170
Kuhara, T., see Haraguchi, S.-I. (227) 1, (230) 7
Kuhnert, B.R., Kuhnert, P.M., Reese, A.L.P. and Knapp, D.R.
Measurement of 2-chloroprocaine in plasma by selected ion monitoring (224) 488
Kuhnert, P.M., see Kuhnert, B.R. (224) 488
Kuksis, A., Myher, J.J., Geher, K., Breckenridge, W.C., Jones, G.J.L. and Little, J.A.
Lipid class and molecular species interrelationships among plasma lipoproteins of normolipemic subjects (224) 1
Kuksis, A., Myher, J.J., Geher, K., Breckenridge, W.C. and Little, J.A.
Lipid class and molecular species interrelationships among plasma lipoproteins of Type III and Type IV hyperlipemic subjects (230) 231
Kullberg, M.P., see Park, G.B. (222) 213
Kumana, C.R., see Gupta, R.N. (224) 496
Kundu, S.K. and Suzuki, A.
Simple micro-method for the isolation of gangliosides by reversed-phase chromatography (224) 249
Kunitani, M., see Muir, K.T. (221) 85
Kunitani, M.G., Johnson, D.A., Upton, R.A. and Riegelman, S.
Convenient and sensitive HPLC assay for cimetidine in plasma or urine (224) 156
Kunze, H.E., see Hooper, W.D. (223) 426
Kuo, J.C. and Yeung, E.S.
Determination of carbohydrates in urine by HPLC and optical activity detection (223) 321

Kuo, J.C. and Yeung, E.S.
Determination of free and esterified cholesterol in human serum by HPLC and optical activity detection (229) 293
Kuo, K.C., see Gehrke, C.W. (230) 297
Kuruma, K., see Konaka, R. (225) 169
Kutt, H., see Dhar, A.K. (222) 203
Kuttan, R., Wilson, N. and Tenni, R.
Determination of $\gamma$-carboxyglutamic acid excretion in urine (223) 182
Kuwada, M., Tateyama, T. and Tsutsumi, J.
Simultaneous determination of verapamil and its seven metabolites by HPLC (222) 507
Kylberg-Hanssen, K., see Ervik, M. (229) 87

# L

Lacotte, J., see Moulin, M.A. (226) 250
Lagerström, P.-O.
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476
Lagerström, P.-O., see Ervik, M. (229) 87
Lahr, P.D., see Mais, D.E. (225) 27
Lai, C.-C., see Lin, J.-K. (227) 369
Lajtha, A., see Hui, K.-S. (222) 512
Laker, M.F., see Mount, J.N. (226) 191
Lambrecht, J., see Hoogewijs, G. (226) 423
Land, G. and Bye, A.
Simple HPLC method for the analysis of 9-(2-hydroxyethoxymethyl)guanine (acyclovir) in human plasma and urine (224) 51
Land, G., Dean, K. and Bye, A.
Determination of cyclizine and norcyclizine in plasma and urine using GLC with nitrogen selective detection (222) 135
Land, G., see Dean, K. (221) 408
Land, G., see Lo, L.Y. (222) 297
Lang, E., see Buskin, J.N. (230) 454
Lankmayr, E.P., Budna, K.W., Müller, K., Nachtmann, F. and Rainer, F.
Determination of D-penicillamine in serum by fluorescence derivatization and LCC (222) 249
Lankmayr, E.P., Maichin, B., Knapp, G. and Nachtmann, F.
Catalytic detection principle for HPLC: determination of enantiomeric iodinated thyronines in blood serum (224) 239
Lapierre, Y.D., see Cooper, S.F. (222) 291
Larking, P.
GC determination of clonazepam (221) 399
Larsen, H.F. and Trostmann, A.F.
Improved TLC assay for monitoring lecithin/sphingomyelin ratios in amniotic fluid (226) 484
Larsen, N.-E. and Marinelli, K.
Mass fragmentographic quantification of naproxen in human plasma (222) 482
Larsen, N.-E., Marinelli, K. and Møller Heilesen, A.
Determination of gentamicin in serum using LCC (221) 182
Lartigue-Mattei, C., Chabard, J.L., Bargnoux, H., Petit, J. and Berger, J.A.
Simultaneous determination of allopurinol and oxipurinol in biological fluids by mass fragmentography (229) 211
Lartigue-Mattei, C., see Chabard, J.L. (221) 9
Lato, M., see Ghebregzabher, M. (222) 191
Lauff, J.J., Kasper, M.E. and Ambrose, R.T.
Separation of bilirubin species in serum and bile by reversed-phase HPLC (226) 391
Lauterburg, B.H., Smith, C.V. and Mitchell, J.R.
Determination of isoniazid and its hydrazino metabolites, acetylisoniazid, acetylhydrazine, and diacetylhydrazine in human plasma by GC-MS (224) 431
Lawson, G.J., Dixon, P.F. and Aherne, G.W.
Rapid and simple method for the measurement of methotrexate and 7-hydroxymethotrexate in serum by HPLC (223) 225
Le Roux, Y., Gaillot, J. and Bieder, A.
HPLC determination of pipotiazine in human plasma and urine (230) 401
Lea, E.J.A. and Jones, S.P.
Simultaneous measurement of sterol and fatty acid composition in small samples (221) 367
Lecaillon, J.B., Rouan, M.C., Souppart, C., Febvre, N. and Juge, F.
Determination of cefsulodin, cefotiam, cephalexin, cefotaxime, desacetyl-cefotaxime, cefuroxime and cefroxadin in plasma and urine by HPLC (228) 257
Lee, C.R., see Gillet, G. (230) 154
Lee, G.J.-L. and Tieckelmann, H.
The application of HPLC in enzymatic assays of chondroitin sulfate isomers in normal human urine (222) 23
Lee, T.L. and Brooks, M.A.
Determination of amdinocillin in plasma and urine by high-performance liquid chromatography (227) 137
Lee, T.L., Brooks, M.A. and De Silva, J.A.F.
Determination of the antihypertensive agent 1-(2-aminoethyl)-3-(2,6-dichlorophenyl)thiourea in plasma and urine by HPLC (224) 275
Lee, W.H., see Pieroni, J.P. (230) 115
Lefebvre, M.A., Julian, B. and Fourtillan, J.B.
HPLC assay for determination of a new β-blocking agent FM 24 (230) 199
Leferink, J.G., Dankers, J. and Maes, R.A.A.
A time-saving method for the determination of the $\beta_2$ sympathomimetics terbutaline, salbutamol and fenoterol. Preliminary results (229) 217

LeGatt, D.F., Baker, G.B. and Coutts, R.T.
Simultaneous extraction and separation of trace amines of biological interest (225) 301
Lehmann, P., see Haegele, E.-O. (223) 69
Lemar, M., see Sommadossi, J.P. (228) 205
Lembeck, F., see Skofitsch, G. (226) 53
Lemke, P.R., see Dokladalova, J. (224) 33
Lemmer, B., see Winkler, H. (228) 223
Leroux, J.P., see Rocchiccioli, F. (226) 325
Leroyer, R., Jarreau, C. and Pays, M.
Specific determination of quinidine and metabolites in biological fluids by reversed-phase HPLC (228) 366
Leszczyńska, D., see Schneider, Z. (229) 77
Lewis, E.C., see Chester, T.L. (225) 17
Lhoste, F.J.M., see Jenner, D.A. (224) 507
Li, J.B., see Wassner, S.J. (227) 497
Liddiard, C. and Nau, H.
Determination of etozolin and ozolinone in human plasma and tissues by reversed-phase HPLC (225) 504
Ligtenstein, D.A., see Benschop, H.P. (225) 107
Lim, C.K. and Chan, J.Y.Y.
Normal-phase HPLC of porphyrin free acids on silica modified with tetraethylenepentamine (228) 305
Lim, C.K., see Crowne, H. (223) 421
Lima, J.J., see Haughey, D.B. (229) 387
Lin, J.-K. and Lai, C.-C.
Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369
Lin, K.-T.D.
Simplified procedure for the analysis of 3- and 4-hydroxyproline (227) 341
Lindström, B., see Alván, G. (229) 241
Lindström, B., see Floberg, S. (221) 166, (225) 73
Lindström, E., see Allenmark, S. (224) 399
Linssen, P., Drenthe-Schonk, A., Wessels, H. and Haanen, C.
Determination of 1-β-D-arabinofuranosyl-cytosine and 1-β-D-arabinofuranosyluracil in human plasma by HPLC (223) 371
Lipe, G.W., see Slikker, W. Jr. (224) 205
Little, G.H., see Cole, K.D. (227) 503
Little, J.A., see Kuksis, A. (224) 1, (230) 231
Lo, L.Y., Land, G. and Bye, A.
Sensitive assay for pseudoephedrine and its metabolite, norpseudoephedrine in plasma and urine using GLC with electron-capture detection (222) 297
Longo, A., Di Toro, M., Pagani, E. and Carenzi, A.
Simple selected ion monitoring method for determination of fosfomycin in blood and urine (224) 257
Lönnerholm, G., see Floberg, S. (221) 166, (225) 73
Loo, T.L., see Benvenuto, J.A. (222) 518
Loo, T.L., see Miller, A.A. (228) 165
Loomis, C.W. and Brien, J.F.
Determination of carbimide in plasma by GLC (222) 421
Loon, G.R. van, see Van Loon, G.R. (228) 123
Lopes, A.A.T., see Gill, R. (226) 117
Losito, R., Gattiker, H. and Bilodeau, G.
Gel chromatography of heparin (226) 61
Lovelace, C.E.A., Njapau, H., Salter, L.F. and Bayley, A.C.
Screening method for the detection of aflatoxin and metabolites in human urine: aflatoxins $B_1$, $G_1$, $M_1$, $B_2a$, $G_2a$, aflatoxicols I and II (227) 256
Lowe, J.R., see Pickup, M.E. (225) 493
Lowenthal, A., see Marescau, B. (224) 185
Luccioni, C.M., see Au, J.L.-S. (228) 245
Ludden, T.M., see Velagapudi, R. (228) 423
Luley, C., Kegel, H. and Jakobs, C.
Improved method for the determination of procetofenic acid in human plasma by GLC (224) 500
Lunan, K.D., see Thomas, D.W. (225) 433
Lunde, H., see Undrum, T. (227) 53
Lutz, T., see Jindal, S.P. (221) 301
Luyten, Y., see Marescau, B. (224) 185

# M

Mabuchi, H. and Nakahashi, H.
Analysis of middle molecular peptides in normal and uremic body fluids by HPLC (224) 322
Mabuchi, H. and Nakahashi, H.
Analysis of small peptides in uremic serum by HPLC (228) 292
McCarthy, T.P., Brodie, B., Milner, J.A. and Bevill, R.F.
Improved method for selenium determination in biological samples by GC (225) 9
McCauley, T., see Wong, S.H.Y. (226) 147
McChesney, J.D., see Baker, J.K. (230) 69
McCloskey, J.A., see Gelijkens, C.F. (225) 291
McCluskey, M., see Egan, C.M. (224) 338
McCormack, J.J., see Ackerly, C.C. (230) 175
McCormack, J.J., see Allen, B.A. (222) 146
McCullough, J., see Rao, G.H.R. (229) 205
McCune, R.A., see Gehrke, C.W. (230) 297
McDermott, J.R., Smith, A.I., Biggins, J.A., Al-Noaemi, M.C. and Edwardson, J.A.
Characterization and determination of neuropeptides by HPLC and radioimmunoassay (222) 371
Macek, K., Deyl, Z., Čoupek, J. and Sanitrák, J.
Separation of collagen types I and III by HPLC (222) 284
Macek, K., see Deyl, Z. (230) 409
McErlane, K.M., see Pillai, G.K. (229) 103
McGaugh, J.L., see Ishikawa, K. (229) 35
McGilveray, I.J., see Charette, C. (224) 128

McGinley, P.A., Braughler, J.M. and Hall, E.D.
Determination of methylprednisolone in central nervous tissue and plasma using normal-phase HPLC (230) 29
Mach, M., Kersten, H. and Kersten, W.
Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51
McKee, R.W., Kang-Lee, Y.A., Panaqua, M. and Swendseid, M.E.
Determination of nicotinamide and metabolic products in urine by HPLC (230) 309
McLean, A.E.M., see Beales, D. (226) 498
McLean, S., see Roberts, M.S. (226) 175
McNair, H.M., see Tangney, C.C. (224) 389
Maddocks, J.L., see Dooley, T. (229) 121
Maeda, K., Kawaguchi, S., Niwa, T., Ohki, T. and Kobayashi, K.
Identification of some abnormal metabolites in psoriatic nail using GC-MS (221) 199
Maeda, K., see Niwa, T. (225) 1, (228) 59, (230) 1
Maeda, K., see Ohki, T. (228) 51
Maeda, M., see Kawasaki, T. (226) 1
Maeda, T., Yamaguchi, T. and Hashimoto, M.
GC determination of the hypoglycaemic agent gliclazide in plasma (223) 357
Maes, R.A.A., see Leferink, J.G. (229) 217
Maes, R.A.A., see Meering, P.G. (225) 407
Magnusson, J.-O., see Eriksson, B.-M. (223) 401
Magnusson, O., Nilsson, L.B. and Westerlund, D.
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237
Maguire, K.P., Norman, T.R., Burrows, G.D. and Scoggins, B.A.
Simultaneous measurement of dothiepin and its major metabolites in plasma and whole blood by GC-MF (222) 399
Mai, J., Goswami, S.K., Bruckner, G. and Kinsella, J.E.
Determination of prostaglandins and thromboxane as their pentafluorobenzyl-trimethylsilyl derivatives by electron-capture GC (230) 15
Maichin, B., see Lankmayr, E.P. (224) 239
Maier, K., see Fischer, C. (225) 498
Mailman, R.B., see Kilts, C.D. (225) 347
Mais, D.E., Lahr, P.D. and Bosin, T.R.
Determination of serotonin, its precursors, metabolites and [$^3$H]serotonin in lung by HPLC with fluorescence detection (225) 27
Makin, H.L.J., see Trafford, D.J.H. (226) 351
Mamont, P., see Wagner, J. (227) 349
Mamont, P.S., see Grove, J. (223) 409
Manabe, T., Okuyama, T., Suzuki, A. and Shigematsu, A.
Detection of the changes in protein distribution of rat plasma induced by carbon tetrachloride administration by means of two-dimensional electrophoresis (225) 65
Mangoni, P., see Sioufi, A. (222) 429
Manthei, R.W., see Gorsen, R.M. (221) 309
Maquart, F.X., see Bellon, G. (230) 420
Marchesini, B., see Boschi, S. (224) 139
Marcucci, F., see Cotellessa, L. (221) 149
Marcucci, F., see Mussini, E. (224) 94
Marescau, B., Lowenthal, A., Esmans, E., Luyten, Y., Alderweireldt, F. and Terheggen, H.G.
Isolation and identification of some guanidino compounds in the urine of patients with hyperargininaemia by LC, TLC and GC-MS (224) 185
Marinelli, K., see Larsen, N.-E. (221) 182, (222) 482
Martin, B., see Caron, J.C. (230) 125
Martin, L.E., see Carey, P.F. (225) 161
Marunaka, T. and Umeno, Y.
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382
Marunaka, T., see Matsushima, E. (230) 188
Marunaka, T., see Umeno, Y. (226) 333
Marzillo, G., see Farina, P. (222) 141
Mascher, H., see Nitsche, V. (227) 521
Massart, D.L., see Hoogewijs, G. (226) 423
Masso, P.D. and Kramer, P.A.
Simultaneous determination of disulfiram and two of its dithiocarbamate metabolites in human plasma by reversed-phase LC (224) 457
Matsuki, M., see Kakita, K. (222) 33
Matsumoto, I., see Haraguchi, S.-I. (227) 1, (230) 7
Matsumura, K., Kubo, O., Sakashita, T., Adachi, Y., Kato, H., Watanabe, K. and Hirobe, M.
Quantitative determination of tulobuterol and its metabolites in human urine by mass fragmentography (222) 53
Matsumura, K., Kubo, O., Tsukada, T., Nishide, K., Kato, H., Watanabe, K. and Hirobe, M.
Determination of tulobuterol in human serum by GLC (230) 148
Matsumura, M., see Kawahara, K. (223) 202
Matsumura, S., see Kakita, K. (222) 33
Matsushima, E., Umeno, Y., Marunaka, T. and Akagi, H.
Determination of pentazirinocyclodiphosphathiazene in biological fluids by HPLC (230) 188
Matsushima, E., see Umeno, Y. (226) 333
Matsushima, Y., see Sadanaga, T. (223) 243
Matsuyama, K., see Noda, A. (230) 345
Matthaei, D., see Pinkston, D. (223) 1
Mattocks, A.M., see Wesley-Hadzija, B. (229) 425
Matzke, G.R., see Guay, D.R.P. (228) 398
Maume, B.F., see Ramirez, L.C. (229) 267
Maurer, H. and Pfleger, K.
Determination of 1,4- and 1,5-

benzodiazepines in urine using a computerized GC-MS technique (222) 409
Maxeiner, J., see Schöneshöfer, M. (224) 229
May, M.E., see Andrews, P.A. (227) 83
Meering, P.G. and Maes, R.A.A.
Simultaneous determination of misonidazole and desmethylmisonidazole in plasma by HPLC (225) 407
Meffin, P.J., see Veenendaal, J.R. (223) 147
Meinertz, T., see Wiegand, U.-W. (223) 238
Merdink, J.L., see Abdel-Monem, M.M. (222) 363
Meredith, P., see Rubin, P.C. (221) 193
Metayer, J.-P., see Sioufi, A. (222) 429
Meulemans, A., Mohler, J., Henzel, D. and Duvaldestin, P.
Quantitation of D-tubocurarine in human plasma using HPLC (226) 255
Meyer, M.C. and Hwang, P.T.R.
Determination of trichlormethiazide in human plasma and urine by HPLC (223) 466
Miceli, J.N., see Aravind, M.K. (221) 176, (228) 418
Michaud, R.L., Bannon, M.J. and Roth, R.H.
The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335
Michielsen, L., see Woestenborghs, R. (224) 25, (224) 122
Michotte, Y., see Hoogewijs, G. (226) 423
Mico, B.A., Baughman, R.A. Jr. and Benet, L.Z.
Determination of tiodazosin in plasma and whole blood by HPLC (230) 203
Midha, K.K., see Charette, C. (224) 128
Mikkers, F., see Oerlemans, F. (225) 369
Miller, A.A., Benvenuto, J.A. and Loo, T.L.
Comparison of the reversed-phase HPLC separations of fluoropyrimidines, pyrimidines, and purines (228) 165
Miller, J.N., see Walmsley, L.M. (226) 441
Miller, K.F., see Weinfeld, R.E. (223) 123
Miller, M., see Ward, L.C. (223) 417
Miller, R., see Jatlow, P.I. (227) 233
Miller, R.D., see Shinohara, Y. (230) 363
Millingen, K.S., see Roberts, M.S. (226) 175
Millot, C., see Ramirez, L.C. (229) 267
Milner, J.A., see McCarthy, T.P. (225) 9
Miners, J.O., see Chivers, D.A. (225) 261
Mita, H., Yasueda, H. and Shida, T.
Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human plasma and urine by GC-MS (221) 1
Mitchell, J.R., see Lauterburg, B.H. (224) 431
Mitomi, M., see Katogi, Y. (228) 404
Miura, K., Nakamura, H. and Tamura, Z.
A new method for detection of drug-binding proteins using a parallel-flow dialysis technique (223) 267
Miyai, K., see Yamaguchi, Y. (224) 111
Miyatake, T., see Suzuki, M. (221) 361
Miyazaki, H., see Idzu, G. (229) 327
Miyazaki, K., Arita, T., Oka, I., Koyama, T. and Yamashita, I.
HPLC determination of haloperidol in plasma (223) 449
Miyazaki, M., see Hayakawa, K. (229) 159
Moerman, E., see Van Aerde, P. (222) 467
Moffat, A.C., see Gill, R. (226) 117
Mohammed, H.Y., Veening, H. and Dayton, D.A.
LC determination and time-concentration studies of riboflavin in hemodialysate from uremic patients (226) 471
Mohler, J., see Meulemans, A. (226) 255
Mokofio, F., see Bonini, M. (224) 332
Moleman, P. and Borstrok, J.J.M.
Analysis of urinary 3-methoxy-4-hydroxy-phenylglycol by HPLC and electrochemical detection (227) 391
Molin, L., see Eriksson, B.-M. (223) 401
Möller, H., see Stüber, W. (224) 327
Møller Heilesen, A., see Larsen, N.-E. (221) 182
Moore, S.A., see Rhodes, G. (228) 33
Moore, S.A., see Smith, S.L. (221) 19
Moran, T.F., see Reiner, E.J. (221) 371
Mori, H., see Iwai, M. (225) 275
Mori, Y., see Konishi, M. (229) 355
Morison, D.H., see Gupta, R.N. (224) 496
Morita, I., see Suzuki, M. (221) 361
Moro, E., Jannuzzo, M.G., Ranghieri, M., Stegnjaich, S. and Valzelli, G.
Determination of 4'-epidoxorubicin and its 13-dihydro derivative in human plasma by HPLC with fluorescence detection (230) 207
Morris, H.S., see Glencross, R.G. (223) 193
Morris, J.E., Canoy, D.W. and Rynd, L.S.
Electrophoresis with two buffers in one dimension in the analysis of glycosaminoglycans on cellulose acetate strips (224) 407
Morrison, P.J., see Barnicoat, A.J. (225) 236
Morselli, P.L., see Gillet, G. (230) 154
Mortimer, R.H., see Cannell, G.R. (226) 492
Moschitto, L.J., see Greenblatt, D.J. (225) 202
Mosnaim, A.D., Callaghan, O.H. and Wolf, M.E.
Determination of noncatecholic phenylethyl-amines and monomethylated derivatives of phenylethylamine (224) 481
Mosser, D.D., see Bols, N.C. (229) 460
Moulin, M.A., Albessard, F., Lacotte, J. and Camsonne, R.
Hydrophilic ion-pair reversed-phase HPLC for the simultaneous assay of isoniazid and acetylisoniazid in serum: a microscale procedure (226) 250
Mount, J.N. and Laker, M.F.
Estimation of sugar alcohols by GLC using a modified acetylation procedure (226) 191
Mrongovius, R., see Müller, H. (226) 450
Mueller, H.W. and Binz, K.
Glass capillary GC of the serum fatty acid fraction via automatic injection of lipid extracts (228) 75
Muir, K.T., Jonkman, J.H.G., Tang, D.-S., Kunitani, M. and Riegelman, S.

Simultaneous determination of theophylline and its major metabolites in urine by reversed-phase ion-pair HPLC (221) 85
Muirhead, D.C., see Vose, C.W. (222) 311
Mulder, T.B.A., see Feenstra, M.G.P. (230) 271
Müller, H., Mrongovius, R. and Seyberth, H.W. Improved sample preparation for the quantitative MS determination of prostaglandins in biological samples (226) 450
Müller, K., see Lankmayr, E.P. (222) 249
Murakami, K., Murakami, K., Ueno, T., Hijikata, J., Shirasawa, K. and Muto, T. Simultaneous determination of chlorpromazine and levomepromazine in human plasma and urine by HPLC using electrochemical detection (227) 103
Murakami, K., Ueno, N. and Hirose, S. Demonstration of the formation of renin and renin-binding protein complex using HPLC (225) 329
Murakami, K., see Murakami, K. (227) 103
Murao, M., see Kaji, H. (221) 145
Murayama, K. and Sugawara, T. Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315
Murdoch, S., see Vijayakumar, R. (225) 47
Murdock, L.L., see Omar, D. (224) 310
Murota, S.-I., see Suzuki, M. (221) 361
Murray, G.R., see Chan, K. (223) 213
Muskiet, F.A.J., Stratingh, C.M. and Fremouw-Ottevangers, D.C. MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142
Muskiet, F.A.J., see Pavel, S. (222) 329
Mussini, E., Cotellessa, L., Colombo, L., Cani, D., Sfondrini, P., Marcucci, F. and Poy, F. Glass capillary quantitative determination of $N^{\pi}$-methylhistidine in urine and muscles (224) 94
Mussini, E., see Cotellessa, L. (221) 149
Muto, T., see Murakami, K. (227) 103
Mutschler, E., see Schäfer-Korting, M. (230) 461
Mutschler, E., see Stüber, W. (227) 193
Myers, C., see Ackerly, C.C. (230) 175
Myher, J.J., see Kuksis, A. (224) 1, (230) 231

# N

Nacht, S., see Yeung, D. (224) 513
Nachtmann, F., see Lankmayr, E.P. (222) 249, (224) 239
Nagase, H., see Suzuki, M. (221) 361
Nagatsu, T., see Rahman, M.K. (221) 265
Nagatsu, T., see Trocewicz, J. (227) 407
Nagatsu, T., see Uji, A. (221) 155
Nagel, G.T., see Pavel, S. (222) 329
Nahata, M.C. HPLC determination of cephalexin in human plasma, urine and saliva (225) 532
Nahata, M.C. Determination of cefaclor by HPLC (228) 429
Nahata, M.C. and Powell, D.A. Simultaneous determination of chloramphenicol and its succinate ester by HPLC (223) 247
Nakae, H., see Takada, K. (222) 156
Nakahashi, H., see Mabuchi, H. (224) 322, (228) 292
Nakai, K., see Umeno, Y. (226) 333
Nakajima, O., Yoshida, Y., Isoda, T., Takemasa, Y., Imamura, Y. and Koyama, Y. Quantitative determination of 1-hexylcarbamoyl-5-fluorouracil and its metabolites in man (225) 91
Nakajima, O., Yoshida, Y., Kubota, T., Takemasa, Y. and Koyama, Y. Quantitative determination of 1-(2-chloroethyl)-3-(β-D-glucopyranosyl)-1-nitrosourea in blood and urine of man (229) 481
Nakamura, H., see Miura, K. (223) 267
Nakamura, T., see Goto, M. (226) 33
Nakanishi, M., see Sadanaga, T. (223) 243
Nakano, S., see Watanabe, H. (229) 95
Nakatsu, K., see Tasker, R.A.R. (228) 346
Nakayama, T., see Iwai, M. (225) 275
Nambara, T., see Goto, J. (226) 13
Nambara, T., see Shimada, K. (223) 33, (227) 445
Narang, P.K., Yeager, R.L. and Chatterji, D.C. Quantitation of 6-mercaptopurine in biologic fluids using HPLC: a selective and novel procedure (230) 373
Narasimhachari, N. Evaluation of $C_{18}$ Sep-Pak cartridges for biological sample clean-up for tricyclic antidepressant assays (225) 189
Naruse, H., see Hayashi, T. (224) 197
Nau, H., Wittfoht, W., Schäfer, H., Jakobs, C., Rating, D. and Helge, H. Valproic acid and several metabolites: quantitative determination in serum, urine, breast milk and tissues by GC-MS using selected ion monitoring (226) 69
Nau, H., see Liddiard, C. (225) 504
Nealon, T.F. Jr., see Ching, N.P.H. (222) 171, (225) 196
Necciari, J., see Sommadossi, J.P. (228) 205
Neeley, W.E., see Alexander, N.M. (230) 137
Negi, D.S., see Thomas, D.W. (225) 433
Neiderhiser, D.H. and Fuller, R.K. HPLC method for the determination of diethylamine a metabolite of disulfiram, in urine (229) 470

Neill, D.B., see Holdiness, M.R. (225) 283
Neill, D.B., see Saraswat, L.D. (222) 353
Nelson, S.D., see Wilson, J.M. (227) 453
Nemoto, E.M., see Shiu, G.K. (227) 207
Newman, R.A., see Ackerly, C.C. (230) 175
Newman, R.A., see Allen, B.A. (222) 146
Newport, G.D., see Slikker, W. Jr. (224) 205
Ng, K.J., Andresen, B.D., Bianchine, J.R., Iams, J.D., O'Shaugnessy, R.W., Stempel, L.E. and Zuspan, F.P.
Capillary GC-MS profiles of trimethylsilyl derivatives of organic acids from amniotic fluids of different gestational age (228) 43
Nguyen-Dinh, P., see Staiger, M.A. (225) 139
Nice, E.C., see Zanelli, J.M. (223) 59
Nielsen, A., see Bechgaard, E. (228) 392
Nilsson, B., see Johansson, B. (229) 439
Nilsson, L.B., see Magnusson, O. (221) 237
Nimura, N., Ishida, K. and Kinoshita, T.
Novel post-column derivatization method for the fluorimetric determination of norepinephrine and epinephrine (221) 249
Nishida, S., see Kakita, K. (222) 33
Nishide, K., see Matsumura, K. (230) 148
Nishimura, R., see Konaka, R. (225) 169
Nishizawa, M., see Suzuki, M. (221) 361
Nissen, P.
Simultaneous determination of allopurinol, oxipurinol and uric acid in human plasma by HPLC (228) 382
Nitsche, V. and Mascher, H.
Rapid HPLC assay of cinnarizine in human plasma (227) 521
Niwa, T., Maeda, K., Asada, H., Shibata, M., Ohki, T., Saito, A. and Furukawa, H.
GC-MS analysis of organic acids in renal tissue biopsy. Identification of 4-hydroxybutyric acid and 4-hydroxy-2-butenoic acid (230) 1
Niwa, T., Maeda, K., Ohki, T., Saito, A. and Tsuchida, I.
GC-MS profile of organic acids in urine and serum of diabetic ketotic patients (225) 1
Niwa, T., Maeda, K., Ohki, T. and Sakakibara, J.
Identification of 2-hydroxy-2-methyllevulinic acid in urine and serum of diabetic patients with ketoacidosis (228) 59
Niwa, T., see Maeda, K. (221) 199
Niwa, T., see Ohki, T. (228) 51
Njapau, H., see Lovelace, C.E.A. (227) 256
Noda, A., Hsu, K.-Y., Aso, Y., Matsuyama, K., Iguchi, S. and Hirata, M.
Studies of isoniazid metabolism in isolated rat hepatocytes by mass fragmentography (230) 345
Nohta, H., Ohtsubo, K., Zaitsu, K. and Ohkura, Y.
Assay for dopamine β-hydroxylase in rat serum and adrenal medulla by HPLC with fluorescence detection (227) 415
Nomiyama, H., see Nomiyama, K. (228) 285
Nomiyama, K. and Nomiyama, H.
HPLC determination of tissue metallothionein in monkeys chronically exposed to cadmium (228) 285
Norman, T.R., see Maguire, K.P. (222) 399
Norris, R.L.G., Ahokas, J.T. and Ravenscroft, P.J.
Simultaneous HPLC assay for quinidine, disopyramide and the mono-N-dealkylated metabolite of disopyramide (230) 433
Novotný, M., see Rhodes, G. (228) 33
Novotný, M., see Smith, S.L. (221) 19
Novotný, M.V., see Smith, S.L. (223) 173

# O

Ober, R.E., see Chang, S.F. (226) 79
Odlind, B., see Floberg, S. (225) 73
Oerlemans, F., De Bruyn, C., Mikkers, F., Verheggen, T. and Everaerts, F.
Isotachophoresis of urinary purines and pyrimidines. The use of spacers and enzymes for identification (225) 369
Ogawa, N., see Watanabe, H. (229) 95
O'Hare, M.J., see Zanelli, J.M. (223) 59
Ohki, T., Saito, A., Yamanaka, N., Ohta, K., Sakakibara, J., Niwa, T. and Maeda, K.
GC-MS analysis of amine metabolites in urine (228) 51
Ohki, T., see Maeda, K. (221) 199
Ohki, T., see Niwa, T. (225) 1, (228) 59, (230) 1
Ohkura, Y., see Hirata, T. (226) 25
Ohkura, Y., see Nohta, H. (227) 415
Ohkura, Y., see Sadanaga, T. (223) 243
Ohkura, Y., see Tsuruta, Y. (224) 105
Ohno, K., see Suzuki, M. (221) 361
Ohno, Y., see Okazaki, M. (221) 257, (223) 285
Ohta, K., see Ohki, T. (228) 51
Ohtsubo, K., see Nohta, H. (227) 415
Ohya, K., Shintani, S. and Sano, M.
Determination of ethambutol in plasma using selected ion monitoring (221) 293
Ohya, K., Shintani, S., Suzuki, W. and Sano, M.
Sensitive and selective method for the determination of chlormezanone in plasma by electron-capture GC (221) 67
Oka, I., see Miyazaki, K. (223) 449
Oka, K., see Trocewicz, J. (227) 407
Oka, K., see Yamada, K. (223) 176
Okamoto, M., Yamada, F., Ishiguro, M. and Yasue, T.
HPLC determination of bromural in serum upon hemoperfusion (223) 473
Okamoto, M., see Koh, S. (226) 461
Okazaki, M., Ohno, Y. and Hara, I.
High-performance aqueous gel permeation chromatography of human serum lipoproteins (221) 257

Okazaki, M., Shiraishi, K., Ohno, Y. and Hara, I.
High-performance aqueous gel permeation chromatography of serum lipoproteins: selective detection of cholesterol by enzymatic reaction (223) 285
Okuyama, T., see Manabe, T. (225) 65
Okuyama, T., see Seta, K. (221) 215
Omar, D. and Murdock, L.L.
Determination of N-acetyldopamine by LC with electrochemical detection (224) 310
Omichi, K. and Ikenaka, T.
Rapid and sensitive sialidase assay by HPLC and its application to detection of sialidase in human urine (230) 415
Ong, H., Sved, S. and Beaudoin, N.
Assay and stability of α-methyldopa in man using HPLC with electrochemical detection (229) 433
Oosterhuis, B., Van den Berg, M. and Van Boxtel, C.J.
Sensitive HPLC method for the determination of labetalol in human plasma using fluorimetric detection (226) 259
Orr, J., see Abbott, F.S. (227) 433
O'Shaugnessy, R.W., see Ng, K.J. (228) 43
Østensen, E.T., see Øyehaug, E. (227) 129
Ottoila, P. and Taskinen, J.
Determination of biperiden in human serum by glass capillary GC with isothermal splitless injection and nitrogen-sensitive detection (226) 488
Overø, K.F.
Fluorescence assay of citalopram and its metabolites in plasma by scanning densitometry of thin-layer chromatograms (224) 526
Owen, P.E., see Carey, P.F. (225) 161
Øyehaug, E., Østensen, E.T. and Salvesen, B.
Determination of the antidepressant agent citalopram and metabolites in plasma by LC with fluorescence detection (227) 129

# P

Pagani, E., see Longo, A. (224) 257
Pahuja, S.L., Albert, J. and Reid, T.W.
Use of reversed-phase HPLC for simultaneous determination of glutamine synthetase and glutamic acid decarboxylase in crude extracts (225) 37
Palmskog, G.
Determination of plasma levels of two aromatic retinoic acid analogues with antipsoriatic activity by HPLC (221) 345
Panaqua, M., see McKee, R.W. (230) 309
Pao, J. and De Silva, J.A.F.
Determination of the quinidine analog, 7'-trifluoromethyldihydrocinchonidine·2HCl in plasma and urine by HPLC (221) 97
Parenti, R.R., see Ratti, B. (225) 526
Park, G.B., Biddlecome, C.E., Koblantz, C. and Edelson, J.
Determination of ciprofibrate in human plasma by HPLC (227) 534
Park, G.B., Erdtmansky, P., Kullberg, M.P. and Edelson, J.
The analysis of arildone in plasma, urine and feces by GLC with electron-capture detection (222) 213
Parkhurst, R.M., see Thomas, D.W. (225) 433
Parkin, J.E.
Determination of alcuronium chloride in biological fluids by HPLC (225) 240
Pasquini, J.M., see Bizzozero, O. (227) 33
Pateman, A.J.
Sensitive GC method for the determination of alphadolone in plasma (226) 213
Patrizi, V.W., see Strauss, J.F. (221) 139
Patton, T.F., see Sternson, L.A. (227) 223
Paunio, K., see Virtanen, P. (221) 49
Pautler, D.B. and Jusko, W.J.
Determination of metoprolol and α-hydroxy-metoprolol in plasma by HPLC (228) 215
Pavel, S., Muskiet, F.A.J., Nagel, G.T., Schwippelová, Z. and Duchoň, J.
Identification of two Thormählen-positive compounds from melanotic urine by GC-MS (222) 329
Pays, M., see Biou, D. (226) 477
Pays, M., see Leroyer, R. (228) 366
Pedersen, A.K., see Jakobsen, P. (223) 460
Peller, J.D., see Rao, G.H.R. (226) 466, (229) 205
Peña, M.I. Arranz, see Arranz Peña, M.I. (222) 486
Perhach, J.L. Jr., see Hartman, C.A. (226) 510
Perrin, J.H., see Fleitman, J. (229) 255
Perrin, J.H., see Fleitman, J.S. (228) 372
Persson, B.-A., see Eriksson, B.-M. (228) 143
Peters, J.G.P., see Ketelaars, H.C.J. (224) 144
Peters, J.H., Hamme, K.J. and Gordon, G.R.
Determination of clofazimine in plasma by HPLC (229) 503
Peterson, R.G., see Rhodes, G. (228) 33
Petit, J., see Chabard, J.L. (221) 9
Petit, J., see Lartigue-Mattei, C. (229) 211
Pettibone, D., see Spindel, E. (222) 381
Pfeifer, P. and Spiteller, G.
Steroid profiles of healthy individuals (223) 21
Pfleger, K., see Maurer, H. (222) 409
Phillipou, G., see Harman, P.J. (225) 131
Pickup, M.E., Lowe, J.R. and Galloway, D.B.
Determination of Ro12-0068, a new anti-inflammatory and analgesic compound, in plasma by means of HPLC (225) 493
Pieroni, J.P., Lee, W.H. and Wong, P.Y.-K.
Separation of six major prostacyclin metabolites by HPLC (230) 115
Pietta, P., Calatroni, A. and Rava, A.
HPLC assay for monitoring indapamide and its major metabolite in urine (228) 377

Pietta, P., Calatroni, A. and Rava, A.
Hydrolysis of riboflavin nucleotides in plasma monitored by HPLC (229) 445
Pifferi, G., see Crolla, T. (222) 257
Pillai, G.K., Axelson, J.E. and McErlane, K.M.
Electron-capture GLC determination of tocainide in biological fluids using fused silica capillary columns (229) 103
Pinkston, D., Spiteller, G., Von Henning, H. and Matthaei, D.
High-resolution GC-MS of the methyl esters of organic acids from uremic hemofiltrates (223) 1
Pittman, K.A., see Fandrich, S.E. (223) 155
Plummer, G.F., see Adam, H.K. (223) 232
Plunkett, W., Chubb, S. and Barlogie, B.
Simultaneous determination of 1-β-D-arabinofuranosylcytosine 5'-triphosphate and 3-deazauridine 5'-triphosphate in human leukemia cells by HPLC (221) 425
Pollard, A.C., see Badcock, N.R. (230) 353
Poma, S., see Cassara, G. (221) 279
Pommier, F., see Sioufi, A. (221) 377, (222) 429, (226) 219, (229) 347
Poon, B.T., see Brown, N.D. (229) 248
Porcelli, M., see Della Ragione, F. (226) 243
Porsius, A.J., see De Neef, J.H. (224) 133
Porsius, A.J., see De Wildt, D.J. (225) 381
Porta, R., Esposito, C. and Sellinger, O.Z.
Rapid assay of spermidine synthase activity by HPLC (226) 208
Powell, D.A., see Nahata, M.C. (223) 247
Powis, G.
Reversed-phase HPLC assay for the antineoplastic agent 9,10-anthracenedicarboxaldehyde bis(4,5-dihydro-1H-imidazol-2-yl hydrazone) dihydrochloride (226) 514
Poy, F., see Cotellessa, L. (221) 149
Poy, F., see Mussini, E. (224) 94
Prasad, V.K., see Rapaka, R.S. (227) 463
Premel-Cabic, A., see Turcant, A. (229) 222
Prescott, L.F., see Balali-Mood, M. (229) 234
Preti, G., see Kostelc, J.G. (226) 315
Pritchett, E.L.C., see Shand, D.G. (224) 343
Prussak, C.E. and Russell, D.H.
Single-step HPLC method for the analysis of acetylated polyamines (229) 47
Puech, R., see Alric, R. (224) 289
Puglisi, C.V. and De Silva, J.A.F.
Determination of the anxiolytic agent 8-chloro-6-(2-chlorophenyl)-4H-imidazo-[1,5-*a*][1,4]-benzodiazepine-3-carboxamide in whole blood, plasma or urine by HPLC (226) 135
Puglisi, C.V., see Cotler, S. (222) 95
Puzic, O., see Anastassiades, T. (225) 309
Puzic, R., see Anastassiades, T. (225) 309

# Q

Quercia, G.T., see Dokladalova, J. (224) 33
Queyrel, N., see Biou, D. (226) 477
Qureshi, G.A., see Allenmark, S. (223) 188

# R

Ragione, F. della, see Della Ragione, F. (226) 243
Rahman, M.K., Nagatsu, T. and Kato, T.
New and highly sensitive assay for L-5-hydroxytryptophan decarboxylase activity by HPLC-voltammetry (221) 265
Rainer, F., see Lankmayr, E.P. (222) 249
Raisys, V.A., see Watts, M.T. (230) 79
Rajagopalan, T.G., Anjaneyulu, B., Shanbag, V.D. and Grewal, R.S.
Electron-capture GC assay for primaquine in blood (224) 265
Ramirez, L.C., Millot, C. and Maume, B.F.
Sample purification using a $C_{18}$-bonded reversed-phase cartridge for the quantitative analysis of corticosteroids in adrenal cell cultures by HPLC or GC-MS (229) 267
Rane, A., see Svensson, J.-O. (230) 427
Ranghieri, M., see Moro, E. (230) 207
Ranise, A., Benassi, E. and Besio, G.
Rapid GC method for the determination of carbamazepine and unrearranged carbamazepine-10,11-epoxide in human plasma (222) 120
Rao, G.H.R., Peller, J.D., Richards, K.L., McCullough, J. and White, J.G.
Rapid separation of nucleotides from granulocytes by isocratic, reversed-phase HPLC (229) 205
Rao, G.H.R., Peller, J.D. and White, J.G.
Rapid separation of platelet nucleotides by reversed-phase, isocratic, HPLC with a radially compressed column (226) 466
Rapaka, R.S., Roth, J., Viswanathan, C., Goehl, T.J., Prasad, V.K. and Cabana, B.E.
Improved method for the analysis of furosemide in plasma by HPLC (227) 463
Ratge, D. and Wisser, H.
Urinary protein profiling by high-performance gel permeation chromatography (230) 47
Rathgen, G.H., see Höckel, M. (221) 205
Rating, D., see Nau, H. (226) 69
Ratti, B., Parenti, R.R., Toselli, A. and Zerilli, L.F.
Quantitative assay of rifampicin and its

main metabolite 25-desacetylrifampicin in human plasma by reversed-phase HPLC (225) 526
Rauscher, E., see Haegele, E.-O. (223) 69
Rava, A., see Pietta, P. (228) 377, (229) 445
Ravenscroft, P.J., see Charles, B.G. (222) 152
Ravenscroft, P.J., see Norris, R.L.G. (230) 433
Reece, P.A.
Quantification of prazosin in plasma by HPLC (221) 188
Reece, P.A., Cozamanis, I. and Zacest, R.
Sensitive HPLC assay for endralazine and two of its metabolites in human plasma (225) 151
Reese, A.L.P., see Kuhnert, B.R. (224) 488
Reeve, J. Jr., Yamada, T., Chew, P., Walsh, J.H. and Dimaline, R.
Rapid high-yield purification of neuropeptides from canine intestinal muscle (229) 57
Rehman, A., Gates, S.C. and Webb, J.W.
Comparison of isolation methods of urinary organic acids HPLC (228) 103
Reid, T.W., see Pahuja, S.L. (225) 37
Reiner, E.J. and Moran, T.F.
High-performance pyrolysis-GC: potential for differentiating cystic fibrosis cells (221) 371
Reisch, J. and El-Sharaky, A.S.
Natural products chemistry. 81. Analysis of rat thymus steroids by liquid-gel chromatography and GC-MS (222) 475
Rennert, O.M., see Griesmann, G.E. (230) 121
Repta, A.J., see Reynolds, D.L. (222) 225
Repta, A.J., see Riley, C.M. (229) 373
Reynolds, D.L., Sternson, L.A. and Repta, A.J.
Clinical analysis for the anti-neoplastic agent 1,4-dihydroxy-5,8-bis{{2-[(2-hydroxyethyl)aminolethyl}amino}9,10-anthracenedione dihydrochloride (NSC 301739) in plasma. Application of temperature control to provide selectivity in paired-ion HPLC (222) 225
Reynolds, G.P., see Riederer, P. (225) 179
Rhodes, G., Holland, M.L., Wiesler, D., Novotný, M., Moore, S.A., Peterson, R.G. and Felten, D.L.
Excretion of urinary volatile metabolites in response to alloxan induced diabetes of short duration in rats (228) 33
Richard, A., see Sioufi, A. (221) 393
Richards, K.L., see Rao, G.H.R. (229) 205
Richter, E., see Heusler, H. (226) 403
Richter, W.O., see Schwandt, P. (225) 185
Ried, W., see Winkler, H. (228) 223
Riedel, E., Kreutz, G. and Hermsdorf, D.
Quantitative TLC determination of dihydroergot alkaloids (229) 417
Riederer, P. and Reynolds, G.P.
Determination of a wide range of urinary amine metabolites using a simple HPLC technique (225) 179
Riegelman, S., see Kunitani, M.G. (224) 156
Riegelman, S., see Muir, K.T. (221) 85
Righetti, P.G., see Cassara, G. (221) 279
Riley, C.M., Sternson, L.A., Repta, A.J. and Siegler, R.W.
HPLC of platinum complexes on solvent generated anion exchangers. III. Application to the analysis of cisplatin in urine using automated column switching (229) 373
Ripani, L., see Di Corcia, A. (229) 365
Risby, D., see Joseph, M.H. (226) 361
Ritz, E., see Tschöpe, W. (221) 59, (222) 41
Riva, R., Albani, F. and Baruzzi, A.
Rapid quantitative determination of underivatized carbamazepine, phenytoin, phenobarbital and *p*-hydroxyphenobarbital in biological fluids by packed column GC (221) 75
Riva, R., De Anna, M., Albani, F. and Baruzzi, A.
Rapid quantitation of flurazepam and its major metabolite, N-desalkylflurazepam, in human plasma by GLC with electron-capture detection (222) 491
Riva, R., Tedeschi, G., Albani, F. and Baruzzi, A.
Quantitative determination of clobazam in the plasma of epileptic patients by GLC with electron-capture detection (225) 219
Riva, R., see Albani, F. (228) 362
Rizzardini, M., see Cantoni, L. (229) 311
Robert, T.A., Hagardorn, A.N., Daigneault, E.A. and Brown, R.D.
Analysis of trimethobenzamide in human saliva by GC-MS (224) 116
Roberts, M.S., Watson, H.M., McLean, S. and Millingen, K.S.
Determination of therapeutic plasma concentrations of tetrabenazine and an active metabolite by HPLC (226) 175
Roberts, M.S., see Rumble, R.H. (225) 252
Robinson, M.M., see DeAngelis, R.L. (221) 353
Rocchiccioli, F., Leroux, J.P. and Cartier, P.
Quantitative GC-chemical ionization MS of 2-ketoglutarate from urine as its O-trimethylsilyl-quinoxalinol derivative (226) 325
Rocci, M.L. Jr. and Jusko, W.J.
Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221
Rohdewald, P. and Drehsen, G.
Rapid TLC of various weak analgesics in saliva (225) 427
Rohdewald, P., see Drehsen, G. (223) 479
Rollema, H., see Feenstra, M.G.P. (230) 271
Rooy, H.H. van, see Van Rooy, H.H. (223) 85, (224) 133, (225) 381
Rosa, J., see Basset, P. (227) 267
Rosenberg, M., see Choi, R.L. (230) 181
Rosset, R., see Guyon, F. (229) 475
Rostron, C., see Chan, K. (223) 213
Roth, J., see Rapaka, R.S. (227) 463
Roth, R.H., see Michaud, R.L. (225) 335
Roth, W., Beschke, K., Jauch, R., Zimmer, A. and Koss, F.W.
Fully automated HPLC. A new

chromatograph for pharmacokinetic drug monitoring by direct injection of body fluids (222) 13
Rouan, M.C., see Lecaillon, J.B. (228) 257
Roux, Y. le, see Le Roux, Y. (230) 401
Rovei, V., see Broquaire, M. (224) 43
Rovei, V., see Guinebault, P.R. (223) 103
Rowland, J.M., see Althaus, Z.R. (227) 11
Rubin, P.C., Brunton, J. and Meredith, P. Determination of the vasodilator UK33274 by HPLC using fluorescence detection (221) 193
Rufini, S., see Ghebregzabher, M. (222) 191
Ruggieri, R., see Cantoni, L. (229) 311
Rumble, R.H., Roberts, M.S. and Wanwimolruk, S. Determination of aspirin and its major metabolites in plasma by HPLC without solvent extraction (225) 252
Russell, D.H., see Prussak, C.E. (229) 47
Rustum, Y.M., see Au, J.L.-S. (228) 245
Ryan, A.J., see Boux, L.J. (227) 149
Ryle, P.R., see Jones, C.R. (224) 492
Rynd, L.S., see Morris, J.E. (224) 407

# S

Sadanaga, T., Hikida, K., Tameto, K., Sugimoto, K., Nakanishi, M., Matsushima, Y. and Ohkura, Y. Determination of osalmid in plasma by HPLC (223) 243
Sagraves, R., see Velagapudi, R. (228) 423
Sahawneh, T., see Vasiliades, J. (228) 195
Sahawneh, T.H., see Vasiliades, J. (225) 266
Sahlberg, B.-L., see Axelson, M. (224) 355
Saito, A., see Niwa, T. (225) 1, (230) 1
Saito, A., see Ohki, T. (228) 51
Saito, K., see Kaji, H. (221) 145
Saito, N., see Kaji, H. (221) 145
Sajgó, M., Gesztesi, A. and Sidó, T. Determination of tofisopam in serum by HPLC (222) 303
Sakai, T., see Yoshida, A. (227) 162
Sakakibara, J., see Niwa, T. (228) 59
Sakakibara, J., see Ohki, T. (228) 51
Sakashita, T., see Matsumura, K. (222) 53
Salamone, J.D., see Holdiness, M.R. (225) 283
Salamone, J.D., see Saraswat, L.D. (222) 353
Sales, Y., see Cano, J.P. (226) 413
Salimi, S.L., see Tsao, C.S. (224) 477
Salinas, F.A., see Silver, H.K.B. (224) 381
Salter, L.F., see Lovelace, C.E.A. (227) 256
Salvesen, B. and Haugland, T. Determination of oxeladin in human sera by GLC with thermionic detection (225) 463
Salvesen, B., see Øyehaug, E. (227) 129
Samperi, R., see Di Corcia, A. (229) 365
Samson, D., see Crowne, H. (223) 421
Sandler, S., see Goldberg, E.M. (226) 291
Sandrenan, N., see Sioufi, A. (221) 377
Sanitrák, J., see Macek, K. (222) 284
Sanjuan, M., see Guinebault, P.R. (223) 103
Sano, M., see Ohya, K. (221) 67, (221) 293
Saraswat, L.D., Holdiness, M.R., Justice, J.B., Salamone, J.D. and Neill, D.B. Determination of dopamine, homovanillic acid and 3,4-dihydroxyphenylacetic acid in rat brain striatum by HPLC with electrochemical detection (222) 353
Saria, A., see Skofitsch, G. (226) 53
Saruta, Y., see Goto, J. (226) 13
Sasaki, H., see Kamimura, H. (225) 115, (229) 337
Sassard, J., see Bernard, N. (228) 355
Sattsangi, S., see Tryfiates, G.P. (227) 181
Säwe, J., see Svensson, J.-O. (230) 427
Scaro, J., see Shihabi, Z.K. (224) 99
Schaal, S.F., see Haughey, D.B. (229) 387
Schäfer, H., see Nau, H. (226) 69
Schäfer-Korting, M. and Mutschler, E. Fluorodensitometric determination of nadolol in plasma and urine (230) 461
Schaich, E., see Haegele, E.-O. (223) 69
Schaik, F. van, see Van Schaik, F. (228) 311
Schick, J., see Breithaupt, H. (225) 99
Schier, G.M. and Eng Tho Gan, I. Measurement of plasma theophylline by GLC on the stationary phase SP-2510 (225) 208
Schlöbe, R. and Thijssen, H.H.W. Quantitative TLC of trimethoprim and tetroxoprim using fluorescence densitometry (230) 212
Schmid, J. Assay of prolintane in plasma by capillary GC with nitrogen-selective detection (222) 129
Schmid, J. and Koss, F.-W. Assay of bromhexine in human plasma by capillary GLC with nitrogen-selective detection and selected ion monitoring (227) 71
Schneider, Z., Leszczyńska, D. and Socha, J. Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77
Schöneshöfer, M. and Fenner, A. Hydrophilic ion-pair reversed-phase chromatography of biogenic peptides prior to immunoassay (224) 472
Schöneshöfer, M., Maxeiner, J. and Fenner, A. Quantitation of pregnenolone and 17-hydroxypregnenolone in human serum by automatic HPLC and subsequent radioimmunoassay (224) 229
Schöneshöfer, M., Skobolo, R. and Dulce, H.J. Reliability of the estimation of serum cortisol by HPLC (222) 478
Schöneshöfer, M., Weber, B., Dulce, H.J. and Belkien, L. Estimation of free urinary aldosterone and

18-hydroxycorticosterone by a combination of automatic HPLC and radioimmunoassay (227) 492
Schöpf, J., see Tinguely, D. (229) 319
Schouten, T.J., see De Abreu, R.A. (227) 526
Schretlen, E.D.A.M., see De Abreu, R.A. (227) 45, (227) 526, (229) 67
Schreuers, W.H.P., see Speek, A.J. (228) 311
Schrijver, J., see Speek, A.J. (228) 311
Schulman, S.G., see Fleitman, J.S. (228) 372
Schultz, B. and Hansen, S.H.
Determination of the γ-aminobutyric acid agonist 4,5,6,7-tetrahydroisoxazolo[5,4-c]pyridine-3-ol in urine by HPLC (228) 279
Schwandt, P., Richter, W.O. and Weisweiler, P.
Separation of human C-apolipoproteins by HPLC (225) 185
Schwarz, S. and Stecher, W.
Simple computer program for a low-cost desk-top calculator applied to the evaluation of GLC analyses of 17-ketosteroids and pregnanes (223) 253
Schwippelová, Z., see Pavel, S. (222) 329
Scoggins, B.A., see Maguire, K.P. (222) 399
Scortecci, V., see Cassara, G. (221) 279
Scriver, C.R., see Cole, D.E.C. (225) 359
Seamark, D.A., see Trafford, D.J.H. (226) 351
Segal, S., see Yandrasitz, J.R. (225) 319
Seiler, N. and Knödgen, B.
HPLC procedure for the simultaneous determination of the natural polyamines and their monoacetyl derivatives (221) 227
Sellinger, O.Z., see Porta, R. (226) 208
Senn, M., see Endele, R. (227) 187
Sennello, L.T., see Granneman, G.R. (229) 149
Seppälä, E., see Kärkkäinen, S. (221) 319
Seta, K., Washitake, M., Tanaka, I., Takai, N. and Okuyama, T.
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215
Severen, R. van, see Van Severen, R. (222) 467
Seyberth, H.W., see Müller, H. (226) 450
Sfondrini, P., see Cotellessa, L. (221) 149
Sfondrini, P., see Mussini, E. (224) 94
Shader, R.I., see Greenblatt, D.J. (225) 202
Shanbag, V.D., see Rajagopalan, T.G. (224) 265
Shand, D.G., Verghese, C., Barchowsky, A., Hammill, S.C. and Pritchett, E.L.C.
HPLC analysis of a new antiarrhythmic drug, pirmenol, in biological fluids (224) 343
Shargel, L., see Ameer, B. (226) 224
Shaywitz, B.A., see Anderson, G.M. (222) 112, (223) 315
Shiba, D.A. and Weinkam, R.J.
Quantitative analysis of procarbazine, procarbazine metabolites and chemical degradation products with application to pharmacokinetic studies (229) 397
Shibata, M., see Niwa, T. (230) 1
Shida, T., see Mita, H. (221) 1
Shigematsu, A., see Manabe, T. (225) 65
Shihabi, Z.K., Scaro, J. and Thomas, B.F.
LC assay of urinary estriol and electrochemical detection with a battery powered detector (224) 99
Shimada, K., Tanaka, M., Nambara, T., Imai, Y., Abe, K. and Yoshinaga, K.
Determination of captopril in human blood by HPLC with electrochemical detection (227) 445
Shimada, K., Tanaka, T. and Nambara, T.
Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33
Shimizu, H., see Suzuki, M. (221) 361
Shinka, T., see Haraguchi, S.-I. (227) 1, (230) 7
Shinkai, K., see Akedo, H. (227) 262
Shinohara, Y., Miller, R.D. and Castagnoli, N. Jr.
Ion-pair HPLC assay of 4-aminopyridine in serum (230) 363
Shintani, S., see Ohya, K. (221) 67, (221) 293
Shiobara, Y., see Kamimura, H. (229) 337
Shiraishi, K., see Okazaki, M. (223) 285
Shirasawa, K., see Murakami, K. (227) 103
Shiu, G.K. and Nemoto, E.M.
Simple, rapid and sensitive reversed-phase HPLC method for thiopental and pentobarbital determination in plasma and brain tissue (227) 207
Shroot, B., see Caron, J.C. (230) 125
Shum, A., Sole, M.J. and Van Loon, G.R.
Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123
Sidó, T., see Sajgó, M. (222) 303
Siegler, R.W., see Riley, C.M. (229) 373
Silink, M., see Kreutzmann, D.J. (228) 95
Silva, J.A.F. de, see De Silva, J.A.F. (221) 97, (223) 111, (224) 275, (226) 135, (229) 167
Silver, H.K.B., Karim, K.A., Gray, M.J. and Salinas, F.A.
HPLC quantitation of N-acetylneuraminic acid in malignant melanoma and breast carcinoma (224) 381
Simmonds, R.J. and Harkness, R.A.
HPLC methods for base and nucleoside analysis in extracellular fluids and in cells (226) 369
Simona, M.G. and Grandjean, E.M.
Simple HPTLC method for the determination of disopyramide and its mono-N-dealkylated metabolite in serum (224) 532
Sims, P., Truscott, R. and Halpern, B.
Improved procedure for the anion-exchange isolation of urinary organic acids (222) 337
Sioufi, A., Godbillon, J. and Caudal, F.
HPLC determination of sulphathiazole in human plasma and urine (221) 419
Sioufi, A. and Pommier, F.
GC determination of clioquinol (Vioform) in human plasma (226) 219

Sioufi, A. and Pommier, F.
GC determination of isosorbide dinitrate in human plasma and urine (229) 347
Sioufi, A., Pommier, F., Mangoni, P., Gauron, S. and Metayer, J.-P.
GC determination of phentolamine (Regitine) in human plasma and urine (222) 429
Sioufi, A. and Richard, A.
GC determination of maprotiline and its N-desmethyl metabolite in human blood using nitrogen detection (221) 393
Sioufi, A., Sandrenan, N. and Pommier, F.
GC determination of N-desmethyl-propyphenazone, a metabolite of propyphenazone, in human urine (221) 377
Sipes, I.G., see Gillespie, T.J. (223) 95
Sistovaris, N., see Uihlein, M. (227) 93
Sjöqvist, F., see Svensson, J.-O. (230) 427
Sjövall, J., see Axelson, M. (224) 355
Skinner, S.J.M.
Rapid method for the purification of the elastin cross-links, desmosine and isodesmosine (229) 200
Skobolo, R., see Schöneshöfer, M. (222) 478
Skofitsch, G., Saria, A., Holzer, P. and Lembeck, F.
Histamine in tissue: determination by HPLC after condensation with *o*-phthaldialdehyde (226) 53
Slack, J.A.
Determination of isoxepac in plasma by HPLC (221) 431
Slattery, J.T., see Wilson, J.M. (227) 453
Slemr, J., see Edenharder, R. (222) 1
Slikker, W. Jr., Lipe, G.W. and Newport, G.D.
HPLC analysis of estradiol-17β and metabolites in biological media (224) 205
Slikker, W. Jr., see Althaus, Z.R. (227) 11
Smit, A.L.C. Jr., see Ching, N.P.H. (222) 171
Smith, A.I., see McDermott, J.R. (222) 371
Smith, C.V., see Lauterburg, B.H. (224) 431
Smith, D.L., see Gelijkens, C.F. (225) 291
Smith, M., see Beales, D. (226) 498
Smith, R.G. and Cheung, L.K.
Determination of two nitrosourea antitumor agents by chemical ionization GC-MS (229) 464
Smith, R.G., see Benvenuto, J.A. (222) 518
Smith, R.V., see Velagapudi, R. (228) 423
Smith, S.L., Novotný, M., Moore, S.A. and Felten, D.L.
Studies of hyperlipidemia in drug-induced diabetic rats by HPLC (221) 19
Smith, S.L., Novotný, M.V. and Karmen, A.
Myo-inositol levels in the cerebrospinal fluid of infants (223) 173
Snelleman, C., see Sorel, R.H.A. (221) 129, (222) 241
Socha, J., see Schneider, Z. (229) 77
Sofia, R.D., see Hartman, C.A. (226) 510
Sofia, R.D., see Valia, K.H. (221) 170
Sokoloski, T.D., see Wu, C.C. (228) 333
Sole, M.J., see Shum, A. (228) 123
Sommadossi, J.P. and Cano, J.P.
Determination of a novel fluoropyrimidine, 5'-deoxy-5-fluorouridine, in plasma by HPLC (225) 516
Sommadossi, J.P., Lemar, M., Necciari, J., Sumirtapura, Y., Cano, J.P. and Gaillot, J.
HPLC method for the determination of plasma and urine metapramine after dansylation (228) 205
Sonoda, T., see Takada, K. (222) 156
Sorel, R.H.A., Hulshoff, A. and Snelleman, C.
HPLC analysis of nalidixic acid and hydroxynalidixic acid in plasma with a dynamic anion-exchange system (221) 129
Sorel, R.H.A., Snelleman, C. and Hulshoff, A.
HPLC analysis of nitroxoline in plasma and urine (222) 241
Sörgel, F., see Buskin, J.N. (230) 454
Soto, E.F., see Bizzozero, O. (227) 33
Soubeyrand, J., see Bannier, A. (227) 213
Souppart, C., see Lecaillon, J.B. (228) 257
Speek, A.J., Van Schaik, F., Schrijver, J. and Schreuers, W.H.P.
Determination of the $B_2$ vitamer flavin-adenine dinucleotide in whole blood by HPLC (228) 311
Spindel, E., Pettibone, D., Fisher, L., Fernstrom, J. and Wurtman, R.
Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381
Spiteller, G., see Grupe, A. (226) 301, (230) 335
Spiteller, G., see Pfeifer, P. (223) 21
Spiteller, G., see Pinkston, D. (223) 1
Staiger, M.A., Nguyen-Dinh, P. and Churchill, F.C. II
Sensitive HPLC analysis for chloroquine in body fluids. Application to studies of drug resistance in *Plasmodium falciparum* (225) 139
Stárka, L., see Justová, V. (230) 319
Stecher, W., see Schwarz, S. (223) 253
Stegnjaich, S., see Moro, E. (230) 207
Steinbach, D., see Stüber, W. (227) 193
Stempel, L.E., see Ng, K.J. (228) 43
Stephens, R.J., see Thomas, D.W. (225) 433
Sternson, L.A., Patton, T.F. and King, T.B.
HPLC analysis of miconazole in plasma (227) 223
Sternson, L.A., see Reynolds, D.L. (222) 225
Sternson, L.A., see Riley, C.M. (229) 373
Stevens, D.A., see Swezey, S.F. (227) 510
Stevens, P.M., see Vose, C.W. (222) 311, (225) 488
Stewart, D.J., see Benvenuto, J.A. (222) 518
Stoltenborg, J.K., see Vane, F.M. (227) 471
Stratingh, C.M., see Muskiet, F.A.J. (230) 142
Strauss, J.F., Kitchens, R.L., Patrizi, V.W. and Frenkel, E.P.
Extraction and quantification of daunomycin and doxorubicin in tissues (221) 139

Strebel, L.E., see Aravind, M.K. (221) 176
Strife, R.J., Jardine, I. and Colvin, M.
Analysis of the anticancer drugs etoposide (VP 16-213) and teniposide (VM 26) by HPLC with fluorescence detection (224) 168
Strojny, N., D'Arconte, L. and De Silva, J.A.F.
Determination of the anti-tumor agent, 10-chloro-5-(2-dimethylaminoethyl)-7H-indolo[2,3-C]quinolin-6(5H)-one in blood or plasma by HPLC (223) 111
Strojny, N., see Brooks, M.A. (229) 167
Stüber, W. and Möller, H.
Determination of benzbromarone, bromobenzarone and benzarone in plasma by GC-MS (224) 327
Stüber, W., Mutschler, E. and Steinbach, D.
Determination of ethacrynic and tienilic acid in plasma by GLC-MS (227) 193
Subbarayan, C., see Ching, N.P.H. (222) 171, (225) 196
Suckow, R.F. and Cooper, T.B.
Simultaneous determination of amitriptyline, nortriptyline and their respective isomeric 10-hydroxy metabolites in plasma by LC (230) 391
Sugawara, T., see Murayama, K. (224) 315
Sugihara, J., Imamura, T., Yanase, T., Yamada, H. and Imoto, T.
Separation of peptides by cellulose-phosphate chromatography for identification of a hemoglobin variant (229) 193
Sugimoto, K., see Sadanaga, T. (223) 243
Sumirtapura, Y., see Sommadossi, J.P. (228) 205
Sumirtapura, Y.C., see Cano, J.P. (226) 413
Sun, T.-H., see Abbott, F.S. (227) 433
Sunberg, R.J., see Chester, T.L. (225) 17
Suzuki, A., see Kundu, S.K. (224) 249
Suzuki, A., see Manabe, T. (225) 65
Suzuki, J., see Iwai, M. (225) 275
Suzuki, M., Morita, I., Kawamura, M., Murota, S.-I., Nishizawa, M., Miyatake, T., Nagase, H., Ohno, K. and Shimizu, H.
Quantitative determination of 6-keto prostaglandin $F_{1\alpha}$ in biological fluids by capillary GC-chemical ionization MS (221) 361
Suzuki, S., see Honda, S. (226) 341
Suzuki, W., see Ohya, K. (221) 67
Sved, S., see Ong, H. (229) 433
Svensson, J.-O., Rane, A., Säwe, J. and Sjöqvist, F.
Determination of morphine, morphine-3-glucuronide and (tentatively) morphine-6-glucuronide in plasma and urine using ion-pair HPLC (230) 427
Swanson, B.N. and Boppana, V.K.
Measurement of sulindac and its metabolites in human plasma and urine by HPLC (225) 123
Swendseid, M.E., see McKee, R.W. (230) 309
Swezey, S.F., Giacomini, K.M., Abang, A., Brass, C., Stevens, D.A. and Blaschke, T.F.
Measurement of ketoconazole, a new antifungal agent, by HPLC (227) 510
Swigar, M., see Jatlow, P.I. (227) 233
Szeto, D.W., see Tse, F.L.S. (226) 231
Szymanowicz, A.
Polymorphism of urinary 4-hydroxyproline-containing polypeptides (225) 55

# T

Takada, K., Nakae, H., Asada, S., Ichikawa, Y., Fukunishi, T. and Sonoda, T.
Rapid method for the HPLC determination of bredinin in human serum (222) 156
Takai, N., see Seta, K. (221) 215
Takai, T., see Honda, S. (226) 341
Takemasa, Y., see Nakajima, O. (225) 91, (229) 481
Takitani, S., see Gidoh, M. (223) 379
Tamaki, N., see Katogi, Y. (228) 404
Tambute, A., see Guyon, F. (229) 475
Tameto, K., see Sadanaga, T. (223) 243
Tamura, Z., see Miura, K. (223) 267
Tamura, Z., see Yoshida, A. (227) 162
Tan, Y., see Kerremans, A.L.M. (229) 129
Tanaka, I., see Seta, K. (221) 215
Tanaka, M., see Shimada, K. (227) 445
Tanaka, T., see Shimada, K. (223) 33
Tang, D.-S., see Muir, K.T. (221) 85
Tangney, C.C., McNair, H.M. and Driskell, J.A.
Quantitation of individual tocopherols in plasma, platelets, lipids, and livers by HPLC (224) 389
Tanimura, T., see Uchikura, K. (223) 41
Tasker, R.A.R. and Nakatsu, K.
Rapid, reliable and sensitive assay for warfarin using normal-phase HPLC (228) 346
Taskinen, J., see Ottoila, P. (226) 488
Tateyama, T., see Kuwada, M. (222) 507
Taylor, G. and Houston, J.B.
Simultaneous determination of promethazine and two of its circulating metabolites by HPLC (230) 194
Tedeschi, G., see Riva, R. (225) 219
Tekenbergs, L., see Eriksson, B.-M. (223) 401
Tenku, A., see Kakita, K. (222) 33
Tenni, R., see Kuttan, R. (223) 182
Terao, J., see Katogi, Y. (228) 404
Terasawa, M., see Haraguchi, S.-I. (230) 7
Terheggen, H.G., see Marescau, B. (224) 185
Tettenhorst, W.C., see Chester, T.L. (225) 17
Thijssen, H.H.W., see Schlöbe, R. (230) 212
Thoanes, H., see Bellon, G. (230) 420
Thomas, B.F., see Shihabi, Z.K. (224) 99
Thomas, D.W., Parkhurst, R.M., Negi, D.S., Lunan, K.D., Wen, A.C., Brandt, A.E. and Stephens, R.J.

Improved assay for α-tocopherol in the picogram range, using GC-MS (225) 433
Thomas, E.W.
Analysis of buflomedil in mouse, rat and rabbit plasma by reversed-phase ion-pair HPLC (228) 387
Thomas, M.J., see Cannell, G.R. (226) 492
Tieckelmann, H., see Lee, G.J.-L. (222) 23
Tillement, J.P., see Houin, G. (223) 351
Timm, U. and Weidekamm, E.
Determination of pyrimethamine in human plasma after administration of Fansidar or Fansidar-mefloquine by means of HPLC with fluorescence detection (230) 107
Timm, U., Zell, M. and Herzfeld, L.
Sensitive GLC method for the determination of oxiconazole in plasma (229) 111
Tinguely, D., Baumann, P. and Schöpf, J.
Microprocedure to determine the polymorphic forms of acid $\alpha_1$-glycoprotein in plasma. Application to depressive patients treated with amitriptyline (229) 319
Todoriki, H., see Hayashi, T. (224) 197
Tokunaga, K., see Kubo, H. (227) 244
Tombe, P.P. de, see De Tombe, P.P. (229) 498, (230) 131
Tonelli, D., see Gattavecchia, E. (224) 465
Tontodonati, R., see Hart, S.J. (225) 387
Tonzetich, J., see Kostelc, J.G. (226) 315
Toren, E.C. Jr., see Vacik, D.N. (228) 1
Toro, M. di, see Di Toro, M. (224) 257
Torosian, G., see Fleitman, J. (229) 255
Toselli, A., see Ratti, B. (225) 526
Toshima, H., see Haraguchi, S.-I. (227) 1, (230) 7
Tournet, M.-C., Girre, C. and Fournier, P.E.
Simultaneous quantitation of glafenine and its major metabolites, glafenic and hydroxyglafenic acid, by HPLC (224) 348
Toutain, P.L., see Alvinerie, M. (222) 308
Towell, J.F. and Erwin, V.G.
Determination of the primary metabolite of central nervous system norepinephrine, 3-methoxy-4-hydroxyphenethyleneglycol, in mouse brain and brain perfusate by HPLC with electrochemical detection (223) 295
Trafford, D.J.H., Seamark, D.A., Turnbull, H. and Makin, H.L.J.
High-performance liquid chromatography of 25-hydroxyvitamin $D_2$ and 25-hydroxyvitamin $D_3$ in human plasma. Use of isotachysterols and a comparison with GC-MS (226) 351
Trocewicz, J., Oka, K. and Nagatsu, T.
Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407
Trostmann, A.F., see Larsen, H.F. (226) 484
Truscott, R., see Sims, P. (222) 337
Tryfiates, G.P. and Sattsangi, S.
Separation of vitamin $B_6$ compounds by paired-ion HPLC (227) 181
Tsao, C.S. and Salimi, S.L.
Ultramicromethod for the measurement of ascorbic acid in plasma and white blood cells by HPLC with electrochemical detection (224) 477
Tschöpe, W., Brenner, R. and Ritz, E.
Isotachophoresis for the determination of oxalate in unprocessed urine (222) 41
Tschöpe, W. and Ritz, E.
Isotachophoretic determination of urinary citrate in native urine (221) 59
Tse, F.L.S. and Szeto, D.W.
Reversed-phase HPLC determination of caffeine and its N-demethylated metabolites in dog plasma (226) 231
Tsuchida, I., see Niwa, T. (225) 1
Tsuji, A., see Kawasaki, T. (226) 1
Tsukada, T., see Matsumura, K. (230) 148
Tsuruta, Y., Kohashi, K. and Ohkura, Y.
Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human urine and rat brain by HPLC with fluorescence detection (224) 105
Tsutsumi, J., see Kuwada, M. (222) 507
Tsutsumi, S., see Gidoh, M. (223) 379
Turcant, A., Premel-Cabic, A., Cailleux, A. and Allain, P.
Micromethod for automated identification and quantitation of fifteen barbiturates in plasma by GLC (229) 222
Turnbull, H., see Trafford, D.J.H. (226) 351
Turnell, D.C., see Cooper, J.D.H. (227) 158
Turner, A. and Warnock, D.W.
Determination of miconazole in human saliva using HPLC (227) 229
Turner, C.E., see Jones, A.B. (226) 99

# U

Uchikura, K., Horikawa, R., Tanimura, T. and Kabasawa, Y.
Determination of catecholamines by radio-enzymatic assay using ion-pair LC (223) 41
Ueno, N., see Murakami, K. (225) 329
Ueno, T., see Murakami, K. (227) 103
Uihlein, M. and Sistovaris, N.
HPLC and TLC determination of human serum glibenclamide at therapeutic levels (227) 93
Uji, A., Kawai, S. and Nagatsu, T.
GC determination of monoamine oxidase activity using mixed substrates (221) 155
Umeno, Y., Nakai, K., Matsushima, E. and Marunaka, T.
GC-MF determination of homopantothenic acid in plasma (226) 333
Umeno, Y., see Marunaka, T. (221) 382
Umeno, Y., see Matsushima, E. (230) 188

Undrum, T., Lunde, H. and Gjessing, L.R.
Determination of ophidine in human urine (227) 53
Upton, R.A., see Buskin, J.N. (230) 438, (230) 443, (230) 454
Upton, R.A., see Kunitani, M.G. (224) 156
Uzan, A., see Jozefczak, C. (230) 87

# V

Vacik, D.N. and Toren, E.C. Jr.
Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1
Valia, K.H., Hartman, C.A., Kucharczyk, N. and Sofia, R.D.
Simultaneous determination of dyphylline and theophylline in human plasma by HPLC (221) 170
Valzelli, G., see Moro, E. (230) 207
Van Aerde, P., Moerman, E., Van Severen, R. and Braeckman, P.
Determination of plasma theophylline by straight-phase HPLC: elimination of interfering caffeine metabolites (222) 467
Van Baal, J.M., see De Abreu, R.A. (227) 45, (227) 526, (229) 67
Van Boxtel, C.J., see Oosterhuis, B. (226) 259
Van den Berg, M., see Oosterhuis, B. (226) 259
Van der Kleijn, E., see Vree, T.B. (224) 519
Van Ginneken, C.A.M., see Kerremans, A.L.M. (229) 129
Van Loon, G.R., see Shum, A. (228) 123
Van Rooy, H.H., Vermeulen, N.P.E. and Bovill, J.G.
The assay of fentanyl and its metabolites in plasma of patients using GC with alkali flame ionisation detection and GC-MS (223) 85
Van Rooy, H.H., see De Neef, J.H. (224) 133
Van Rooy, H.H., see De Wildt, D.J. (225) 381
Van Schaik, F., see Speek, A.J. (228) 311
Van Severen, R., see Van Aerde, P. (222) 467
Van 't Hoff, W.G., see Barnicoat, A.J. (225) 236
Vane, F.M., Stoltenborg, J.K. and Buggé, C.J.L.
Determination of 13-*cis*-retinoic acid and its major metabolite, 4-oxo-13-*cis*-retinoic acid, in human blood by reversed-phase HPLC (227) 471
Varteni-Farina, M., see Della Ragione, F. (226) 243
Vasiliades, J. and Sahawneh, T.
Midazolam determination by GC, LC and GC-MS (228) 195
Vasiliades, J. and Sahawneh, T.H.
Determination of midazolam by HPLC (225) 266
Vedrine, F., see Chabard, J.L. (221) 9
Veenendaal, J.R. and Meffin, P.J.
The simultaneous analysis of clofibric acid and probenecid and the direct analysis of clofibric acid glucuronide by HPLC (223) 147
Veening, H., see Mohammed, H.Y. (226) 471
Velagapudi, R., Smith, R.V., Ludden, T.M. and Sagraves, R.
Simultaneous determination of chloramphenicol and chloramphenicol succinate in plasma using HPLC (228) 423
Vereczkey, L., see Klebovich, I. (221) 403
Verghese, C., see Shand, D.G. (224) 343
Vergin, H., see Bishop-Freudling, G.B. (224) 301
Verheggen, T., see Oerlemans, F. (225) 369
Vermeulen, N.P.E., see Heusler, H. (226) 403
Vermeulen, N.P.E., see Van Rooy, H.H. (223) 85
Verweij, A. and Kientz, C.E.
HPLC separation, isolation and identification of 1,2,3-thiadiazole-5-carboxaldoxime glucuronide in rabbit urine (226) 165
Vessman, J., Johansson, M. and Gröningsson, K.
Determination of moroxydine in biological fluids by electron-capture GC (229) 227
Viala, A., Deturmeny, E., Estadieu, M., Durand, A. and Cano, J.P.
Determination of chloroquine in blood by GC with nitrogen-selective detection using an internal standard (224) 503
Vicentini, L., see Cassara, G. (221) 279
Vijayakumar, R., Murdoch, S., Deheny, T. and Walters, W.A.W.
Radioimmunoassay of prostaglandin $F_{2\alpha}$ using Sephadex G-75 gel chromatography (225) 47
Virtanen, P. and Paunio, K.
The effect of pH and ionic strength on the electrophoretic separation of acidic glycosaminoglycans (221) 49
Visseaux, M.N., see Biou, D. (226) 477
Viswanathan, C., see Rapaka, R.S. (227) 463
Vliegenthart, J.F.G., see Kamerling, J.P. (222) 276
Voisin, C., see Baune, A. (223) 219
Von Bahr, C., see Hermansson, J. (221) 109, (227) 113
Von Henning, H., see Pinkston, D. (223) 1
Vose, C.W., Muirhead, D.C., Evans, G.L., Stevens, P.M. and Burford, S.R.
Quantitation of metolazone in plasma and urine by HPLC with fluorescence detection (222) 311
Vose, C.W., Stevens, P.M. and Dickens, J.P.
LC analysis of an antimicrobial 5-nitroimidazolyl-2-sulphide derivative in biological fluids (225) 488
Vovopivec, P., see Vukušić, I. (222) 324
Vree, T.B., Baars, A.M., Hekster, Y.A. and Van der Kleijn, E.
Simultaneous determination of

chlordiazepoxide and its metabolites in human plasma and urine by means of reversed-phase HPLC (224) 519
Vries, J.X. de, see De Vries, J.X. (221) 161
Vukušić, I. and Vovopivec, P.
Quantitative TLC determination of ticrynafen in plasma of the dog (222) 324

# W

Wachter, H., see Hausen, A. (227) 61
Wadman, S.K., see Kamerling, J.P. (222) 276
Wagner, J., Danzin, C. and Mamont, P.
Reversed-phase ion-pair LC procedure for the simultaneous analysis of S-adenosylmethionine, its metabolites and the natural polyamines (227) 349
Wagner, J.G., see Bergstrom, R.F. (222) 445
Wahrgren, P., see Kinberger, B. (224) 449
Walker, S.E. and Coates, P.E.
HPLC method for determination of gentamicin in biological fluids (223) 131
Walmsley, L.M. and Chasseaud, L.F.
HPLC determination of lorazepam in monkey plasma (226) 155
Walmsley, L.M., Chasseaud, L.F. and Miller, J.N.
Determination of bumetanide in the plasma of non-human primates by HPLC (226) 441
Walmsley, L.M., see Brodie, R.R. (226) 526
Walsh, J.H., see Reeve, J. Jr. (229) 57
Walters, W.A.W., see Vijayakumar, R. (225) 47
Wang, Y.-M., see Howell, S.K. (227) 174
Wanwimolruk, S., see Rumble, R.H. (225) 252
Ward, L.C., Miller, M. and Hawgood, S.
Simple and rapid HPLC method for the quantification of 3-methylhistidine (223) 417
Warnock, D.W., see Turner, A. (227) 229
Warsh, J.J., Chiu, A. and Godse, D.D.
Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131
Washitake, M., see Seta, K. (221) 215
Wassner, S.J. and Li, J.B.
HPLC separation of six essential amino acids and its use as an aid in the diagnosis of branched-chain ketoaciduria (227) 497
Watanabe, H., Nakano, S. and Ogawa, N.
Selected ion monitoring of apomorphine in whole rat brain (229) 95
Watanabe, K., see Matsumura, K. (222) 53, (230) 148
Watkins, G.M., see Whelpton, R. (228) 321
Watson, H.M., see Roberts, M.S. (226) 175
Watts, M.T., Raisys, V.A. and Bauer, L.A.
Determination of thiabendazole and 5-hydroxythiabendazole in human serum by fluorescence-detected HPLC (230) 79
Weatherley, B.C., see Jones, C.R. (224) 492
Webb, J.W., see Rehman, A. (228) 103
Weber, B., see Schöneshöfer, M. (227) 492
Weidekamm, E., see Timm, U. (230) 107
Weinfeld, R.E. and Miller, K.F.
Determination of the major urinary metabolite of flurazepam in man by HPLC (223) 123
Weinkam, R.J., see Shiba, D.A. (229) 397
Weiss, A.J., see Gorsen, R.M. (221) 309
Weisweiler, P., see Schwandt, P. (225) 185
Welch, R.M., see DeAngelis, R.L. (221) 353
Wen, A.C., see Thomas, D.W. (225) 433
Wermeille, M.M. and Huber, G.A.
GLC determination of free dimethindene in human serum and urine at low concentrations (228) 187
Wesley-Hadzija, B. and Mattocks, A.M.
TLC determination of furosemide and 4-chloro-5-sulfamoyl anthranilic acid in plasma and urine (229) 425
Wessels, H., see Linssen, P. (223) 371
Westerink, B.H.C., see Feenstra, M.G.P. (230) 271
Westerlund, D., see Magnusson, O. (221) 237
Whelpton, R., Curry, S.H. and Watkins, G.M.
Analysis of plasma trifluoperazine by GC and selected ion monitoring (228) 321
White, J.G., see Rao, G.H.R. (226) 466, (229) 205
White, M.H., see Chambers, D.M. (225) 231
Wiegand, U.-W., Meinertz, T., Kasper, W. and Jähnchen, E.
Determination of alinidine in human plasma by HPLC (223) 238
Wientjes, M.G., see Au, J.L.-S. (228) 245
Wiesler, D., see Rhodes, G. (228) 33
Wightman, R.M., see Caudill, W.L. (227) 331
Wildt, D.J. de, see De Wildt, D.J. (225) 381
Wildtová, Z., see Justová, V. (230) 319
Wilfling, M., see Breithaupt, H. (230) 97
Wilkinson, P.A., see Brown, J.E. (226) 521
Williams, C. II, Huang, C.S., Erb, R. and Gonzalez, M.A.
HPLC assay for plasma dipyridamole monitoring (225) 225
Williams, C.H., see Davis, T.P. (228) 113
Williams, N.E., see Chan, K. (223) 213
Williams, R.L., see Buskin, J.N. (230) 438, (230) 443, (230) 454
Willis, J.V., see Jack, D.B. (223) 484
Wilson, I.D., see Beales, D. (226) 498
Wilson, J.M., Slattery, J.T., Forte, A.J. and Nelson, S.D.
Analysis of acetaminophen metabolites in urine by HPLC with UV and amperometric detection (227) 453
Wilson, J.T., see Kearns, G.L. (226) 183
Wilson, M., see Jacob, P. III (222) 61
Wilson, M.R., see Ellin, R.I. (228) 235
Wilson, N., see Kuttan, R. (223) 182
Winkler, H., Ried, W. and Lemmer, B.
HPLC method for the quantitative analysis of the aryloxypropanolamines propranolol,

metoprolol and atenolol in plasma and tissue (228) 223
Wisser, H., see Ratge, D. (230) 47
Witter, J.P., Gatley, S.J. and Balish, E.
HPLC analysis of nitrate in urine and feces (229) 450
Wittfoht, W., see Nau, H. (226) 69
Wladichuk, D., see Abbott, F.S. (227) 433
Woestenborghs, R., Michielsen, L. and Heykants, J.
Determination of levamisole in plasma and animal tissues by GC with thermionic specific detection (224) 25
Woestenborghs, R., Michielsen, L. and Heykants, J.
Rapid and sensitive GC method for the determination of alfentanil and sufentanil in biological samples (224) 122
Wolf, M.E., see Mosnaim, A.D. (224) 481
Wong, P.Y.-K., see Pieroni, J.P. (230) 115
Wong, S.H.Y., McCauley, T. and Kramer, P.A.
Determination of 2-hydroxydesipramine by HPLC (226) 147
Wright, D.J., see Boux, L.J. (227) 149
Wu, C.C., Sokoloski, T.D., Burkman, A.M., Blanford, M.F. and Wu, L.S.
Methods for the quantitation of nitroglycerin and its metabolites in human plasma (228) 333
Wu, L.S., see Wu, C.C. (228) 333
Wurtman, R., see Spindel, E. (222) 381

# Y

Yajima, T., see Yuki, H. (221) 271
Yamada, F., see Okamoto, M. (223) 473
Yamada, H., see Sugihara, J. (229) 193
Yamada, K., Kayama, E., Aizawa, Y., Oka, K. and Hara, S.
Determination of vanillylmandelic acid in urine by pre-column dansylation using micro HPLC with fluorescence detection (223) 176
Yamada, T., see Reeve, J. Jr. (229) 57
Yamaguchi, T., see Maeda, T. (223) 357
Yamaguchi, Y.
Enzymatic determination of urinary 17β-hydroxysteroids on thin-layer chromatograms (228) 317
Yamaguchi, Y. and Hayashi, C.
Enzymatic detection of urinary conjugated steroids after gel chromatography (230) 263
Yamaguchi, Y., Hayashi, C. and Miyai, K.
Enzymatic detection of urinary acidic 3α-hydroxysteroids on thin-layer chromatograms (224) 111
Yamamoto, K., see Idzu, G. (229) 327
Yamanaka, N., see Ohki, T. (228) 51
Yamashita, I., see Miyazaki, K. (223) 449
Yanaihara, T., see Iwai, M. (225) 275
Yanase, T., see Sugihara, J. (229) 193
Yandrasitz, J.R., Berry, G. and Segal, S.
HPLC of phospholipids with UV detection: optimization of separations on silica (225) 319
Yasue, T., see Okamoto, M. (223) 473
Yasueda, H., see Mita, H. (221) 1
Yeager, R.L., see Narang, P.K. (230) 373
Yee, Y.-G. and Kates, R.E.
HPLC analysis of lorcainide and its active metabolite, norlorcainide, in human plasma (223) 454
Yeh, S.Y., see Cone, E.J. (230) 57
Yeung, D., Nacht, S. and Gans, E.H.
HPLC determination of free resorcinol in plasma and in urine (224) 513
Yeung, E.S., see Kuo, J.C. (223) 321, (229) 293
Yoshida, A., Yoshioka, M., Sakai, T. and Tamura, Z.
Simple method for the determination of homovanillic acid and vanillylmandelic acid in urine by HPLC (227) 162
Yoshida, T., see Konaka, R. (225) 169
Yoshida, Y., see Nakajima, O. (225) 91, (229) 481
Yoshinaga, K., see Shimada, K. (227) 445
Yoshioka, M., see Yoshida, A. (227) 162
Young, J.G., see Anderson, G.M. (222) 112, (223) 315, (228) 155
Young, S.N., see Anderson, G.M. (223) 315, (228) 155
Yousefnejad, D., see Cone, E.J. (223) 331
Yuki, H., Hirano, N., Kawasaki, H. and Yajima, T.
Analysis of serum iron by gel permeation HPLC (221) 271

# Z

Zacest, R., see Reece, P.A. (225) 151
Zaitsu, K., see Nohta, H. (227) 415
Zakaria, M. and Brown, P.R.
HPLC of nucleotides, nucleosides and bases (Review) (226) 267
Zakaria, M. and Brown, P.R.
Rapid assay for plasma chlorambucil and phenyl acetic mustard using reversed-phase LC (230) 381
Zanelli, J.M., O'Hare, M.J., Nice, E.C. and Corran, P.H.
Purification and assay of bovine parathyroid hormone by reversed-phase HPLC (223) 59
Zappia, V., see Della Ragione, F. (226) 243
Zecca, L., Guadagni, L. and Bareggi, S.R.
Determination of sodium flavodate in body

fluids by HPLC. Application to clinical pharmacokinetic studies (230) 168
Zell, M., see Timm, U. (229) 111
Zelson, P.R., see Kostelc, J.G. (226) 315
Zerilli, L.F., see Ratti, B. (225) 526
Zimmer, A., see Roth, W. (222) 13
Zodda, J.P., Heineman, W.R., Gilbert, T.W. and Deutsch, E.
Quantitative determination of pertechnetate by HPLC with UV detection (227) 249
Zumbo, A.M., see Abernethy, D.R. (223) 432
Zuspan, F.P., see Ng, K.J. (228) 43
Zvirblis, P., see Ellin, R.I. (228) 235
Zwaan, C.L., see Barends, D.M. (222) 316
Zwaan, C.L., see Barends, DM. (225) 417

# SUBJECT INDEX

## VOLUMES 221—230

*The volume number is placed in parentheses before the page number*

# A

**Acebutolol**
HPLC assay of acebutolol and two of its metabolites in plasma and urine (230) 438

**Acetaminophen**
Rapid GC method for emergency determination of paracetamol in human serum (221) 414

Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479

Reversed-phase chromatography of urinary metabolites of paracetamol using ion suppression and ion pairing (225) 387

HPLC determination of acetaminophen in plasma: single-dose pharmacokinetic studies (226) 224

Analysis of acetaminophen metabolites in urine by HPLC with UV and amperometric detection (227) 453

Simultaneous determination of acetaminophen, theophylline and salicylate in serum by HPLC (229) 492

Improved acetaminophen assay sensitivity by modification of a HPLC technique (230) 443

**Acetazolamide**
Efficient extraction and reversed-phase HPLC-UV quantitation of acetazolamide in serum (225) 231

**Acetoacetate**
Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455

**Acetylaminophenazone**
Rapid TLC of various weak analgesics in saliva (225) 427

**Acetylaminosalicylic acid**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

**Acetylcysteine**
Reversed-phase ion-pair HPLC of mercaptoacetate and N-acetylcysteine after derivatization with N-(1-pyrene)maleimide and N-(7-dimethylamino-4-methyl-3-coumarinyl)maleimide (229) 409

**Acetyldopamine**
Determination of N-acetyldopamine by LC with electrochemical detection (224) 310

**Acetylhydrazine**
Determination of isoniazid and its hydrazino metabolites, acetylisoniazid, acetylhydrazine, and diacetylhydrazine in human plasma by GC-MS (224) 431

**Acetylisoniazid**
Determination of isoniazid and its hydrazino metabolites, acetylisoniazid, acetylhydrazine, and diacetylhydrazine in human plasma by GC-MS (224) 431

Hydrophilic ion-pair reversed-phase HPLC for the simultaneous assay of isoniazid and acetylisoniazid in serum: a microscale procedure (226) 250

**Acetylmethadol**
Determination of acetylmethadol and metabolites by use of HPLC (222) 81

**Acetylneuraminic acid**
HPLC quantitation of N-acetylneuraminic acid in malignant melanoma and breast carcinoma (224) 381

**Acetylsalicylamide**
Rapid TLC of various weak analgesics in saliva (225) 427

**Acetylsalicylic acid**
Determination of aspirin and its major metabolites in plasma by HPLC without solvent extraction (225) 252

**Acetylserotonin**
Determination of indoles in human and rat pineal (228) 155

**Acetylsulphamethoxazole**
Simultaneous determination of trimethoprim, sulphamethoxazole and $N^4$-acetylsulphamethoxazole in serum and urine by HPLC (223) 139

**Acid glycoprotein**
Microprocedure to determine the polymorphic forms of acid $\alpha_1$-glycoprotein in plasma. Application to depressive patients treated with amitriptyline (229) 319

**Acid metabolites**
New polar acid metabolites in human urine (226) 301

**Acyclovir**
Simple HPLC method for the analysis of 9-(2-hydroxyethoxymethyl)guanine (acyclovir) in human plasma and urine (224) 51

**Adenine nucleotides**
Simultaneous determination of myocardial adenine nucleotides and creatine phosphate by HPLC (230) 131

**Adenosine**
Analysis of adenosine, inosine and hypoxanthine in suspensions of cardiac myocytes by HPLC (226) 202

Determination of adenosine in normal human plasma and serum by HPLC (227) 168

**Adenosine triphosphate**
Direct assay for creatine kinase by reversed-phase HPLC (221) 39

**Adenosylmethionine**
Reversed-phase ion-pair LC procedure for the simultaneous analysis of S-adenosylmethionine, its metabolites and the natural polyamines (227) 349

**Adrenaline**
Determination of $\alpha$-methyldopa, $\alpha$-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407

Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143

**Adriamycin**
Isotachophoretic determination of adriamycin and adriamycinol in human plasma (227) 262

**Adriamycinol**
Isotachophoretic determination of adriamycin and adriamycinol in human plasma (227) 262

**Aflatoxins**
Screening method for the detection of aflatoxin and metabolites in human urine: aflatoxins $B_1$, $G_1$, $M_1$, $B_2a$, $G_2a$, aflatoxicols I and II (227) 256

**Albumin**
A new method for detection of drug-binding proteins using a parallel-flow dialysis technique (223) 267

**Alcuronium chloride**
Determination of alcuronium chloride in biological fluids by HPLC (225) 240

**Alditol acetates**
Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

**Aldoses**
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Aldosterone**
Estimation of free urinary aldosterone and 18-hydroxycorticosterone by a combination of automatic HPLC and radioimmunoassay (227) 492

**Aldosterone antagonist**
Determination of plasma levels of spirorenone, a new aldosterone antagonist, and one of its metabolites by HPLC (230) 37

**Alfentanil**
Rapid and sensitive GC method for the determination of alfentanil and sufentanil in biological samples (224) 122

**Alinidine**
Determination of alinidine in human plasma by HPLC (223) 238

**Allopurinol**
Determination of allopurinol and oxipurinol in biological fluids by HPLC (226) 237

Simultaneous determination of allopurinol, oxipurinol and uric acid in human plasma by HPLC (228) 382

Simultaneous determination of allopurinol and oxipurinol in biological fluids by mass fragmentography (229) 211

**Alloxan**
Excretion of urinary volatile metabolites in response to alloxan induced diabetes of short duration in rats (228) 33

**Almitrine**
Trace determination of almitrine in plasma by GLC using a nitrogen-phosphorus detector (223) 219

**Alphadolone**
Sensitive GC method for the determination of alphadolone in plasma (226) 213

**Alprazolam**
Electron-capture GC analysis of the triazolobenzodiazepines alprazolam and triazolam (225) 202

**Alprenolol**
Determination of (*R*)- and (*S*)-alprenolol and (*R*)- and (*S*)-metoprolol as their diastereomeric derivatives in human plasma by reversed-phase LC (227) 113

**Amantadine**
GLC determination of plasma and urinary levels of amantadine in man (228) 327

**Amdinocillin**
Determination of amdinocillin in plasma and urine by high-performance liquid chromatography (227) 137

**Amidino penicillins**
Determination of amdinocillin in plasma and urine by high-performance liquid chromatography (227) 137

**Amines**
HPLC procedure for the simultaneous determination of the natural polyamines and their monoacetyl derivatives (221) 227

Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

Novel post-column derivatization method for the fluorimetric determination of norepinephrine and epinephrine (221) 249

Determination of rat brain tissue catecholamines using LC with electrochemical detection (222) 107

Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363

Determination of catecholamines by radioenzymatic assay using ion-pair LC (223) 41

Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51

Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198

Determination of N-acetyldopamine by LC with electrochemical detection (224) 310

HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

Determination of noncatecholic phenylethylamines and monomethylated derivatives of phenylethylamine (224) 481

Determination of α-methyldopa, α-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

Determination of a wide range of urinary amine metabolites using a simple HPLC technique (225) 179

Simultaneous extraction and separation of trace amines of biological interest (225) 301

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347

Micro HPLC system with micro precolumn and dual electrochemical detector for direct injection analysis of catecholamines in body fluids (226) 33

Reversed-phase ion-pair LC procedure for the simultaneous analysis of S-adenosylmethionine, its metabolites and the natural polyamines (227) 349

Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369

GC-MS analysis of amine metabolites in urine (228) 51

Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113

Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143

Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1

Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35

Single-step HPLC method for the analysis of acetylated polyamines (229) 47

Measurement of catecholamines in biological fluids by HPLC. A comparison of fluorimetric with electrochemical detection (229) 301

Assay and stability of α-methyldopa in man using HPLC with electrochemical detection (229) 433

MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142

Determination of α-methyldopa and methyldopate in human breast milk and plasma by ion-exchange chromatography using electrochemical detection (230) 162

Reversed-phase LC with amperometric detection of lipophilic dopamine analogues and determination of brain and serum concentrations after sample clean-up on small Sephadex G-10 columns (230) 271

**Amino acid decarboxylase**
New and highly sensitive assay for L-5-hydroxytryptophan decarboxylase activity by HPLC-voltammetry (221) 265

**Amino acids**
Glass capillary GC determination of $N^{\tau}$-methylhistidine in urine (221) 149

GC analysis of free amino acids in the hyaloplasm of the hypophysis, pineal gland, thyroid gland, spinal cord, thymus and lymph nodes of the cow (222) 345

Separation of taurine and glycerophosphoryl ethanolamine on amino acid analyzers (223) 179

Determination of γ-carboxyglutamic acid excretion in urine (223) 182

Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198

Assay of α-difluoromethylornithine in body fluids and tissues by automatic amino-acid analysis (223) 409

Simple and rapid HPLC method for the quantification of 3-methylhistidine (223) 417

Glass capillary quantitative determination of $N^{\pi}$-methylhistidine in urine and muscles (224) 94

GC determination of glutamine in the presence of glutamic acid in biological fluids (224) 177

Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315

Analysis of middle molecular peptides in normal and uremic body fluids by HPLC (224) 322

Separation and identification of dansylated human serum and urinary amino acids by two-dimensional TLC. Application to aminoacidopathies (226) 477

Fluorescence detection of cystine by *o*-phthalaldehyde derivatisation and its separation using HPLC (227) 158

Determination of urinary amino acids by means of glass capillary GLC with alkali-flame ionisation detection and flame ionisation detection (227) 305

Simplified procedure for the analysis of 3- and 4-hydroxyproline (227) 341

HPLC separation of six essential amino acids and its use as an aid in the diagnosis of branched-chain ketoaciduria (227) 497

Rapid analysis of 3-methylhistidine in urine, plasma, muscle and amniotic fluid with a single HPLC system but with different ion-pairing reagents (228) 273

Chromatographic analysis of glutamine in plasma (229) 188

Improved method for the evaluation of 3-hydroxyproline and 4-hydroxyproline in the urine (230) 420

**Aminobutyric acid**
Use of reversed-phase HPLC for simultaneous determination of glutamine synthetase and glutamic acid decarboxylase in crude extracts (225) 37

GC-MS determination of glutamic acid decarboxylase activity in subregions of rat brain (225) 283

Determination of γ-aminobutyric acid by LC with electrochemical detection (227) 331

Determination of γ-aminobutyric acid by reversed-phase HPLC and pre-column labeling for fluorescence detection (230) 121

**Aminobutyric acid agonist**
Determination of the γ-aminobutyric acid agonist 4,5,6,7-tetrahydroisoxazolo[5,4-*c*]pyridine-3-ol in urine by HPLC (228) 279

**1-(2-Aminoethyl)-3-(2,6-dichlorophenyl)thiourea**
Determination of the antihypertensive agent 1-(2-aminoethyl)-3-(2,6-dichlorophenyl)thiourea in plasma and urine by HPLC (224) 275

**Aminoglycoside antibiotics**
Determination of gentamicin in serum using LCC (221) 182

Improved microdetermination of gentamicin and sisomicin in serum by HPLC with ultraviolet detection (222) 316

HPLC method for determination of gentamicin in biological fluids (223) 131

Micro-determination of tobramycin in serum by HPLC with UV detection (225) 417

Micro determination of gentamicin in serum by HPLC (227) 244

**Aminolaevulinic acid dehydrase**
Determination of 5-aminolaevulinic acid dehydrase activity in erythrocytes by HPLC (223) 421

**Aminophenazone**
Rapid TLC of various weak analgesics in saliva (225) 427

**Aminopyridine**
Ion-pair HPLC assay of 4-aminopyridine in serum (230) 363

**Aminosalicylate**
Assay of 5-aminosalicylate and its acetylated metabolite in biological fluids by HPLC on dynamically modified silica (226) 504

**Aminosalicylic acid**
Simplified HPLC method for 5-aminosalicylic acid in plasma and urine (225) 498

**Aminosugars**
Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

**Aminothiadiazole**
LC separation and quantitation of 2-amino-1,3,4-thiadiazole (NSC-4728) from human and murine serum (230) 175

**Amitriptyline**
Simultaneous determination of amitriptyline, nortriptyline and their respective isomeric 10-hydroxy metabolites in plasma by LC (230) 391

**Amperometric detector**
Differential pulse amperometric detection of drugs in plasma using a dropping mercury electrode as a HPLC detector (222) 179

**AMP nucleotidase**
Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77

**Amylase**
Action pattern of human pancreatic α-amylase on maltoheptaose, a substrate for determining α-amylase in serum (223) 69

**Amylobarbitone**
Analysis of barbiturates in blood by HPLC (226) 117

**Analgesics**
Rapid TLC of various weak analgesics in saliva (225) 427

**Androgens**
Combined PC and reversed-phase HPLC method for the study of pregnenolone and progesterone metabolites (224) 399

**Androsterone**
Steroid profiles of healthy individuals (223) 21

**9,10-Anthracenedicarboxaldehyde bis(4,5-dihydro-1H-imidazol-2-yl hydrazone) dihydrochloride**
Reversed-phase HPLC assay for the antineoplastic agent 9,10-anthracenedicarboxaldehyde bis(4,5-dihydro-1H-imidazol-2-yl hydrazone) dihydrochloride (226) 514

**Anthracenediones**
Clinical analysis for the anti-neoplastic agent 1,4-dihydroxy-5,8-bis{[2-[(2-hydroxyethyl)aminolethyl]amino}9,10-anthracenedione dihydrochloride (NSC 301739) in plasma. Application of temperature control to provide selectivity in paired-ion HPLC (222) 225

**Anthracenes**
Reversed-phase HPLC assay for the antineoplastic agent 9,10-anthracenedicarboxaldehyde bis(4,5-dihydro-1H-imidazol-2-yl hydrazone) dihydrochloride (226) 514

**Anthracycline antibiotics**
Extraction and quantification of daunomycin and doxorubicin in tissues (221) 139

Analysis of carminomycin in human serum by fluorometric HPLC (223) 155

Rapid HPLC assay for the anthracyclines daunorubicin and 7-con-O-methylnogarol in plasma (226) 521

Isotachophoretic determination of adriamycin and adriamycinol in human plasma (227) 262

**Anthraquinone glycosides**
Determination of 4'-epidoxorubicin and its 13-dihydro derivative in human plasma by HPLC with fluorescence detection (230) 207

**Anticonvulsant drugs**
Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

**Antidepressant drugs**
Evaluation of $C_{18}$ Sep-Pak cartridges for biological sample clean-up for tricyclic antidepressant assays (225) 189

**Antioxidants**
Determination of the antioxidant 3-*tert.* -butyl-4-hydroxyanisole in rat plasma using high-resolution GC-MS (225) 83

**Antipyrine**
Antipyrine determination in human plasma by GLC using nitrogen-phosphorus detection (223) 432

**Antrafenine**
Determination of antrafenine and its main acid metabolite, 2-{[7-(trifluoromethyl)-4-quinolinyl]amino}-benzoic acid, in biological fluids using HPLC with large volume automatic injection and GLC with derivative formation (223) 103

**Apolipoproteins**
Separation of human C-apolipoproteins by HPLC (225) 185

**Apomorphine**
Selected ion monitoring of apomorphine in whole rat brain (229) 95

**Arabinofuranosylcytosine**
Simultaneous determination of 1-β-D-arabinofuranosylcytosine 5'-triphosphate and 3-deaza-

uridine 5'-triphosphate in human leukemia cells by HPLC (221) 425

Determination of 1-β-D-arabinofuranosyl-cytosine and 1-β-D-arabinofuranosyluracil in human plasma by HPLC (223) 371

**Arabinofuranosyluracil**
Determination of 1-β-D-arabinofuranosyl-cytosine and 1-β-D-arabinofuranosyluracil in human plasma by HPLC (223) 371

**Arildone**
The analysis of arildone in plasma, urine and feces by GLC with electron-capture detection (222) 213

**Ascorbic acid**
Ultramicromethod for the measurement of ascorbic acid in plasma and white blood cells by HPLC with electrochemical detection (224) 477

**Aspirin**
Determination of aspirin and its major metabolites in plasma by HPLC without solvent extraction (225) 252

**Atenolol**
HPLC method for the quantitative analysis of the aryloxypropanolamines propranolol, metoprolol and atenolol in plasma and tissue (228) 223

**Aziridinylbenzoquinones**
LC determination of an antineoplastic aziridinylbenzoquinone in human and murine serum (222) 146

**Azlocillin**
Improved procedure for the determination of the ureidopenicillins azlocillin and mezlocillin in plasma by HPLC (228) 409

**Azodipyrroles**
Isocratic HPLC of bile pigments (227) 503

# B

**Barbiturates**
Simultaneous assay of methylphenobarbital and phenobarbital in plasma using GC-MS with selected ion monitoring (223) 426

Determination of barbiturates and some neutral drugs in serum using quartz glass capillary GC (224) 449

Analysis of barbiturates in blood by HPLC (226) 117

Simultaneous determination of blood concentrations of methohexital and its hydroxy metabolite by GC and identification of 4'-hydroxymethohexital by combined GLC-MS (226) 403

Simple, rapid and sensitive reversed-phase HPLC method for thiopental and pentobarbital determination in plasma and brain tissue (227) 207

Rapid analysis of hexobarbital and its main metabolites in mice by HPLC (228) 350

Micromethod for automated identification and quantitation of fifteen barbiturates in plasma by GLC (229) 222

Chromatographic micro-procedure for trace determination of phenobarbital in blood serum (229) 365

**Benzarone**
Determination of benzbromarone, bromo-benzarone and benzarone in plasma by GC-MS (224) 327

**Benzbromarone**
Determination of benzbromarone, bromo-benzarone and benzarone in plasma by GC-MS (224) 327

**Benzodiazepines**
GC determination of clonazepam (221) 399

Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95

Plasma concentrations of temazepam, a 3-hydroxy benzodiazepine, determined by electron-capture GLC (222) 125

Improved GC procedure for the determination of clonazepam levels in plasma using a nitrogen-sensitive detector (222) 203

Determination of tofisopam in serum by HPLC (222) 303

Determination of 1,4- and 1,5-benzodiazepines in urine using a computerized GC-MS technique (222) 409

Rapid quantitation of flurazepam and its major metabolite, N-desalkylflurazepam, in human plasma by GLC with electron-capture detection (222) 491

Determination of bromazepam by GLC and its application for pharmacokinetic studies in man (222) 501

Determination of the major urinary metabolite of flurazepam in man by HPLC (223) 123

Assay of triazolam in plasma by capillary GC (223) 438

Electron-capture GC analysis of the triazolobenzodiazepines alprazolam and triazolam (225) 202

Quantitative determination of clobazam in the plasma of epileptic patients by GLC with electron-capture detection (225) 219

Determination of midazolam by HPLC (225) 266

Determination of the anxiolytic agent 8-chloro-6-(2-chlorophenyl)-4H-imidazo-[1,5-*a*][1,4]-benzodiazepine-3-carboxamide in whole blood, plasma or urine by HPLC (226) 135

HPLC determination of lorazepam in monkey plasma (226) 155

Analysis of the metabolites of ethyl loflazepate by GC with electron-capture detection (226) 413

Assay of flunitrazepam, temazepam and desalkylflurazepam in plasma by capillary GC with electron-capture detection (227) 199

Midazolam determination by GC, LC and GC-MS (228) 195

Micro-determination of clonazepam in plasma or serum by electron-capture GLC (230) 353

**Benzofurans**
Determination of benzbromarone, bromo-benzarone and benzarone in plasma by GC-MS (224) 327

**Benzyl-P2A**
Determination of some pyridinium aldoxime compounds by means of ion-pair reversed-phase HPLC: application in biological material (225) 107

**Biguanides**
Determination of moroxydine in biological fluids by electron-capture GC (229) 227

**Bile acids**
GC and MS analysis of bile acids as trifluoro-acetyl-hexafluoroisopropyl and heptafluorobutyryl derivatives (222) 1

Studies on steroids. CLXX. Separation and determination of bile acid 3-sulfates in human bile by HPLC (226) 13

**Bile pigments**
Isocratic HPLC of bile pigments (227) 503

**Bilirubin**
Separation of bilirubin species in serum and bile by reversed-phase HPLC (226) 391

**Bilirubins**
Isocratic HPLC of bile pigments (227) 503

**Biperiden**
Determination of biperiden in human serum by glass capillary GC with isothermal splitless injection and nitrogen-sensitive detection (226) 488

**4,5-Bis-(4-methoxyphenyl)-2-(2-hydroxyethylsulfinyl)-imidazole**
Determination of plasma and urine levels of a new anti-inflammatory agent, 4,5-bis-(4-methoxyphenyl)-2-(2-hydroxymethylsulfinyl)-imidazole, by HPLC and electrochemical or UV detection (222) 71

**Body fluids**
Fully automated HPLC. A new chromatograph for pharmacokinetic drug monitoring by direct injection of body fluids (222) 13

**Bombesin**
Rapid high-yield purification of neuropeptides from canine intestinal muscle (229) 57

**Breath-hydrogen**
A new chromatographic instrument for measuring trace concentrations of breath-hydrogen (229) 259

**Bredinin**
Rapid method for the HPLC determination of bredinin in human serum (222) 156

**Bromazepam**
Determination of bromazepam by GLC and its application for pharmacokinetic studies in man (222) 501

**Bromhexine**
Assay of bromhexine in human plasma by capillary GLC with nitrogen-selective detection and selected ion monitoring (227) 71

Determination of bromhexine in human plasma and urine by HPLC (228) 392

**Bromobenzarone**
Determination of benzbromarone, bromo-benzarone and benzarone in plasma by GC-MS (224) 327

**Bromoethylcinnamamide**
Quantitation of the anticonvulsant cinromide (3-bromo-N-ethylcinnamamide) and its major plasma metabolites by TLC (221) 353

**Bromo-lasalocid**
Determination of bromo-lasalocid in plasma by HPLC with fluorimetric detection (229) 167

**Bromovincamine**
Determination of 11-bromovincamine in human plasma by HPLC (228) 413

**Bromural**
HPLC determination of bromural in serum upon hemoperfusion (223) 473

**Bucetin**
Rapid TLC of various weak analgesics in saliva (225) 427

**Buflomedil**
Analysis of buflomedil in mouse, rat and rabbit plasma by reversed-phase ion-pair HPLC (228) 387

**Bufuralol**
Determination of bufuralol and its major metabolites in plasma by HPLC (221) 327

**Bumetanide**
Determination of bumetanide in the plasma of non-human primates by HPLC (226) 441

**Butobarbitone**
Analysis of barbiturates in blood by HPLC (226) 117

**Butylhydroxyanisole**
Determination of the antioxidant 3-*tert.* -butyl-4-hydroxyanisole in rat plasma using high-resolution GC-MS (225) 83

# C

**Cadaverine**
GC-MS analysis of amine metabolites in urine (228) 51

Single-step HPLC method for the analysis of acetylated polyamines (229) 47

**Caffeine**
Simultaneous determination of theophylline and caffeine after extractive alkylation in small volumes of plasma by GC-MS (221) 166

Simultaneous microdetermination of theophylline, caffeine and phenobarbital in blood collected on paper (224) 439

Reversed-phase HPLC determination of caffeine and its N-demethylated metabolites in dog plasma (226) 231

LC determination of caffeine in biologic fluids (229) 387

**Calcium carbimide**
Determination of carbimide in plasma by GLC (222) 421

**Cannabidiol**
Determination of cannabidiol in plasma by electron-capture GC (226) 99

**Captopril**
Determination of captopril in human blood by HPLC with electrochemical detection (227) 445

**Carbamazepine**
Rapid quantitative determination of underivatized carbamazepine, phenytoin, phenobarbital and *p*-hydroxyphenobarbital in biological fluids by packed column GC (221) 75

Rapid GC method for the determination of carbamazepine and unrearranged carbamazepine-10,11-epoxide in human plasma (222) 120

Rapid microanalysis of anticonvulsants by HPTLC (222) 265

Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

Simple and rapid GLC method for estimating carbamazepine in serum (222) 496

**Carbimide**
Determination of carbimide in plasma by GLC (222) 421

**Carbohydrates**
see also sugars

Determination of carbohydrates in urine by HPLC and optical activity detection (223) 321

Estimation of sugar alcohols by GLC using a modified acetylation procedure (226) 191

Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Carbon tetrachloride**
Detection of the changes in protein distribution of rat plasma induced by carbon tetrachloride administration by means of two-dimensional electrophoresis (225) 65

**Carboxyglutamic acid**
Determination of $\gamma$-carboxyglutamic acid excretion in urine (223) 182

Simplified procedure for the analysis of 3- and 4-hydroxyproline (227) 341

**Carboxylic acids**
see also organic acids

Determination of the absolute configuration of some biologically important urinary 2-hydroxydicarboxylic acids by capillary GLC (222) 276

Improved procedure for the anion-exchange isolation of urinary organic acids (222) 337

New polar acid metabolites in human urine (226) 301

**Carminomycin**
Analysis of carminomycin in human serum by fluorometric HPLC (223) 155

**Carnosine**
Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315

**Catecholamines**
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

Novel post-column derivatization method for the fluorimetric determination of norepinephrine and epinephrine (221) 249

Determination of rat brain tissue catecholamines using LC with electrochemical detection (222) 107

Determination of dopamine, homovanillic acid and 3,4-dihydroxyphenylacetic acid in rat

brain striatum by HPLC with electrochemical detection (222) 353

Determination of catecholamines by radio-enzymatic assay using ion-pair LC (223) 41

Determination of vanillylmandelic acid in urine by pre-column dansylation using micro HPLC with fluorescence detection (223) 176

Studies on dopamine-converting enzymes in human plasma (223) 188

Determination of the primary metabolite of central nervous system norepinephrine, 3-methoxy-4-hydroxyphenethyleneglycol, in mouse brain and brain perfusate by HPLC with electrochemical detection (223) 295

Quantitative determination of 3-methoxy-4-hydroxyphenylethyleneglycol and its sulfate conjugate in human lumbar cerebrospinal fluid using LC with aperometric detection (223) 305

Determination of N-acetyldopamine by LC with electrochemical detection (224) 310

Determination of α-methyldopa, α-methyl-noradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

Determination of a wide range of urinary amine metabolites using a simple HPLC technique (225) 179

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347

Micro HPLC system with micro precolumn and dual electrochemical detector for direct injection analysis of catecholamines in body fluids (226) 33

Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

Assay of catechol O-methyltransferase activity by HPLC with electrochemical detection (226) 461

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

Simultaneous determination of 4-hydroxy-3-methoxy-phenylacetic (homovanillic) acid and other monoamine metabolites in human lumbar cerebrospinal fluid. An improved HPLC study with electrochemical detection (227) 379

Analysis of urinary 3-methoxy-4-hydroxy-phenylglycol by HPLC and electrochemical detection (227) 391

Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407

Assay for dopamine β-hydroxylase in rat serum and adrenal medulla by HPLC with fluorescence detection (227) 415

Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113

Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143

Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1

Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35

Measurement of catecholamines in biological fluids by HPLC. A comparison of fluorimetric with electrochemical detection (229) 301

Assay and stability of α-methyldopa in man using HPLC with electrochemical detection (229) 433

Determination of α-methyldopa and methyldopate in human breast milk and plasma by ion-exchange chromatography using electrochemical detection (230) 162

Reversed-phase LC with amperometric detection of lipophilic dopamine analogues and determination of brain and serum concentrations after sample clean-up on small Sephadex G-10 columns (230) 271

**Catechol oestrogens**

Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33

**Catechol O-methyltransferase**
Assay of catechol O-methyltransferase activity by HPLC with electrochemical detection (226) 461

**Catechols**
Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

**Cefaclor**
Determination of cefaclor by HPLC (228) 429

**Cefmenoxime**
A very precise HPLC procedure for the determination of cefmenoxime, a new cephalosporin antibiotic, in plasma (229) 149

**Cefotaxime**
Determination of cefotaxime and desacetyl-cefotaxime in plasma and urine by HPLC (226) 431

**Celiprolol**
Specific and sensitive assay of celiprolol in blood, plasma and urine using HPLC (230) 454

**Cephalexin**
HPLC determination of cephalexin in human plasma, urine and saliva (225) 532

**Cephalosporins**
Determination of cephalosporins in biological material by reversed-phase LCC (223) 393

HPLC determination of cephalexin in human plasma, urine and saliva (225) 532

Determination of cefotaxime and desacetyl-cefotaxime in plasma and urine by HPLC (226) 431

Determination of cefsulodin, cefotiam, cephalexin, cefotaxime, desacetyl-cefotaxime, cefuroxime and cefroxadin in plasma and urine by HPLC (228) 257

Determination of cefaclor by HPLC (228) 429

A very precise HPLC procedure for the determination of cefmenoxime, a new cephalosporin antibiotic, in plasma (229) 149

**Chlorambucil**
Rapid assay for plasma chlorambucil and phenyl acetic mustard using reversed-phase LC (230) 381

**Chloramphenicol**
Simultaneous measurement of chloramphenicol and chloramphenicol succinate by HPLC (221) 176

Simultaneous determination of chloramphenicol and its succinate ester by HPLC (223) 247

Simultaneous determination of chloramphenicol and chloramphenicol succinate in plasma using HPLC (228) 423

**Chlordiazepoxide**
Differential pulse amperometric detection of drugs in plasma using a dropping mercury electrode as a HPLC detector (222) 179

Simultaneous determination of chlordiazepoxide and its metabolites in human plasma and urine by means of reversed-phase HPLC (224) 519

**Chlormezanone**
Sensitive and selective method for the determination of chlormezanone in plasma by electron-capture GC (221) 67

**8-Chloro-6-(2-chlorophenyl)-4H-imidazo-[1,5-*a*]-[1,4]-benzodiazepine-3-carboxamide**
Determination of the anxiolytic agent 8-chloro-6-(2-chlorophenyl)-4H-imidazo-[1,5-*a*][1,4]-benzodiazepine-3-carboxamide in whole blood, plasma or urine by HPLC (226) 135

**10-Chloro-5-(2-dimethylaminoethyl)-7H-indolo[2,3-C]-quinolin-6(5H)-one**
Determination of the anti-tumor agent, 10-chloro-5-(2-dimethylaminoethyl)-7H-indolo[2,3-C]quinolin-6(5H)-one in blood or plasma by HPLC (223) 111

**Chloroprocaine**
Measurement of 2-chloroprocaine in plasma by selected ion monitoring (224) 488

**Chloroquine**
Sensitive method for the determination of chloroquine and its metabolite desethyl-chloroquine in human plasma and urine by HPLC (221) 119

Contribution to the determination of four antimalaria drugs, chloroquine, quinine, pyrimethamine and sulfadoxine, alone or in mixtures in biological fluids. II. GC (in French) (224) 332

Determination of chloroquine in blood by GC with nitrogen-selective detection using an internal standard (224) 503

Sensitive HPLC analysis for chloroquine in body fluids. Application to studies of drug resistance in *Plasmodium falciparum* (225) 139

Simultaneous determination of chloroquine and its desethyl metabolites in human plasma by GC (226) 91

Determination of chloroquine and its desethyl metabolite in plasma, red blood cells in urine by LC (229) 241

Determination of chloroquine and its de-ethyl-ated metabolites in human plasma by ion-pair HPLC (229) 248

**Chlorosulphamoyl anthranilic acid**
TLC determination of furosemide and 4-chloro-5-sulfamoyl anthranilic acid in plasma and urine (229) 425

**Chlorpheniramine**
Rapid method for the determination of chlorpheniramine in urine (223) 208

**Chlorpromazine**
Simultaneous determination of chlorpromazine and levomepromazine in human plasma and urine by HPLC using electrochemical detection (227) 103

**Cholesterol**
High-performance aqueous gel permeation chromatography of serum lipoproteins: selective detection of cholesterol by enzymatic reaction (223) 285

Rapid determination of lipids in healthy human lymphocytes (227) 423

Direct quantitation of free cholesterol in total serum lipid extracts by computer-assisted GLC (228) 268

Determination of free and esterified cholesterol in human serum by HPLC and optical activity detection (229) 293

**Cholesterol esters**
Studies of hyperlipidemia in drug-induced diabetic rats by HPLC (221) 19

**Chondroitin sulphate**
The application of HPLC in enzymatic assays of chondroitin sulfate isomers in normal human urine (222) 23

**Chromatografic profiles**
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215

High-performance pyrolysis-GC: potential for differentiating cystic fibrosis cells (221) 371

**Chromatographic profiles**
Studies of hyperlipidemia in drug-induced diabetic rats by HPLC (221) 19

High-resolution GC-MS of the methyl esters of organic acids from uremic hemofiltrates (223) 1

Steroid profiles of healthy individuals (223) 21

Analysis of profiles of conjugated steroids in urine by ion-exchange separation and GC-MS (224) 355

GC-MS profile of organic acids in urine and serum of diabetic ketotic patients (225) 1

A GC-MS study of profiles of volatile metabolites in hepatic encephalopathy (226) 291

New polar acid metabolites in human urine (226) 301

Excretion of urinary volatile metabolites in response to alloxan induced diabetes of short duration in rats (228) 33

Capillary GC-MS profiles of trimethylsilyl derivatives of organic acids from amniotic fluids of different gestational age (228) 43

Comparison of isolation methods of urinary organic acids HPLC (228) 103

Analysis of small peptides in uremic serum by HPLC (228) 292

**Chrondroitin**
Electrophoresis with two buffers in one dimension in the analysis of glycosaminoglycans on cellulose acetate strips (224) 407

**Cimetidine**
Convenient and sensitive HPLC assay for cimetidine in plasma or urine (224) 156

HPLC analysis of cimetidine in serum and urine (228) 398

Improved HPLC assay for cimetidine using ranitidine as an internal standard (229) 255

**Cinnamamides**
Quantitation of the anticonvulsant cinromide (3-bromo-N-ethylcinnamamide) and its major plasma metabolites by TLC (221) 353

**Cinnarizine**
Rapid HPLC assay of cinnarizine in human plasma (227) 521

**Cinromide**
Quantitation of the anticonvulsant cinromide (3-bromo-N-ethylcinnamamide) and its major plasma metabolites by TLC (221) 353

**Ciprofibrate**
Determination of ciprofibrate in human plasma by HPLC (227) 534

**Cisplatin**
HPLC of platinum complexes on solvent generated anion exchangers. III. Application to the analysis of cisplatin in urine using automated column switching (229) 373

**Citalopram**
Fluorescence assay of citalopram and its metabolites in plasma by scanning densitometry of thin-layer chromatograms (224) 526

Determination of the antidepressant agent citalopram and metabolites in plasma by LC with fluorescence detection (227) 129

**Citrate**
Isotachophoretic determination of urinary citrate in native urine (221) 59

**Clioquinol**
GC determination of clioquinol (Vioform) in human plasma (226) 219

HPLC determination of clioquinol and its conjugates in biological materials (229) 159

**Clobazam**
Quantitative determination of clobazam in the plasma of epileptic patients by GLC with electron-capture detection (225) 219

**Clofazimine**
Determination of three main antileprosy drugs and their main metabolites in serum by HPLC (223) 379

Determination of clofazimine in plasma by HPLC (229) 503

**Clofibric acid**
The simultaneous analysis of clofibric acid and probenecid and the direct analysis of clofibric acid glucuronide by HPLC (223) 147

**Clomethiazole**
Unexpected metabolites produced from clomethiazole (230) 335

**Clomiphene**
HPLC determination of clomiphene using post-column on-line photolysis and fluorescence detection (225) 131

**Clonazepam**
GC determination of clonazepam (221) 399

Improved GC procedure for the determination of clonazepam levels in plasma using a nitrogen-sensitive detector (222) 203

Micro-determination of clonazepam in plasma or serum by electron-capture GLC (230) 353

**Codeine**
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

**Collagens**
Separation of collagen types I and III by HPLC (222) 284

Polymorphism of urinary 4-hydroxyproline-containing polypeptides (225) 55

Simplified procedure for the analysis of 3- and 4-hydroxyproline (227) 341

High-performance gel permeation chromatography of collagens (230) 409

Improved method for the evaluation of 3-hydroxyproline and 4-hydroxyproline in the urine (230) 420

**Computer program**
Simple computer program for a low-cost desk-top calculator applied to the evaluation of GLC analyses of 17-ketosteroids and pregnanes (223) 253

**7-Con-O-methylnogarol**
Rapid HPLC assay for the anthracyclines daunorubicin and 7-con-O-methylnogarol in plasma (226) 521

**Contrathion**
Determination of N-methylpyridinium 2-aldoxime methylsulfate (Contrathion) in rat plasma and urine by copper(II)-silica ligand-exchange HPLC (229) 475

**Corticosteroids**
Reliability of the estimation of serum cortisol by HPLC (222) 478

Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221

Separation of some natural and synthetic corticosteroids in biological fluids and tissues by HPLC (227) 11

Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17α-hydroxyprogesterone from small plasma or tissue samples (228) 95

Sample purification using a $C_{18}$-bonded reversed-phase cartridge for the quantitative analysis of corticosteroids in adrenal cell cultures by HPLC or GC-MS (229) 267

Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283

**Cortisol**
Reliability of the estimation of serum cortisol by HPLC (222) 478

Separation of some natural and synthetic corticosteroids in biological fluids and tissues by HPLC (227) 11

**Cotinine**
Improved GC method for the determination of nicotine and cotinine in biologic fluids (222) 61

**C-peptide**
Gel chromatographic separation of human C-peptide and proinsulin (222) 33

**Creatine kinase**
Direct assay for creatine kinase by reversed-phase HPLC (221) 39

Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1

**Creatine phosphate**
Simultaneous determination of myocardial adenine nucleotides and creatine phosphate by HPLC (230) 131

**Cresol**
Determination of urinary hippuric acid and *o*-cresol, as indices of toluene exposure, by LC on dynamically modified silica (229) 141

**Cyclandelate**
Simultaneous determination of cyclandelate and its metabolite in human plasma by capillary column GLC (223) 365

**Cyclic AMP phosphodiesterase**
Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77

**Cyclic ribonucleotides**
HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

**Cyclizine**
Determination of cyclizine and norcyclizine in plasma and urine using GLC with nitrogen selective detection (222) 135

**Cyclobarbitone**
Analysis of barbiturates in blood by HPLC (226) 117

**Cyclodiphosphathiazenes**
Determination of pentazirinocyclodiphosphathiazene in biological fluids by HPLC (230) 188

**Cyproterone acetate**
HPLC estimation of cyproterone acetate in human plasma (226) 492

**Cystic fibrosis cells**
High-performance pyrolysis-GC: potential for differentiating cystic fibrosis cells (221) 371

**Cystine**
Fluorescence detection of cystine by *o*-phthaldehyde derivatisation and its separation using HPLC (227) 158

**Cytarabine**
Determination of cytarabine and uracil arabinoside in human plasma and cerebrospinal fluid by HPLC (225) 99

**Cytosine**
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382

# D

**Dantrolene**
Simultaneous determination of dantrolene and its metabolite, 5-hydroxydantrolene, in human plasma by HPLC (228) 404

**Daunomycin**
Extraction and quantification of daunomycin and doxorubicin in tissues (221) 139

**Daunorubicin**
Rapid HPLC assay for the anthracyclines daunorubicin and 7-con-O-methylnogarol in plasma (226) 521

**Deacetylmetipranolol**
Determination of deacetylmetipranolol in body fluids by GC-chemical-ionization MS (227) 187

**Deazauridine**
Simultaneous determination of 1-β-D-arabinofuranosylcytosine 5'-triphosphate and 3-deazauridine 5'-triphosphate in human leukemia cells by HPLC (221) 425

**Dehydroepiandrosterone**
Steroid profiles of healthy individuals (223) 21

Determination of urinary dehydroepiandrosterone sulphate by combined HPLC and radioimmunoassay (227) 485

**4'-Demethyl-epipodophyllotoxin-9-(4,6-0-ethylidene β-D-glucopyranoside)**
HPLC determination of 4'-demethylepipodophyllotoxin-9-(4,6-0-ethylidene β-D-glucopyranoside) (VP 16-213) in human plasma (222) 141

**Deoxyaldonic acids**
Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

Analysis of organic acids in the hearts of patients with idiopathic cardiomyopathy by GC-MS (230) 7

**Deoxycorticosterone**
Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17*α*-hydroxyprogesterone from small plasma or tissue samples (228) 95

**Deoxyfluorouridine**
Determination of a novel fluoropyrimidine, 5'-deoxy-5-fluorouridine, in plasma by HPLC (225) 516

Reversed-phase ion-pair HPLC assay of 5-fluorouracil, 5'-deoxy-5-fluorouridine, their nucleosides, mono-, di-, and triphosphate nucleotides with a mixture of quaternary ammonium ions (228) 245

**Deoxyribonucleosides**
HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

**Deproteinization methods**
Evaluation of the relative efficacy of various techniques for deproteinizing plasma samples prior to HPLC (226) 455

**Derivatization**
Extractive alkylation of 6-mercaptopurine and determination in plasma by GC-MS (225) 73

Fluorescence detection of cystine by *o*-phthalaldehyde derivatisation and its separation using HPLC (227) 158

**Desacetyl-Bisacodyl**
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389

**Desacetylcefotaxime**
Determination of cefotaxime and desacetylcefotaxime in plasma and urine by HPLC (226) 431

**Desacetylrifampicin**
Quantitative assay of rifampicin and its main metabolite 25-desacetylrifampicin in human plasma by reversed-phase HPLC (225) 526

**Desalkylflurazepam**
Rapid quantitation of flurazepam and its major metabolite, N-desalkylflurazepam, in human plasma by GLC with electron-capture detection (222) 491

Assay of flunitrazepam, temazepam and desalkylflurazepam in plasma by capillary GC with electron-capture detection (227) 199

**Desethylchloroquine**
Sensitive method for the determination of chloroquine and its metabolite desethylchloroquine in human plasma and urine by HPLC (221) 119

Simultaneous determination of chloroquine and its desethyl metabolites in human plasma by GC (226) 91

**Desmethylmisonidazole**
Simultaneous determination of misonidazole and desmethylmisonidazole in plasma by HPLC (225) 407

**Desmethylpropyphenazone**
GC determination of N-desmethyl-propyphenazone, a metabolite of propyphenazone, in human urine (221) 377

**Desmosine**
Rapid method for the purification of the elastin cross-links, desmosine and isodesmosine (229) 200

**Diacetylhydrazine**
Determination of isoniazid and its hydrazino metabolites, acetylisoniazid, acetylhydrazine, and diacetylhydrazine in human plasma by GC-MS (224) 431

**Diamines**
HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

**Diaminobenzylpyrimidines**
Determination of 2,4-diamino-5-benzylpyrimidines in combination with sulphadiazine in biological fluids using HPLC (224) 301

**Diaminodiphenyl sulphone**
Determination of three main antileprosy drugs and their main metabolites in serum by HPLC (223) 379

**Diazepam**
Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95

**Dibutyl phthalate**
GC-MS detection of circulating plasticizers in surgical patients (222) 171

GC quantitation of two plasticizers contaminating intravenous fluids stored in plastic containers (225) 196

**(Dichloromethylene)diphosphonate**
Determination of (dichloromethylene)diphosphonate in physiological fluids by ion-exchange chromatography with phosphorus-selective detection (225) 17

**Diclofenac**
Comments to the article "Improved GC method of determining diclofenac in plasma" (223) 484

**Diethylamine**
HPLC method for the determination of diethylamine a metabolite of disulfiram, in urine (229) 470

**Diethyldithiocarbamate**
Simultaneous determination of disulfiram and two of its dithiocarbamate metabolites in human plasma by reversed-phase LC (224) 457

**Di(ethylhexyl) phthalate**
GC-MS detection of circulating plasticizers in surgical patients (222) 171

GC quantitation of two plasticizers contaminating intravenous fluids stored in plastic containers (225) 196

**Diflunisal**
Rapid estimation of diflunisal in plasma and urine by HPLC and a comparison with a fluorometric method (229) 234

**Difluoromethylornithine**
Assay of α-difluoromethylornithine in body fluids and tissues by automatic amino-acid analysis (223) 409

**Digoxin**
Determination of tritiated digoxin and metabolites in urine by LC (223) 401

Specific assay for radiolabelled digoxin and its known apolar metabolites in biological fluids. I (224) 67

**Dihydroergot alkaloids**
Quantitative TLC determination of dihydroergot alkaloids (229) 417

**6,11-Dihydro-11-oxodibenz[*b,e*]oxepin-2-acetic acid**

Determination of 6,11-dihydro-11-dibenz[*b,e*]-oxepin-2-acetic acid (isoxepac) in plasma by HPLC (225) 482

**Dihydroxyphenylacetic acid**
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

Determination of dopamine, homovanillic acid and 3,4-dihydroxyphenylacetic acid in rat brain striatum by HPLC with electrochemical detection (222) 353

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

**Dihydroxyphenylalanine**
The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

**Dihydroxyphenylethyleneglycol**
GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

**Diisopropylphenol**
Estimation of ICI 35,868 (Diprivan) in blood by HPLC, following coupling with Gibbs' reagent (223) 232

**Dimethindene**
GLC determination of free dimethindene in human serum and urine at low concentrations (228) 187

**Dioxyanthraquinone**
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389

**Diphosphonates**
Determination of (dichloromethylene)diphosphonate in physiological fluids by ion-exchange chromatography with phosphorus-selective detection (225) 17

**Diprivan**
Estimation of ICI 35,868 (Diprivan) in blood by HPLC, following coupling with Gibbs' reagent (223) 232

**Dipyridamole**
HPLC assay for plasma dipyridamole monitoring (225) 225

**Disopyramide**
Simple HPTLC method for the determination of disopyramide and its mono-N-dealkylated metabolite in serum (224) 532

Simultaneous HPLC assay for quinidine, disopyramide and the mono-N-dealkylated metabolite of disopyramide (230) 433

**Disulphiram**
Simultaneous determination of disulfiram and two of its dithiocarbamate metabolites in human plasma by reversed-phase LC (224) 457

HPLC method for the determination of diethylamine a metabolite of disulfiram, in urine (229) 470

**DOPAC**
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

**Dopamine**
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

Determination of dopamine, homovanillic acid and 3,4-dihydroxyphenylacetic acid in rat brain striatum by HPLC with electrochemical detection (222) 353

Determination of catecholamines by radio-enzymatic assay using ion-pair LC (223) 41

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143

Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1

Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35

**Dopamine analogues**
Reversed-phase LC with amperometric detection of lipophilic dopamine analogues and determination of brain and serum concentrations after sample clean-up on small Sephadex G-10 columns (230) 271

**Dopamine-β-hydroxylase**
Studies on dopamine-converting enzymes in human plasma (223) 188

Assay for dopamine β-hydroxylase in rat serum and adrenal medulla by HPLC with fluorescence detection (227) 415

**Dothiepin**
Simultaneous measurement of dothiepin and its major metabolites in plasma and whole blood by GC-MF (222) 399

**Doxorubicin**
Extraction and quantification of daunomycin and doxorubicin in tissues (221) 139

Isotachophoretic determination of adriamycin and adriamycinol in human plasma (227) 262

**Dropping mercury electrode**
Differential pulse amperometric detection of drugs in plasma using a dropping mercury electrode as a HPLC detector (222) 179

**Drug-binding proteins**
A new method for detection of drug-binding proteins using a parallel-flow dialysis technique (223) 267

**Drug monitoring**
Fully automated HPLC. A new chromatograph for pharmacokinetic drug monitoring by direct injection of body fluids (222) 13

**Dyphylline**
Simultaneous determination of dyphylline and theophylline in human plasma by HPLC (221) 170

# E

**Elastin**
Rapid method for the purification of the elastin cross-links, desmosine and isodesmosine (229) 200

**Enantiomer separation**
Catalytic detection principle for HPLC: determination of enantiomeric iodinated thyronines in blood serum (224) 239

**Endralazine**
Sensitive HPLC assay for endralazine and two of its metabolites in human plasma (225) 151

**Enzymes**
see also proteins

Direct assay for creatine kinase by reversed-phase HPLC (221) 39

GC determination of monoamine oxidase activity using mixed substrates (221) 155

New and highly sensitive assay for L-5-hydroxytryptophan decarboxylase activity by HPLC-voltammetry (221) 265

Isoelectric focusing studies of human pancreatic secretion (221) 279

Action pattern of human pancreatic α-amylase on maltoheptaose, a substrate for determining α-amylase in serum (223) 69

Studies on dopamine-converting enzymes in human plasma (223) 188

Determination of 5-aminolaevulinic acid dehydrase activity in erythrocytes by HPLC (223) 421

Use of reversed-phase HPLC for simultaneous determination of glutamine synthetase and glutamic acid decarboxylase in crude extracts (225) 37

GC-MS determination of glutamic acid decarboxylase activity in subregions of rat brain (225) 283

Rapid assay of spermidine synthase activity by HPLC (226) 208

Assay of catechol O-methyltransferase activity by HPLC with electrochemical detection (226) 461

Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407

Assay for dopamine β-hydroxylase in rat serum and adrenal medulla by HPLC with fluorescence detection (227) 415

Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1

Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77

The determination of uroporphyrinogen decarboxylase in tissues by HPLC coupled to fluorescence detection (229) 311

Rapid and sensitive sialidase assay by HPLC and its application to detection of sialidase in human urine (230) 415

**Epidoxorubicin**
Determination of 4'-epidoxorubicin and its 13-dihydro derivative in human plasma by HPLC with fluorescence detection (230) 207

**Epinephrine**
Novel post-column derivatization method for the fluorimetric determination of norepinephrine and epinephrine (221) 249

Determination of catecholamines by radio-enzymatic assay using ion-pair LC (223) 41

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1

**Epoxyphosphonic acid**
Simple selected ion monitoring method for determination of fosfomycin in blood and urine (224) 257

**Ergot alkaloids**
Determination of ergot alkaloids in plasma by HPLC and fluorescence detection (226) 107

**Ergotamine**
Determination of ergot alkaloids in plasma by HPLC and fluorescence detection (226) 107

**Estradiol**
Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33

The use of oestradiol-17β antiserum covalently coupled to Sepharose to extract oestradiol-17β from biological fluids (223) 193

HPLC analysis of estradiol-17β and metabolites in biological media (224) 205

Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

**Estriol**
LC assay of urinary estriol and electrochemical detection with a battery powered detector (224) 99

Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

**Estrogens**
see oestrogens

**Estrone**
Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33

Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

**Ethacrynic acid**
Determination of ethacrynic and tienilic acid in plasma by GLC-MS (227) 193

**Ethambutol**
Determination of ethambutol in plasma using selected ion monitoring (221) 293

GC-MS determination of ethambutol in human plasma (224) 415

**Ethionamide**
HPLC determination of ethionamide and prothionamide in body fluids (225) 245

**Ethosuximide**
Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

**Ethoxybenzamide**
Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479

**Ethyl loflazepate**
Analysis of the metabolites of ethyl loflazepate by GC with electron-capture detection (226) 413

**Etiocholanolone**
Steroid profiles of healthy individuals (223) 21

**Etoposide**
Analysis of the anticancer drugs etoposide (VP 16-213) and teniposide (VM 26) by HPLC with fluorescence detection (224) 168

**Etozolin**
Determination of etozolin and ozolinone in human plasma and tissues by reversed-phase HPLC (225) 504

**Exogenous volatiles**
Volatiles of exogenous origin from the human oral cavity (226) 315

# F

**Fansidar**
Determination of pyrimethamine in human plasma after administration of Fansidar or Fansidar-mefloquine by means of HPLC with fluorescence detection (230) 107

**Fatty acids**
A microliter method for the GC determination of long-chain non-esterified fatty acids in human serum or plasma (221) 205

Simultaneous measurement of sterol and fatty acid composition in small samples (221) 367

Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

Glass capillary GC of the serum fatty acid fraction via automatic injection of lipid extracts (228) 75

Lipids in human neutrophils determined by a microanalytical method (229) 455

**Fenbufen**
HPLC assay for fenbufen and two serum metabolites (228) 372

**Fenofibrate**
Rapid determination of procetofenic acid in plasma by HPLC (227) 219

**Fenoterol**
A time-saving method for the determination of the $\beta_2$ sympathomimetics terbutaline, salbutamol and fenoterol. Preliminary results (229) 217

**Fentanyl**
The assay of fentanyl and its metabolites in plasma of patients using GC with alkali flame ionisation detection and GC-MS (223) 85

**Ferritin**
The effect of ampholytes on ferritin isoelectric focussing patterns (230) 327

**Flavin-adenine dinucleotide**
Determination of the $B_2$ vitamer flavin-adenine dinucleotide in whole blood by HPLC (228) 311

Hydrolysis of riboflavin nucleotides in plasma monitored by HPLC (229) 445

**Flavonoids**
Determination of sodium flavodate in body fluids by HPLC. Application to clinical pharmacokinetic studies (230) 168

**Flecainide**
Fluorometric LC assay of the antiarrhythmic agent flecainide in blood plasma (229) 498

**Flumecinol**
GC method for the determination of flumecinol in biological fluids (221) 403

**Flunitrazepam**
Assay of flunitrazepam, temazepam and desalkylflurazepam in plasma by capillary GC with electron-capture detection (227) 199

**Fluorescence detection**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

**Fluoropyrimidine**
Determination of a novel fluoropyrimidine, 5'-deoxy-5-fluorouridine, in plasma by HPLC (225) 516

**Fluoropyrimidines**
Comparison of the reversed-phase HPLC separations of fluoropyrimidines, pyrimidines, and purines (228) 165

**Fluorouracil**
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382

GC-MS determination of plasma 5-fluorouracil after administration of 1-hexylcarbamoyl-5-fluorouracil to dogs and humans (224) 423

Quantitative determination of 1-hexylcarbamoyl-5-fluorouracil and its metabolites in man (225) 91

Reversed-phase ion-pair HPLC assay of 5-fluorouracil, 5'-deoxy-5-fluorouridine, their nucleosides, mono-, di-, and triphosphate nucleotides with a mixture of quaternary ammonium ions (228) 245

**Flurazepam**
Rapid quantitation of flurazepam and its major metabolite, N-desalkylflurazepam, in human plasma by GLC with electron-capture detection (222) 491

Determination of the major urinary metabolite of flurazepam in man by HPLC (223) 123

**Flurbiprofen**
Determination of flurbiprofen in human plasma using GC-MS with selected ion monitoring (223) 202

**FM 24**
HPLC assay for determination of a new β-blocking agent FM 24 (230) 199

**Food additives**
Determination of the antioxidant 3-*tert.* -butyl-4-hydroxyanisole in rat plasma using high-resolution GC-MS (225) 83

**Formoterol**
Quantitative determination of the β-adrenoceptor stimulant formoterol in urine by GC-MS (229) 337

**Fosfomycin**
Simple selected ion monitoring method for determination of fosfomycin in blood and urine (224) 257

**Furosemide**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

Improved method for the analysis of furosemide in plasma by HPLC (227) 463

Specimen handling and HPLC determination of furosemide (229) 129

TLC determination of furosemide and 4-chloro-5-sulfamoyl anthranilic acid in plasma and urine (229) 425

# G

**Gangliosides**
Simple micro-method for the isolation of gangliosides by reversed-phase chromatography (224) 249

**Gas analyzer**
A new chromatographic instrument for measuring trace concentrations of breath-hydrogen (229) 259

**Gentamicin**
Determination of gentamicin in serum using LCC (221) 182

Improved microdetermination of gentamicin and sisomicin in serum by HPLC with ultraviolet detection (222) 316

HPLC method for determination of gentamicin in biological fluids (223) 131

Micro determination of gentamicin in serum by HPLC (227) 244

**Gentisic acid**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

A method for HPLC screening of UV-positive components in urine eluate from Sephadex G-10 and modifications for determination of urinary salicylic, salicyluric and gentisic acids (230) 253

**Glafenine**
Simultaneous quantitation of glafenine and its major metabolites, glafenic and hydroxyglafenic acid, by HPLC (224) 348

**Glibenclamide**
HPLC and TLC determination of human serum glibenclamide at therapeutic levels (227) 93

**Gliclazide**
GC determination of the hypoglycaemic agent gliclazide in plasma (223) 357

**Globin chains**
An update on electrophoretic and chromatographic methods in the diagnosis of hemoglobinopathies (Review) (227) 267

Rapid separation of human globin chains by high flow-rate chromatography under low pressure using CM-Trisacryl M (228) 298

**Glucocorticoids**
Determination of methylprednisolone in central nervous tissue and plasma using normal-phase HPLC (230) 29

**Glutamic acid**
GC determination of glutamine in the presence of glutamic acid in biological fluids (224) 177

**Glutamic acid decarboxylase**
Use of reversed-phase HPLC for simultaneous determination of glutamine synthetase and glutamic acid decarboxylase in crude extracts (225) 37

GC-MS determination of glutamic acid decarboxylase activity in subregions of rat brain (225) 283

**Glutamine**
GC determination of glutamine in the presence of glutamic acid in biological fluids (224) 177

Chromatographic analysis of glutamine in plasma (229) 188

**Glutamine synthetase**
Use of reversed-phase HPLC for simultaneous determination of glutamine synthetase and glutamic acid decarboxylase in crude extracts (225) 37

**Glycerophosphoryl ethanolamine**
Separation of taurine and glycerophosphoryl ethanolamine on amino acid analyzers (223) 179

**Glyceryl trinitrate**
Determination of glyceryl trinitrate in human plasma by GC-negative ion chemical ionization-selected ion monitoring (229) 327

**Glycoconjugates**
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Glycolic acid**
Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

**Glycoproteins**
Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

Microprocedure to determine the polymorphic forms of acid $\alpha_1$-glycoprotein in plasma. Application to depressive patients treated with amitriptyline (229) 319

**Glycosaminoglycans**
The effect of pH and ionic strength on the electrophoretic separation of acidic glycosaminoglycans (221) 49

The application of HPLC in enzymatic assays of chondroitin sulfate isomers in normal human urine (222) 23

Electrophoresis with two buffers in one dimension in the analysis of glycosaminoglycans on cellulose acetate strips (224) 407

**Gonadotropic hormones**
HPLC estimation of cyproterone acetate in human plasma (226) 492

**Gradient elution**
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215

**Guaiphenesin**
Determination of guaiphenesin and its metabolite, β-(2-methoxyphenoxy)lactic acid, in plasma by HPLC (224) 144

**Guanidine**
GC-MS analysis of amine metabolites in urine (228) 51

**Guanidino compounds**
Isolation and identification of some guanidino compounds in the urine of patients with hyperargininaemia by LC, TLC and GC-MS (224) 185

Post-column derivatization of guanidino compounds in HPLC using ninhydrin (226) 43

# H

**Haemoglobins**
Carrier-ampholyte displacement chromatography (chromatofocusing) on ion-exchange papers. Application to the separation of haemoglobin variants (222) 116

An update on electrophoretic and chromatographic methods in the diagnosis of hemoglobinopathies (Review) (227) 267

Chromatofocusing of human hemoglobins: application to the quantitation of hemoglobin $A_2$ (228) 177

Rapid separation of human globin chains by high flow-rate chromatography under low pressure using CM-Trisacryl M (228) 298

Separation of peptides by cellulose-phosphate chromatography for identification of a hemoglobin variant (229) 193

Chromatofocusing of human hemoglobins (230) 137

**Haloperidol**
HPLC determination of haloperidol in plasma (223) 449

Measurement of haloperidol in human plasma using reversed-phase HPLC (227) 233

**Hemoglobins**
see haemoglobins

**Heparan sulphate**
Electrophoresis with two buffers in one dimension in the analysis of glycosaminoglycans on cellulose acetate strips (224) 407

**Heparin**
Gel chromatography of heparin (226) 61

Gel chromatography of heparin (230) 216

**Hexadecanoic acid octadecyl ester**
Identification of some abnormal metabolites in psoriatic nail using GC-MS (221) 199

**Hexobarbital**
Rapid analysis of hexobarbital and its main metabolites in mice by HPLC (228) 350

**Hexosamines**
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Hexylcarbamoylfluorouracil**
GC-MS determination of plasma 5-fluorouracil after administration of 1-hexylcarbamoyl-5-fluorouracil to dogs and humans (224) 423

Quantitative determination of 1-hexylcarbamoyl-5-fluorouracil and its metabolites in man (225) 91

**HI-6**
Determination of some pyridinium aldoxime compounds by means of ion-pair reversed-phase HPLC: application in biological material (225) 107

**Hippuric acid**
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215

Determination of urinary hippuric acid and *o*-cresol, as indices of toluene exposure, by LC on dynamically modified silica (229) 141

**Histamine**
Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human plasma and urine by GC-MS (221) 1

Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human urine and rat brain by HPLC with fluorescence detection (224) 105

Histamine in tissue: determination by HPLC after condensation with *o*-phthaldialdehyde (226) 53

**Homocarnosine**
Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315

**Homopantothenic acid**
GC-MF determination of homopantothenic acid in plasma (226) 333

**Homovanillic acid**
Simultaneous determination of dopamine, DOPAC and homovanillic acid. Direct injection of supernatants from brain tissue homogenates in a LC-electrochemical detection system (221) 237

Determination of dopamine, homovanillic acid and 3,4-dihydroxyphenylacetic acid in rat brain striatum by HPLC with electrochemical detection (222) 353

Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

HPLC determination of homovanillic acid in urine using Sephadex G-10 for isolation (225) 440

Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

Simple method for the determination of homovanillic acid and vanillylmandelic acid in urine by HPLC (227) 162

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

Simultaneous determination of 4-hydroxy-3-methoxy-phenylacetic (homovanillic) acid and other monoamine metabolites in human lumbar cerebrospinal fluid. An improved HPLC study with electrochemical detection (227) 379

**Hyaluronic acid**
Electrophoresis with two buffers in one dimension in the analysis of glycosaminoglycans on cellulose acetate strips (224) 407

**Hydrogen**
A new chromatographic instrument for measuring trace concentrations of breath-hydrogen (229) 259

**Hydroxybutyrate**
Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455

**Hydroxybutyric acid**
GC-MS analysis of organic acids in renal tissue biopsy. Identification of 4-hydroxybutyric acid and 4-hydroxy-2-butenoic acid (230) 1

**Hydroxycorticosterone**
Estimation of free urinary aldosterone and 18-hydroxycorticosterone by a combination of automatic HPLC and radioimmunoassay (227) 492

**Hydroxydantrolene**
Simultaneous determination of dantrolene and its metabolite, 5-hydroxydantrolene, in human plasma by HPLC (228) 404

**Hydroxydesipramine**
Determination of 2-hydroxydesipramine by HPLC (226) 147

**Hydroxydicarboxylic acids**
Determination of the absolute configuration of some biologically important urinary 2-hydroxydicarboxylic acids by capillary GLC (222) 276

**(Hydroxyethoxymethyl)guanine**
Simple HPLC method for the analysis of 9-(2-hydroxyethoxymethyl)guanine (acyclovir) in human plasma and urine (224) 51

**Hydroxyethylflurazepam**
Determination of the major urinary metabolite of flurazepam in man by HPLC (223) 123

**Hydroxyglafenic acid**
Simultaneous quantitation of glafenine and its major metabolites, glafenic and hydroxyglafenic acid, by HPLC (224) 348

**Hydroxyindoleacetic acid**
Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

Determination of indoles in human and rat pineal (228) 155

**Hydroxymandelic acid**
Excretion of *m*-hydroxymandelic acid in human urine (222) 271

**Hydroxymethohexital**
Simultaneous determination of blood concentrations of methohexital and its hydroxy metabolite by GC and identification of 4'-hydroxymethohexital by combined GLC-MS (226) 403

**Hydroxymethotrexate**
Rapid and simple method for the measurement of methotrexate and 7-hydroxymethotrexate in serum by HPLC (223) 225

**Hydroxymethoxyindole**
Identification of two Thormählen-positive compounds from melanotic urine by GC-MS (222) 329

**Hydroxymethyllevulinic acid**
Identification of 2-hydroxy-2-methyllevulinic acid in urine and serum of diabetic patients with ketoacidosis (228) 59

**Hydroxymetoprolol**
Determination of metoprolol and α-hydroxymetoprolol in plasma by HPLC (228) 215

**Hydroxynalidixic acid**
HPLC analysis of nalidixic acid and hydroxynalidixic acid in plasma with a dynamic anion-exchange system (221) 129

**Hydroxynitroquinoline**
HPLC analysis of nitroxoline in plasma and urine (222) 241

**Hydroxyphenobarbital**
Rapid quantitative determination of underivatized carbamazepine, phenytoin, phenobarbital and *p*-hydroxyphenobarbital in biological fluids by packed column GC (221) 75

**Hydroxyphenylacetic acid**
Longitudinal urinary excretion of some "trace" acids in a human male (222) 161

Plasma concentrations of *p*- and *m*-hydroxyphenylacetic acid and phenylacetic acid in humans. GC-high-resolution MS analysis (230) 219

**Hydroxyprednisolone**
Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283

**Hydroxyprednisone**
Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221

**Hydroxypregnenolone**
Quantitation of pregnenolone and 17-hydroxypregnenolone in human serum by automatic HPLC and subsequent radioimmunoassay (224) 229

**Hydroxyprogesterone**
Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17α-hydroxyprogesterone from small plasma or tissue samples (228) 95

**Hydroxyproline**
Polymorphism of urinary 4-hydroxyproline-containing polypeptides (225) 55

Simplified procedure for the analysis of 3- and 4-hydroxyproline (227) 341

Improved method for the evaluation of 3-hydroxyproline and 4-hydroxyproline in the urine (230) 420

**Hydroxypropranolol**
Simple and rapid determination of propranolol and its active metabolite, 4-hydroxypropranolol, in human plasma by LC with fluorescence detection (228) 362

**Hydroxypropyldiaminobutane**
GC-MS analysis of amine metabolites in urine (228) 51

**Hydroxysalicylanilide**
Determination of osalmid in plasma by HPLC (223) 243

**Hydroxysteroids**
Enzymatic detection of urinary acidic 3α-hydroxysteroids on thin-layer chromatograms (224) 111

Enzymatic determination of urinary 17β-hydroxysteroids on thin-layer chromatograms (228) 317

**Hydroxythiabendazole**
Determination of thiabendazole and 5-hydroxythiabendazole in human serum by fluorescence-detected HPLC (230) 79

**Hydroxytryptamine**
Determination of serotonin, its precursors, metabolites and [$^3$H]serotonin in lung by HPLC with fluorescence detection (225) 27

Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347

Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35

**Hydroxytryptophan**
Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

**Hydroxytryptophan decarboxylase**
New and highly sensitive assay for L-5-hydroxytryptophan decarboxylase activity by HPLC-voltammetry (221) 265

**Hydroxytryptophol**
Determination of indoles in human and rat pineal (228) 155

**Hypoxanthine**
Mass fragmentographic determination of xanthine and hypoxanthine in biological fluids (221) 9

Analysis of adenosine, inosine and hypoxanthine in suspensions of cardiac myocytes by HPLC (226) 202

Simultaneous determination of allopurinol and oxipurinol in biological fluids by mass fragmentography (229) 211

**Hypoxanthine arabinoside**
Reversed-phase HPLC method to determine vidarabine and hypoxanthine arabinoside in biological fluids (229) 487

# I

**Ibuprofen**
Determination of ibuprofen in serum by HPLC and application to ibuprofen disposition (226) 183

**Imidazobenzodiazepines**
Determination of the anxiolytic agent 8-chloro-6-(2-chlorophenyl)-4H-imidazo-[1,5-*a*][1,4]-benzodiazepine-3-carboxamide in whole blood, plasma or urine by HPLC (226) 135

**Indalpine**
HPLC determination of indalpine, a new non-tricyclic antidepressant, in human plasma. Identification and simultaneous measurement of its major plasma metabolite (230) 87

**Indapamide**
HPLC assay for monitoring indapamide and its major metabolite in urine (228) 377

HPLC analysis of indapamide (RHC 2555) in urine, plasma and blood (230) 181

**Indoleamines**
Determination of a wide range of urinary amine metabolites using a simple HPLC technique (225) 179

Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase HPLC with electrochemical detection (225) 347

Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection.

Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

**Indoles**
Analysis of some tryptophan and phenylalanine metabolites in urine by a straight-phase HPLC technique (222) 191

Identification of two Thormählen-positive compounds from melanotic urine by GC-MS (222) 329

Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

Determination of serotonin, its precursors, metabolites and [$^3$H]serotonin in lung by HPLC with fluorescence detection (225) 27

Determination of indoles in human and rat pineal (228) 155

**Indoloquinolinones**
Determination of the anti-tumor agent, 10-chloro-5-(2-dimethylaminoethyl)-7H-indolo[2,3-C]quinolin-6(5H)-one in blood or plasma by HPLC (223) 111

**Indomethacin**
HPLC-fluorescence analysis for indomethacin and metabolites in biological fluids (229) 179

**Inosine**
Analysis of adenosine, inosine and hypoxanthine in suspensions of cardiac myocytes by HPLC (226) 202

**Iodinated thyronines**
Catalytic detection principle for HPLC: determination of enantiomeric iodinated thyronines in blood serum (224) 239

**Iodohippurate**
HPLC method for the simultaneous determination of iothalamate and *o*-iodohippurate (224) 139

**Iodothyronines**
Simultaneous determination of D- and L-thyroxine in human serum by LC with electrochemical detection (226) 383

**Iothalamate**
HPLC method for the simultaneous determination of iothalamate and *o*-iodohippurate (224) 139

**Iron**
Analysis of serum iron by gel permeation HPLC (221) 271

**Isoamylases**
Isoelectric focusing studies of human pancreatic secretion (221) 279

**1-Isobutyl-3,4-diphenylpyrazole-5-acetic acid**
Determination of a new anti-inflammatory agent, 1-isobutyl-3,4-diphenylpyrazole-5-acetic acid, by HPLC (227) 213

**Isodesmosine**
Rapid method for the purification of the elastin cross-links, desmosine and isodesmosine (229) 200

**Isoenzymes**
Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1

**Isoniazid**
Determination of isoniazid and its hydrazino metabolites, acetylisoniazid, acetylhydrazine, and diacetylhydrazine in human plasma by GC-MS (224) 431

Hydrophilic ion-pair reversed-phase HPLC for the simultaneous assay of isoniazid and acetylisoniazid in serum: a microscale procedure (226) 250

Studies of isoniazid metabolism in isolated rat hepatocytes by mass fragmentography (230) 345

**Isoputreanine**
MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142

**Isosorbide dinitrate**
GC determination of isosorbide dinitrate in human plasma and urine (229) 347

**Isoxepac**
Determination of isoxepac in plasma by HPLC (221) 431

Determination of 6,11-dihydro-11-dibenz[*b,e*]-oxepin-2-acetic acid (isoxepac) in plasma by HPLC (225) 482

**Isozymes**
Isoelectric focusing studies of human pancreatic secretion (221) 279

# K

**Keto acids**
HPLC determination of α-keto acids (224) 197

**Ketoconazole**
Determination of the antifungal agent, ketoconazole, in human plasma by HPLC (221) 337

Measurement of ketoconazole, a new antifungal agent, by HPLC (227) 510

**Ketoglutarate**
Quantitative GC-chemical ionization MS of 2-ketoglutarate from urine as its O-trimethylsilyl-quinoxalinol derivative (226) 325

**Ketomethylthiobutyric acid**
GC and GC-MS studies on α-keto-γ-methylthiobutyric acid in urine following ingestion of optical isomers of methionine (221) 145

**Ketones**
Excretion of urinary volatile metabolites in response to alloxan induced diabetes of short duration in rats (228) 33

**Keto prostaglandin**
Quantitative determination of 6-keto prostaglandin $F_{1\alpha}$ in biological fluids by capillary GC-chemical ionization MS (221) 361

**Ketosteroids**
Simple computer program for a low-cost desktop calculator applied to the evaluation of GLC analyses of 17-ketosteroids and pregnanes (223) 253

# L

**Labetalol**
Sensitive HPLC method for the determination of labetalol in human plasma using fluorimetric detection (226) 259

**Lactam antibiotics**
Determination of cephalosporins in biological material by reversed-phase LCC (223) 393

HPLC analysis of a new β-lactam antibiotic, 6059-S (moxalactam) (225) 169

HPLC determination of cephalexin in human plasma, urine and saliva (225) 532

Determination of cefotaxime and desacetyl-cefotaxime in plasma and urine by HPLC (226) 431

Determination of cefsulodin, cefotiam, cephalexin, cefotaxime, desacetyl-cefotaxime, cefuroxime and cefroxadin in plasma and urine by HPLC (228) 257

Determination of moxalactam by HPLC (228) 418

Determination of cefaclor by HPLC (228) 429

A very precise HPLC procedure for the determination of cefmenoxime, a new cephalosporin antibiotic, in plasma (229) 149

**Lactate**
Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455

**Lactate dehydrogenase**
Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1

**Lactic acid**
Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

**Laxatives**
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389

**Lecithin**
HPLC determination of the lecithin/sphingomyelin ratio in amniotic fluid (223) 277

Improved TLC assay for monitoring lecithin/sphingomyelin ratios in amniotic fluid (226) 484

**Levamisole**
Ion-pair HPLC assay of levamisole in biological fluids (223) 445

Determination of levamisole in plasma and animal tissues by GC with thermionic specific detection (224) 25

**Levomepromazine**
Simultaneous determination of chlorpromazine and levomepromazine in human plasma and urine by HPLC using electrochemical detection (227) 103

**Librium**
Simultaneous determination of chlordiazepoxide and its metabolites in human plasma and urine by means of reversed-phase HPLC (224) 519

**Lignocaine**
Simple method for the measurement of tocainide and lignocaine in blood plasma or serum using GLC with flame ionisation detection (225) 469

**Lipases**
Isoelectric focusing studies of human pancreatic secretion (221) 279

**Lipids**
Studies of hyperlipidemia in drug-induced diabetic rats by HPLC (221) 19

Simultaneous measurement of sterol and fatty acid composition in small samples (221) 367

Lipid class and molecular species interrelationships among plasma lipoproteins of normolipemic subjects (224) 1

Simple micro-method for the isolation of gangliosides by reversed-phase chromatography (224) 249

Rapid determination of lipids in healthy human lymphocytes (227) 423

Glass capillary GC of the serum fatty acid fraction via automatic injection of lipid extracts (228) 75

Lipids in human neutrophils determined by a microanalytical method (229) 455

Lipid class and molecular species interrelationships among plasma lipoproteins of Type III and Type IV hyperlipemic subjects (230) 231

**Lipoproteins**
High-performance aqueous gel permeation chromatography of human serum lipoproteins (221) 257

High-performance aqueous gel permeation chromatography of serum lipoproteins: selective detection of cholesterol by enzymatic reaction (223) 285

Lipid class and molecular species interrelationships among plasma lipoproteins of normolipemic subjects (224) 1

Separation of human C-apolipoproteins by HPLC (225) 185

Lipid class and molecular species interrelationships among plasma lipoproteins of Type III and Type IV hyperlipemic subjects (230) 231

**Lorazepam**
HPLC determination of lorazepam in monkey plasma (226) 155

**Lorcainide**
HPLC analysis of lorcainide and its active metabolite, norlorcainide, in human plasma (223) 454

# M

**Maltoheptaose**
Action pattern of human pancreatic $\alpha$-amylase on maltoheptaose, a substrate for determining $\alpha$-amylase in serum (223) 69

**Mandelic acid**
Simultaneous determination of cyclandelate and its metabolite in human plasma by capillary column GLC (223) 365

**Maprotiline**
GC analysis of therapeutic concentrations of maprotiline in serum, using flame-ionization detection (221) 319

GC determination of maprotiline and its N-desmethyl metabolite in human blood using nitrogen detection (221) 393

GLC procedure with alkali flame ionization detection for the determination of maprotiline in plasma (224) 128

**Mecillinam**
Determination of amdinocillin in plasma and urine by high-performance liquid chromatography (227) 137

**Mefloquine**
Determination of pyrimethamine in human plasma after administration of Fansidar or Fansidar-mefloquine by means of HPLC with fluorescence detection (230) 107

**Melanotropin-inhibiting factor**
Determination of the melanotropin-inhibiting factor analogue pareptide in urine by HPLC (222) 512

**Melatonin**
Determination of indoles in human and rat pineal (228) 155

**Melphalan**
Quantitative analysis of melphalan and its major hydrolysate in patients and animals by reversed-phase HPLC (222) 453

Method for the measurement of melphalan in biological samples by HPLC (224) 338

**Mercaptoacetate**
Reversed-phase ion-pair HPLC of mercaptoacetate and N-acetylcysteine after derivatization with N-(1-pyrene)maleimide and N-(7-dimethylamino-4-methyl-3-coumarinyl)maleimide (229) 409

**Mercaptopurine**
Extractive alkylation of 6-mercaptopurine and determination in plasma by GC-MS (225) 73

HPLC determination of plasma 6-mercaptopurine in clinically relevant concentrations (227) 526

Quantitation of 6-mercaptopurine in biologic fluids using HPLC: a selective and novel procedure (230) 373

**Metabolic profiles**
Identification of some abnormal metabolites in psoriatic nail using GC-MS (221) 199

High-resolution GC-MS of the methyl esters of organic acids from uremic hemofiltrates (223) 1

Myo-inositol levels in the cerebrospinal fluid of infants (223) 173

Analysis of profiles of conjugated steroids in urine by ion-exchange separation and GC-MS (224) 355

GC-MS profile of organic acids in urine and serum of diabetic ketotic patients (225) 1

A GC-MS study of profiles of volatile metabolites in hepatic encephalopathy (226) 291

Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

Excretion of urinary volatile metabolites in response to alloxan induced diabetes of short duration in rats (228) 33

GC-MS analysis of amine metabolites in urine (228) 51

Comparison of isolation methods of urinary organic acids HPLC (228) 103

GC-MS analysis of organic acids in renal tissue biopsy. Identification of 4-hydroxybutyric acid and 4-hydroxy-2-butenoic acid (230) 1

Analysis of organic acids in the hearts of patients with idiopathic cardiomyopathy by GC-MS (230) 7

**Metabolic profils**
Improved procedure for the anion-exchange isolation of urinary organic acids (222) 337

**Metallothionein**
HPLC determination of tissue metallothionein in monkeys chronically exposed to cadmium (228) 285

**Metals**
Analysis of serum iron by gel permeation HPLC (221) 271

Improved method for selenium determination in biological samples by GC (225) 9

**Metapramine**
HPLC method for the determination of plasma and urine metapramine after dansylation (228) 205

**Methohexital**
Simultaneous determination of blood concentrations of methohexital and its hydroxy metabolite by GC and identification of 4'-hydroxymethohexital by combined GLC-MS (226) 403

**Methotrexate**
Rapid and simple method for the measurement of methotrexate and 7-hydroxymethotrexate in serum by HPLC (223) 225

Sensitive and rapid HPLC method for the simultaneous determination of methotrexate and its metabolites in plasma, saliva and urine (226) 125

**Methoxyhydroxyphenethyleneglycol**
Determination of the primary metabolite of central nervous system norepinephrine, 3-methoxy-4-hydroxyphenethyleneglycol, in mouse brain and brain perfusate by HPLC with electrochemical detection (223) 295

**Methoxyhydroxyphenylethyleneglycol**
Amperometric determination of 3-methoxy-4-hydroxyphenylethyleneglycol in human cerebrospinal fluid (222) 112

Quantitative determination of 3-methoxy-4-hydroxyphenylethyleneglycol and its sulfate conjugate in human lumbar cerebrospinal fluid using LC with aperometric detection (223) 305

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

**Methoxyhydroxyphenylglycol**
Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

Analysis of urinary 3-methoxy-4-hydroxyphenylglycol by HPLC and electrochemical detection (227) 391

**(Methoxyphenoxy)lactic acid**
Determination of guaiphenesin and its metabolite, β-(2-methoxyphenoxy)lactic acid, in plasma by HPLC (224) 144

**Methoxytyramine**
Simultaneous extraction and separation of trace amines of biological interest (225) 301

GC-MS analysis of amine metabolites in urine (228) 51

**Methyclothiazide**
Determination of methyclothiazide in human plasma by HPLC (226) 510

**Methylaminophenazone**
Rapid TLC of various weak analgesics in saliva (225) 427

**Methylbenz[*c*]acridine**
TLC and HPLC separation of metabolites of the weak carcinogen, 7-methylbenz[*c*]acridine (227) 149

**Methyldopa**
Determination of α-methyldopa, α-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

Assay and stability of α-methyldopa in man using HPLC with electrochemical detection (229) 433

Determination of α-methyldopa and methyldopate in human breast milk and plasma by ion-exchange chromatography using electrochemical detection (230) 162

**Methylergometrine**
Determination of ergot alkaloids in plasma by HPLC and fluorescence detection (226) 107

**Methylguanidine**
GC-MS analysis of amine metabolites in urine (228) 51

**Methylhistamine**
Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human plasma and urine by GC-MS (221) 1

Simultaneous determination of histamine and $N^{\tau}$-methylhistamine in human urine and rat brain by HPLC with fluorescence detection (224) 105

**Methylhistidine**
Glass capillary GC determination of $N^{\tau}$-methylhistidine in urine (221) 149

Simple and rapid HPLC method for the quantification of 3-methylhistidine (223) 417

Glass capillary quantitative determination of $N^{\pi}$-methylhistidine in urine and muscles (224) 94

Rapid analysis of 3-methylhistidine in urine, plasma, muscle and amniotic fluid with a single HPLC system but with different ion-pairing reagents (228) 273

**Methylnoradrenaline**
Determination of α-methyldopa, α-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

**Methylphenobarbital**
Simultaneous assay of methylphenobarbital and phenobarbital in plasma using GC-MS with selected ion monitoring (223) 426

**Methylprednisolone**
Determination of methylprednisolone in central nervous tissue and plasma using normal-phase HPLC (230) 29

**Methylpyridinium aldoxime**
Determination of N-methylpyridinium 2-aldoxime methylsulfate (Contrathion) in rat plasma and urine by copper(II)-silica ligand-exchange HPLC (229) 475

**Methylthioadenosine**
HPLC analysis of 5'-methylthioadenosine in rat tissues (226) 243

**Metioprim**
Determination of 2,4-diamino-5-benzylpyrimidines in combination with sulphadiazine in biological fluids using HPLC (224) 301

**Metipranolol**
Determination of deacetylmetipranolol in body fluids by GC-chemical-ionization MS (227) 187

**Metolazone**
Quantitation of metolazone in plasma and urine by HPLC with fluorescence detection (222) 311

Determination of the diuretic agent metolazone in plasma by HPLC (226) 526

**Metoprolol**
Determination of metoprolol in plasma and urine by GLC with electron-capture detection (225) 213

Determination of (*R*)- and (*S*)-alprenolol and (*R*)- and (*S*)-metoprolol as their diastereomeric derivatives in human plasma by reversed-phase LC (227) 113

Determination of metoprolol and α-hydroxy-metoprolol in plasma by HPLC (228) 215

HPLC method for the quantitative analysis of the aryloxypropanolamines propranolol, metoprolol and atenolol in plasma and tissue (228) 223

**Metronidazole**
Determination of metronidazole, misonidazole and its metabolite in serum and urine on RP-18 HPTLC plates (224) 465

**Mexiletine**
Determination of mexiletine in biological fluids by HPLC (230) 97

**Mezlocillin**
Improved procedure for the determination of the ureidopenicillins azlocillin and mezlocillin in plasma by HPLC (228) 409

**Miconazole**
HPLC analysis of miconazole in plasma (227) 223

Determination of miconazole in human saliva using HPLC (227) 229

**Midazolam**
Determination of midazolam by HPLC (225) 266

Midazolam determination by GC, LC and GC-MS (228) 195

**Minaxolone**
Determination of minaxolone (CCI 12923) by GC with nitrogen-sensitive detection (224) 496

**Misonidazole**
Determination of metronidazole, misonidazole and its metabolite in serum and urine on RP-18 HPTLC plates (224) 465

Simultaneous determination of misonidazole and desmethylmisonidazole in plasma by HPLC (225) 407

**Mobile phase composition**
The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

**Monoacetyldiamines**
Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363

**Monoamine oxidase**
GC determination of monoamine oxidase activity using mixed substrates (221) 155

**Monosaccharides**
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Moroxydine**
Determination of moroxydine in biological fluids by electron-capture GC (229) 227

**Morphine**
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

Determination of morphine, morphine-3-glucuronide and (tentatively) morphine-6-glucuronide in plasma and urine using ion-pair HPLC (230) 427

**Moxalactam**
HPLC analysis of a new β-lactam antibiotic, 6059-S (moxalactam) (225) 169

Determination of moxalactam by HPLC (228) 418

**Myo-inositol**
Myo-inositol levels in the cerebrospinal fluid of infants (223) 173

# N

**Nadolol**
Fluorodensitometric determination of nadolol in plasma and urine (230) 461

**Nalidixic acid**
HPLC analysis of nalidixic acid and hydroxy-nalidixic acid in plasma with a dynamic anion-exchange system (221) 129

TLC method for the quantitative analysis of nalidixic acid in human plasma (223) 165

**Naproxen**
Mass fragmentographic quantification of naproxen in human plasma (222) 482

Quantitative determination of naproxen in plasma by a simple HPLC method (224) 43

**Nefopam**
Quantitative determination of nefopam in human plasma, saliva and cerebrospinal fluid by GLC using a nitrogen-selective detector (226) 79

**Neopterine**
Determination of neopterine in human urine by reversed-phase HPLC (227) 61

**Neuropeptides**
Characterization and determination of neuropeptides by HPLC and radioimmunoassay (222) 371

Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381

Rapid high-yield purification of neuropeptides from canine intestinal muscle (229) 57

**Neutral drugs**
Determination of barbiturates and some neutral drugs in serum using quartz glass capillary GC (224) 449

**Nicardipine**
Specific determination of plasma nicardipine hydrochloride levels by GC-MS (223) 341

**Nicotinamide**
Determination of nicotinamide in human plasma and urine by ion-pair reversed-phase HPLC (221) 161

Determination of nicotinamide and metabolic products in urine by HPLC (230) 309

**Nicotine**
Improved GC method for the determination of nicotine and cotinine in biologic fluids (222) 61

**Niflumic acid**
Quantitative determination of niflumic acid and its β-morpholinoethyl ester in human plasma by GLC (223) 351

**Ninhydrin-positive compounds**
Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315

**Nitrate**
HPLC analysis of nitrate in urine and feces (229) 450

**Nitroglycerin**
Methods for the quantitation of nitroglycerin and its metabolites in human plasma (228) 333

**Nitroimidazoles**
LC analysis of an antimicrobial 5-nitroimidazolyl-2-sulphide derivative in biological fluids (225) 488

**Nitrosoureas**
Determination of two nitrosourea antitumor agents by chemical ionization GC-MS (229) 464

Quantitative determination of 1-(2-chloroethyl)-3-(β-D-glucopyranosyl)-1-nitrosourea in blood and urine of man (229) 481

**Nitroxoline**
HPLC analysis of nitroxoline in plasma and urine (222) 241

**Noradrenaline**
Determination of α-methyldopa, α-methylnoradrenaline, noradrenaline and adrenaline in plasma using HPLC with electrochemical detection (224) 507

Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407

Determination of catecholamines in rat heart tissue and plasma by LC with electrochemical detection (228) 143

Simultaneous determination of monoamine transmitters, precursors and metabolites in a single mouse brain (229) 35

**Norcyclizine**
Determination of cyclizine and norcyclizine in plasma and urine using GLC with nitrogen selective detection (222) 135

**Nordiazepam**
Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95

**Norepinephrine**
Novel post-column derivatization method for the fluorimetric determination of norepinephrine and epinephrine (221) 249

Amperometric determination of 3-methoxy-4-hydroxyphenylethyleneglycol in human cerebrospinal fluid (222) 112

Determination of catecholamines by radioenzymatic assay using ion-pair LC (223) 41

Studies on dopamine-converting enzymes in human plasma (223) 188

Determination of the primary metabolite of central nervous system norepinephrine, 3-methoxy-4-hydroxyphenethyleneglycol, in mouse brain and brain perfusate by HPLC with electrochemical detection (223) 295

Quantitative determination of 3-methoxy-4-hydroxyphenylethyleneglycol and its sulfate conjugate in human lumbar cerebrospinal fluid using LC with aperometric detection (223) 305

The use of $C_8$-octyl columns for the analysis of catecholamines by ion-pair reversed-phase LC with amperometric detection (225) 335

GC method for the simultaneous determination of dopamine and norepinephrine metabolites (227) 323

Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

Investigation of catecholamine metabolism using HPLC. Analytical methodology and clinical applications (229) 1

**Norlorcainide**
HPLC analysis of lorcainide and its active metabolite, norlorcainide, in human plasma (223) 454

**Normetanephrine**
Simultaneous extraction and separation of trace amines of biological interest (225) 301

**Norpseudoephedrine**
Sensitive assay for pseudoephedrine and its metabolite, norpseudoephedrine in plasma and urine using GLC with electron-capture detection (222) 297

**Northiaden**
Simultaneous measurement of dothiepin and its major metabolites in plasma and whole blood by GC-MF (222) 399

**Nortriptyline**
Simultaneous determination of amitriptyline, nortriptyline and their respective isomeric 10-hydroxy metabolites in plasma by LC (230) 391

**Noscapine**
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

**Nucleic acid bases**
Capillary GC of pyrimidines and purines: N,O-peralkyl and trifluoroacetyl-N,O-alkyl derivatives (225) 291

HPLC of nucleotides, nucleosides and bases (Review) (226) 267

Separation by TLC of nucleotides from bases and nucleosides in trichloroacetic acid extracts of cells (229) 460

**Nucleosides**
Simultaneous determination of 1-β-D-arabino-furanosylcytosine 5'-triphosphate and 3-deaza-uridine 5'-triphosphate in human leukemia cells by HPLC (221) 425

Isotachophoresis of urinary purines and pyrimidines. The use of spacers and enzymes for identification (225) 369

Analysis of adenosine, inosine and hypoxanthine in suspensions of cardiac myocytes by HPLC (226) 202

HPLC analysis of 5'-methylthioadenosine in rat tissues (226) 243

HPLC of nucleotides, nucleosides and bases (Review) (226) 267

HPLC methods for base and nucleoside analysis in extracellular fluids and in cells (226) 369

Determination of adenosine in normal human plasma and serum by HPLC (227) 168

Comparison of the reversed-phase HPLC separations of fluoropyrimidines, pyrimidines, and purines (228) 165

Reversed-phase ion-pair HPLC assay of 5-fluorouracil, 5'-deoxy-5-fluorouridine, their nucleosides, mono-, di-, and triphosphate nucleotides with a mixture of quaternary ammonium ions (228) 245

HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

Separation by TLC of nucleotides from bases and nucleosides in trichloroacetic acid extracts of cells (229) 460

Reversed-phase HPLC method to determine vidarabine and hypoxanthine arabinoside in biological fluids (229) 487

Quantitative enzymatic hydrolysis of tRNAs. Reversed-phase HPLC of tRNA nucleosides (230) 297

**Nucleotidase**
Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77

**Nucleotides**
Simultaneous determination of 1-β-D-arabino-furanosylcytosine 5'-triphosphate and 3-deaza-uridine 5'-triphosphate in human leukemia cells by HPLC (221) 425

Determination of pyridine dinucleotides in cell extracts by HPLC (225) 446

HPLC of nucleotides, nucleosides and bases (Review) (226) 267

Rapid separation of platelet nucleotides by reversed-phase, isocratic, HPLC with a radially compressed column (226) 466

HPLC assay for identification and quantitation of nucleotides in lymphocytes and malignant lymphoblasts (227) 45

Reversed-phase ion-pair HPLC assay of 5-fluorouracil, 5'-deoxy-5-fluorouridine, their nucleosides, mono-, di-, and triphosphate nucleotides with a mixture of quaternary ammonium ions (228) 245

HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

Assay of an active metabolite of 6-thioguanine, 6-thioguanosine 5'-monophosphate, in human red blood cells (229) 121

Rapid separation of nucleotides from granulocytes by isocratic, reversed-phase HPLC (229) 205

Hydrolysis of riboflavin nucleotides in plasma monitored by HPLC (229) 445

Separation by TLC of nucleotides from bases and nucleosides in trichloroacetic acid extracts of cells (229) 460

Simultaneous determination of myocardial adenine nucleotides and creatine phosphate by HPLC (230) 131

# O

**Obidoxime**
Determination of some pyridinium aldoxime compounds by means of ion-pair reversed-phase HPLC: application in biological material (225) 107

**Octadecanoic acid octadecyl ester**
Identification of some abnormal metabolites in psoriatic nail using GC-MS (221) 199

**Octopamine**
Simultaneous extraction and separation of trace amines of biological interest (225) 301

Assay for dopamine β-hydroxylase in rat serum and adrenal medulla by HPLC with fluorescence detection (227) 415

**Octopamines**
Determination of noncatecholic phenylethyl-amines and monomethylated derivatives of phenylethylamine (224) 481

**Oestradiol**
see Estradiol

**Oestrogens**
Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33

The use of oestradiol-17β antiserum covalently coupled to Sepharose to extract oestradiol-17β from biological fluids (223) 193

LC assay of urinary estriol and electrochemical detection with a battery powered detector (224) 99

HPLC analysis of estradiol-17β and metabolites in biological media (224) 205

Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

**Ophidine**
Determination of ophidine in human urine (227) 53

**Opium alkaloids**
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

**Optimization**
HPLC of phospholipids with UV detection: optimization of separations on silica (225) 319

**Organic acids**
see also carboxylic acids

Determination of the absolute configuration of some biologically important urinary 2-hydroxydicarboxylic acids by capillary GLC (222) 276

Improved procedure for the anion-exchange isolation of urinary organic acids (222) 337

High-resolution GC-MS of the methyl esters of organic acids from uremic hemofiltrates (223) 1

HPLC determination of α-keto acids (224) 197

GC-MS profile of organic acids in urine and serum of diabetic ketotic patients (225) 1

Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455

Changes of organic acids in rat heart muscle under ischemic-like conditions (227) 1

Capillary GC-MS profiles of trimethylsilyl derivatives of organic acids from amniotic fluids of different gestational age (228) 43

Study of the extraction of organic acids from urine. Preliminary step to their GC separation (in French) (228) 67

Comparison of isolation methods of urinary organic acids HPLC (228) 103

GC-MS analysis of organic acids in renal tissue biopsy. Identification of 4-hydroxybutyric acid and 4-hydroxy-2-butenoic acid (230) 1

Analysis of organic acids in the hearts of patients with idiopathic cardiomyopathy by GC-MS (230) 7

**Osalmid**
Determination of osalmid in plasma by HPLC (223) 243

**Oxalate**
Isotachophoresis for the determination of oxalate in unprocessed urine (222) 41

**Oxazepam**
Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95

**Oxeladin**
Determination of oxeladin in human sera by GLC with thermionic detection (225) 463

**Oxiconazole**
Sensitive GLC method for the determination of oxiconazole in plasma (229) 111

**Oxipurinol**
Determination of allopurinol and oxipurinol in biological fluids by HPLC (226) 237

Simultaneous determination of allopurinol, oxipurinol and uric acid in human plasma by HPLC (228) 382

Simultaneous determination of allopurinol and oxipurinol in biological fluids by mass fragmentography (229) 211

**Oxoretinoic acid**
Determination of 13-*cis*-retinoic acid and its major metabolite, 4-oxo-13-*cis*-retinoic acid, in human blood by reversed-phase HPLC (227) 471

**Oxosteroids**
Determination of 17-oxosteroids in serum and urine by fluorescence HPLC using dansyl hydrazine as a pre-labeling reagent (226) 1

**Oxyphenisatin**
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389

**Ozolinone**
Determination of etozolin and ozolinone in human plasma and tissues by reversed-phase HPLC (225) 504

# P

**Pancreatic secretion**
Isoelectric focusing studies of human pancreatic secretion (221) 279

**Papaverine**
HPLC determination of papaverine in whole blood (226) 423

Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

**Paracetamol**
Rapid GC method for emergency determination of paracetamol in human serum (221) 414

Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479

Reversed-phase chromatography of urinary metabolites of paracetamol using ion suppression and ion pairing (225) 387

HPLC determination of acetaminophen in plasma: single-dose pharmacokinetic studies (226) 224

Analysis of acetaminophen metabolites in urine by HPLC with UV and amperometric detection (227) 453

Improved acetaminophen assay sensitivity by modification of a HPLC technique (230) 443

**Paraoxon**
Quantitative analysis of the cholinesterase inhibitor paraoxon in brain tissue using HPLC (224) 133

**Parathyroid hormone**
Purification and assay of bovine parathyroid hormone by reversed-phase HPLC (223) 59

**Pareptide**
Determination of the melanotropin-inhibiting factor analogue pareptide in urine by HPLC (222) 512

**Penbutolol**
Determination of penbutolol and its hydroxylated metabolite in biological fluids by reversed-phase HPLC (228) 355

**Penicillamine**
Determination of D-penicillamine in serum by fluorescence derivatization and LCC (222) 249

HPLC determination of penicillamine in whole blood, plasma, and urine (222) 445

Determination of penicillamine and other thiols by combined HPLC and post-column reaction with Ellman's reagent: application to human urine (226) 498

**Penicillins**
Determination of amdinocillin in plasma and urine by high-performance liquid chromatography (227) 137

Improved procedure for the determination of the ureidopenicillins azlocillin and mezlocillin in plasma by HPLC (228) 409

**Pentamethylmelamine**
HPLC analysis of pentamethylmelamine and its metabolites in biological fluids (222) 518

**Pentazirinocyclodiphosphathiazene**
Determination of pentazirinocyclodiphosphathiazene in biological fluids by HPLC (230) 188

**Pentazocine**
HPLC analysis of pentazocine in blood and plasma (227) 239

**Pentobarbital**
Simple, rapid and sensitive reversed-phase HPLC method for thiopental and pentobarbital determination in plasma and brain tissue (227) 207

**Pentobarbitone**
Analysis of barbiturates in blood by HPLC (226) 117

**Pentoxifylline**
Simultaneous determination of pentoxifylline and its hydroxy metabolite in plasma by HPLC (225) 261

**Peptides**
Gel chromatographic separation of human C-peptide and proinsulin (222) 33

Characterization and determination of neuropeptides by HPLC and radioimmunoassay (222) 371

Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381

Determination of the melanotropin-inhibiting factor analogue pareptide in urine by HPLC (222) 512

Purification and assay of bovine parathyroid hormone by reversed-phase HPLC (223) 59

Resolution of 52 ninhydrin-positive compounds with a high-speed amino acid analyser. Determination of carnosine and homocarnosine in biological materials (224) 315

Analysis of middle molecular peptides in normal and uremic body fluids by HPLC (224) 322

Hydrophilic ion-pair reversed-phase chromatography of biogenic peptides prior to immunoassay (224) 472

Polymorphism of urinary 4-hydroxyproline-containing polypeptides (225) 55

Determination of ophidine in human urine (227) 53

Analysis of small peptides in uremic serum by HPLC (228) 292

Rapid high-yield purification of neuropeptides from canine intestinal muscle (229) 57

Separation of peptides by cellulose-phosphate chromatography for identification of a hemoglobin variant (229) 193

HPLC purification of endogenous tumour cell growth inhibitory peptides (230) 289

**Pertechnetate**
Quantitative determination of pertechnetate by HPLC with UV detection (227) 249

**Pharmacokinetic drug monitoring**
Fully automated HPLC. A new chromatograph for pharmacokinetic drug monitoring by direct injection of body fluids (222) 13

**Phenacetin**
Rapid TLC of various weak analgesics in saliva (225) 427

**Phenazone**
Rapid TLC of various weak analgesics in saliva (225) 427

**Phencyclidine**
Simultaneous determination of phencyclidine and monohydroxylated metabolites in urine of man by GC-MF with methane chemical ionization (223) 331

**Phenelzine**
Determination of phenelzine in human plasma with GC-MS using an isotope labeled internal standard (221) 301

**Phenobarbital**
Rapid quantitative determination of underivatized carbamazepine, phenytoin, phenobarbital and *p*-hydroxyphenobarbital in biological fluids by packed column GC (221) 75

Rapid microanalysis of anticonvulsants by HPTLC (222) 265

Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

Simultaneous assay of methylphenobarbital and phenobarbital in plasma using GC-MS with selected ion monitoring (223) 426

Simultaneous microdetermination of theophylline, caffeine and phenobarbital in blood collected on paper (224) 439

Chromatographic micro-procedure for trace determination of phenobarbital in blood serum (229) 365

**Phenolphthalein**
Qualitative and quantitative analysis of some synthetic, chemically acting laxatives in urine by GC-MS (222) 389

**Phenoperidine**
Quantitative GLC method for the determination of phenoperidine in human plasma (223) 213

**Phenothiazines**
GLC determination of pipotiazine in plasma of psychiatric patients (222) 291

Sensitive GC determination of trifluoperazine in human plasma (223) 95

Simultaneous determination of chlorpromazine and levomepromazine in human plasma and urine by HPLC using electrochemical detection (227) 103

Analysis of plasma trifluoperazine by GC and selected ion monitoring (228) 321

HPLC determination of pipotiazine in human plasma and urine (230) 401

**Phentolamine**
GC determination of phentolamine (Regitine) in human plasma and urine (222) 429

Determination of the major metabolite of phentolamine in human plasma and urine by HPLC (222) 461

**Phenylacetic acid**
Longitudinal urinary excretion of some "trace" acids in a human male (222) 161

Plasma concentrations of *p*- and *m*-hydroxyphenylacetic acid and phenylacetic acid in humans. GC-high-resolution MS analysis (230) 219

**Phenyl acetic mustard**
Rapid assay for plasma chlorambucil and phenyl acetic mustard using reversed-phase LC (230) 381

**Phenylalanine**
Analysis of some tryptophan and phenylalanine metabolites in urine by a straight-phase HPLC technique (222) 191

**Phenylethanolamine N-methyltransferase**
Highly sensitive assay for phenylethanolamine N-methyltransferase activity in rat brain by HPLC with electrochemical detection (227) 407

**Phenylethylamine**
Simultaneous extraction and separation of trace amines of biological interest (225) 301

**Phenylethylamines**
Determination of noncatecholic phenylethylamines and monomethylated derivatives of phenylethylamine (224) 481

Determination of the α,β-adrenoceptor blocker YM-09538 in plasma by HPLC with fluorescence detection (225) 115

**Phenylethylhydrazine**
Determination of phenelzine in human plasma with GC-MS using an isotope labeled internal standard (221) 301

**Phenylpyruvic acid**
Determination of phenylpyruvic acid in urine and serum by HPLC with fluorescence detection (226) 25

**Phenytoin**
Rapid quantitative determination of underivatized carbamazepine, phenytoin, phenobarbital and *p*-hydroxyphenobarbital in biological fluids by packed column GC (221) 75

Rapid microanalysis of anticonvulsants by HPTLC (222) 265

Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

**Phosphodiesterase**
Rapid assay of cyclic AMP phosphodiesterase and 5'-nucleotidase by means of chromatography on cellulose-nitrate membrane strips (229) 77

**Phospholipids**
HPLC determination of the lecithin/sphingomyelin ratio in amniotic fluid (223) 277

Lipid class and molecular species interrelationships among plasma lipoproteins of normolipemic subjects (224) 1

HPLC of phospholipids with UV detection: optimization of separations on silica (225) 319

Nano-scale densitometric quantitation of phospholipids (225) 450

Improved TLC assay for monitoring lecithin/sphingomyelin ratios in amniotic fluid (226) 484

Improved procedure for the separation of phospholipids by HPLC (227) 25

Rapid determination of lipids in healthy human lymphocytes (227) 423

Lipids in human neutrophils determined by a microanalytical method (229) 455

**Physostigmine**
Quantitative determination of the cholinesterase inhibitor physostigmine in brain tissue samples using reversed-phase HPLC (225) 381

**Pindolol**
Determination of pindolol in biological fluids by an electron-capture GLC method on a wall-coated open tubular column (221) 387

**Pipotiazine**
GLC determination of pipotiazine in plasma of psychiatric patients (222) 291

HPLC determination of pipotiazine in human plasma and urine (230) 401

**Pirmenol**
HPLC analysis of a new antiarrhythmic drug, pirmenol, in biological fluids (224) 343

**Plasticizers**
GC-MS detection of circulating plasticizers in surgical patients (222) 171

GC quantitation of two plasticizers contaminating intravenous fluids stored in plastic containers (225) 196

**Platinum complexes**
HPLC of platinum complexes on solvent generated anion exchangers. III. Application to the analysis of cisplatin in urine using automated column switching (229) 373

**Polyamines**
HPLC procedure for the simultaneous determination of the natural polyamines and their monoacetyl derivatives (221) 227

Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363

Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51

Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198

HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

Rapid assay of spermidine synthase activity by HPLC (226) 208

Reversed-phase ion-pair LC procedure for the simultaneous analysis of S-adenosylmethionine, its metabolites and the natural polyamines (227) 349

Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369

GC-MS analysis of amine metabolites in urine (228) 51

Single-step HPLC method for the analysis of acetylated polyamines (229) 47

MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142

**Polyols**
Myo-inositol levels in the cerebrospinal fluid of infants (223) 173

**Polypeptides**
Separation of collagen types I and III by HPLC (222) 284

Polymorphism of urinary 4-hydroxyproline-containing polypeptides (225) 55

High-performance gel permeation chromatography of collagens (230) 409

**Polythiazide**
Determination of polythiazide and prazosin in human plasma by HPLC (224) 33

**Porphyrins**
Normal-phase HPLC of porphyrin free acids on silica modified with tetraethylenepentamine (228) 305

The determination of uroporphyrinogen decarboxylase in tissues by HPLC coupled to fluorescence detection (229) 311

Separation and quantitation of urinary porphyrins by HPLC (229) 439

**Prazosin**
Quantification of prazosin in plasma by HPLC (221) 188

Determination of polythiazide and prazosin in human plasma by HPLC (224) 33

**Prazosin analogue**
Determination of the vasodilator UK33274 by HPLC using fluorescence detection (221) 193

**Prednisolone**
Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221

Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283

**Prednisone**
Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221

Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283

**Pregnanes**
Simple computer program for a low-cost desk-top calculator applied to the evaluation of GLC analyses of 17-ketosteroids and pregnanes (223) 253

**Pregnenedione**
Natural products chemistry. 81. Analysis of rat thymus steroids by liquid-gel chromatography and GC-MS (222) 475

**Pregnenolone**
Quantitation of pregnenolone and 17-hydroxy-pregnenolone in human serum by automatic HPLC and subsequent radioimmunoassay (224) 229

Combined PC and reversed-phase HPLC method for the study of pregnenolone and progesterone metabolites (224) 399

**Prenalterol**
Determination of prenalterol in plasma and in urine by GLC (222) 437

Assay of prenalterol in plasma and urine by GC-MS (229) 87

**Primaquine**
Electron-capture GC assay for primaquine in blood (224) 265

HPLC analysis of the metabolism of primaquine and the identification of a new mammalian metabolite (230) 69

**Primidone**
Rapid microanalysis of anticonvulsants by HPTLC (222) 265

Rapid determination of anticonvulsant drugs by isothermal GLC (222) 486

**Probenecid**
The simultaneous analysis of clofibric acid and probenecid and the direct analysis of clofibric acid glucuronide by HPLC (223) 147

Rapid LC determination of probenecid in human cerebrospinal fluid (225) 521

Extractive alkylation of probenecid in plasma and cerebrospinal fluid and determination by electron-capture GC (228) 340

**Procarbazine**
Analysis of procarbazine and metabolites by GC-MS (221) 309

Quantitative analysis of procarbazine, procarbazine metabolites and chemical degradation products with application to pharmacokinetic studies (229) 397

**Procetofenic acid**
Improved method for the determination of procetofenic acid in human plasma by GLC (224) 500

Rapid determination of procetofenic acid in plasma by HPLC (227) 219

**Procyclidine**
Analysis of procyclidine in human plasma and urine by GLC (221) 408

**Progabide**
GC method for the determination of progabide (SL 76.002) in biological fluids (230) 154

**Progesterone**
Combined PC and reversed-phase HPLC method for the study of pregnenolone and progesterone metabolites (224) 399

Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17*α*-hydroxyprogesterone from small plasma or tissue samples (228) 95

**Proinsulin**
Gel chromatographic separation of human C-peptide and proinsulin (222) 33

**Prolintane**
Assay of prolintane in plasma by capillary GC with nitrogen-selective detection (222) 129

**Promethazine**
Simultaneous determination of promethazine and two of its circulating metabolites by HPLC (230) 194

**Propafenone**
HPLC analysis of propafenone in human plasma samples (230) 448

**Propranolol**
Simultaneous determination of *d*- and *l*-propranolol in human plasma by HPLC (221) 109

HPLC method for the quantitative analysis of the aryloxypropanolamines propranolol, metoprolol and atenolol in plasma and tissue (228) 223

Simple and rapid determination of propranolol and its active metabolite, 4-hydroxypropranolol, in human plasma by LC with fluorescence detection (228) 362

**Propyphenazone**
GC determination of N-desmethyl-propyphenazone, a metabolite of propyphenazone, in human urine (221) 377

Rapid TLC of various weak analgesics in saliva (225) 427

**Prostacyclin**
Separation of six major prostacyclin metabolites by HPLC (230) 115

**Prostaglandins**
Quantitative determination of 6-keto prostaglandin $F_{1\alpha}$ in biological fluids by capillary GC-chemical ionization MS (221) 361

Radioimmunoassay of prostaglandin $F_{2\alpha}$ using Sephadex G-75 gel chromatography (225) 47

Improved sample preparation for the quantitative MS determination of prostaglandins in biological samples (226) 450

Determination of prostaglandins and thromboxane as their pentafluorobenzyl-trimethylsilyl derivatives by electron-capture GC (230) 15

Separation of six major prostacyclin metabolites by HPLC (230) 115

**Proteases**
Isoelectric focusing studies of human pancreatic secretion (221) 279

**Protein distribution**
Detection of the changes in protein distribution of rat plasma induced by carbon tetrachloride administration by means of two-dimensional electrophoresis (225) 65

**Protein profiling**
Urinary protein profiling by high-performance gel permeation chromatography (230) 47

**Protein removing**
Evaluation of the relative efficacy of various techniques for deproteinizing plasma samples prior to HPLC (226) 455

**Proteins**
see also enzymes

High-performance aqueous gel permeation chromatography of human serum lipoproteins (221) 257

Carrier-ampholyte displacement chromatography (chromatofocusing) on ion-exchange papers. Application to the separation of haemoglobin variants (222) 116

A new method for detection of drug-binding proteins using a parallel-flow dialysis technique (223) 267

High-performance aqueous gel permeation chromatography of serum lipoproteins:

selective detection of cholesterol by enzymatic reaction (223) 285

Lipid class and molecular species inter-relationships among plasma lipoproteins of normolipemic subjects (224) 1

Analysis of middle molecular peptides in normal and uremic body fluids by HPLC (224) 322

Separation of human C-apolipoproteins by HPLC (225) 185

Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

Demonstration of the formation of renin and renin-binding protein complex using HPLC (225) 329

Rapid purification of proteolipids from rat brain subcellular fractions by chromatography on a lipophilic dextran gel (227) 33

An update on electrophoretic and chromatographic methods in the diagnosis of hemoglobinopathies (Review) (227) 267

Separation and measurement of isoenzymes and other proteins by HPLC (Review) (228) 1

Chromatofocusing of human hemoglobins: application to the quantitation of hemoglobin $A_2$ (228) 177

Rapid separation of human globin chains by high flow-rate chromatography under low pressure using CM-Trisacryl M (228) 298

Separation of peptides by cellulose-phosphate chromatography for identification of a hemoglobin variant (229) 193

Rapid method for the purification of the elastin cross-links, desmosine and isodesmosine (229) 200

Microprocedure to determine the polymorphic forms of acid $\alpha_1$-glycoprotein in plasma. Application to depressive patients treated with amitriptyline (229) 319

Urinary protein profiling by high-performance gel permeation chromatography (230) 47

Chromatofocusing of human hemoglobins (230) 137

The effect of ampholytes on ferritin isoelectric focussing patterns (230) 327

**Proteolipids**
Rapid purification of proteolipids from rat brain subcellular fractions by chromatography on a lipophilic dextran gel (227) 33

**Prothionamide**
HPLC determination of ethionamide and prothionamide in body fluids (225) 245

**Pseudoephedrine**
Sensitive assay for pseudoephedrine and its metabolite, norpseudoephedrine in plasma and urine using GLC with electron-capture detection (222) 297

**Pteridines**
Determination of neopterine in human urine by reversed-phase HPLC (227) 61

**Purines**
Capillary GC of pyrimidines and purines: N,O-peralkyl and trifluoroacetyl-N,O-alkyl derivatives (225) 291

Isotachophoresis of urinary purines and pyrimidines. The use of spacers and enzymes for identification (225) 369

HPLC methods for base and nucleoside analysis in extracellular fluids and in cells (226) 369

HPLC determination of plasma 6-mercaptopurine in clinically relevant concentrations (227) 526

Comparison of the reversed-phase HPLC separations of fluoropyrimidines, pyrimidines, and purines (228) 165

HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

**Putrescine**
Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363

Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51

Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198

HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369

GC-MS analysis of amine metabolites in urine (228) 51

Single-step HPLC method for the analysis of acetylated polyamines (229) 47

**Pyridine dinucleotides**
Determination of pyridine dinucleotides in cell extracts by HPLC (225) 446

**Pyridinium aldoximes**
Determination of some pyridinium aldoxime compounds by means of ion-pair reversed-phase HPLC: application in biological material (225) 107

**Pyridostigmine**
Method for isolation and determination of pyridostigmine and metabolites in urine and blood (228) 235

**Pyrimethamine**
Contribution to the determination of four antimalaria drugs, chloroquine, quinine, pyrimethamine and sulfadoxine, alone or in mixtures in biological fluids. II. GC (in French) (224) 332

Measurement of pyrimethamine in human plasma by GLC (224) 492

Determination of pyrimethamine in human plasma after administration of Fansidar or Fansidar-mefloquine by means of HPLC with fluorescence detection (230) 107

**Pyrimidines**
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382

Determination of 1-β-D-arabinofuranosyl-cytosine and 1-β-D-arabinofuranosyluracil in human plasma by HPLC (223) 371

Capillary GC of pyrimidines and purines: N,O-peralkyl and trifluoroacetyl-N,O-alkyl derivatives (225) 291

Isotachophoresis of urinary purines and pyrimidines. The use of spacers and enzymes for identification (225) 369

HPLC methods for base and nucleoside analysis in extracellular fluids and in cells (226) 369

Comparison of the reversed-phase HPLC separations of fluoropyrimidines, pyrimidines, and purines (228) 165

HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

**Pyruvate**
Determination of pyruvate, lactate, acetoacetate, and 3-hydroxybutyrate in serum by capillary isotachophoresis (225) 455

# Q

**Quinalbarbitone**
Analysis of barbiturates in blood by HPLC (226) 117

**Quinidine**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

Specific determination of quinidine and metabolites in biological fluids by reversed-phase HPLC (228) 366

Simultaneous HPLC assay for quinidine, disopyramide and the mono-N-dealkylated metabolite of disopyramide (230) 433

**Quinidine analogues**
Determination of the quinidine analog, 7'-trifluoromethyldihydrocinchonidine·2HCl in plasma and urine by HPLC (221) 97

**Quinine**
Contribution to the determination of four antimalaria drugs, chloroquine, quinine, pyrimethamine and sulfadoxine, alone or in mixtures in biological fluids. II. GC (in French) (224) 332

# R

**Ranitidine**
Determination of ranitidine and its metabolites in human urine by reversed-phase ion-pair HPLC (225) 161

**Regitine**
GC determination of phentolamine (Regitine) in human plasma and urine (222) 429

Determination of the major metabolite of phentolamine in human plasma and urine by HPLC (222) 461

**Renin-binding protein**
Demonstration of the formation of renin and renin-binding protein complex using HPLC (225) 329

**Resorcinol**
HPLC determination of free resorcinol in plasma and in urine (224) 513

**Retinoic acid**
Determination of 13-*cis*-retinoic acid and its major metabolite, 4-oxo-13-*cis*-retinoic acid, in human blood by reversed-phase HPLC (227) 471

**Retinoic acid analogues**
Determination of plasma levels of two aromatic retinoic acid analogues with antipsoriatic activity by HPLC (221) 345

**Riboflavin**
LC determination and time-concentration studies of riboflavin in hemodialysate from uremic patients (226) 471

**Riboflavin nucleotides**
Hydrolysis of riboflavin nucleotides in plasma monitored by HPLC (229) 445

**Ribonucleic acids**
see RNA

**Ribonucleosides**
HPLC determination of purine and pyrimidine bases, ribonucleosides, deoxyribonucleosides and cyclic ribonucleotides in biological fluids (229) 67

**Rifampicin**
Determination of three main antileprosy drugs and their main metabolites in serum by HPLC (223) 379

Quantitative assay of rifampicin and its main metabolite 25-desacetylrifampicin in human plasma by reversed-phase HPLC (225) 526

**tRNA**
Quantitative enzymatic hydrolysis of tRNAs. Reversed-phase HPLC of tRNA nucleosides (230) 297

**RNAase**
Isoelectric focusing studies of human pancreatic secretion (221) 279

**Ro12-0068**
Determination of Ro12-0068, a new anti-inflammatory and analgesic compound, in plasma by means of HPLC (225) 493

# S

**Salbutamol**
A time-saving method for the determination of the $\beta_2$ sympathomimetics terbutaline, salbutamol and fenoterol. Preliminary results (229) 217

**Salicylamide**
Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479

**Salicylate**
Simultaneous determination of acetaminophen, theophylline and salicylate in serum by HPLC (229) 492

**Salicylazosulphapyridine**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

Simplified HPLC method for 5-aminosalicylic acid in plasma and urine (225) 498

Assay of 5-aminosalicylate and its acetylated metabolite in biological fluids by HPLC on dynamically modified silica (226) 504

**Salicylhydroxamic acid**
Determination of salicylhydroxamic acid, a trypanocidal agent, by reversed-phase HPLC (225) 236

**Salicylic acid**
Rapid HPTLC of salicylic acid, salicylamide, ethoxybenzamide and paracetamol in saliva (223) 479

LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

A method for HPLC screening of UV-positive components in urine eluate from Sephadex G-10 and modifications for determination of urinary salicylic, salicyluric and gentisic acids (230) 253

**Salicyluric acid**
A method for HPLC screening of UV-positive components in urine eluate from Sephadex G-10 and modifications for determination of urinary salicylic, salicyluric and gentisic acids (230) 253

**Sample preparation**
Evaluation of the relative efficacy of various techniques for deproteinizing plasma samples prior to HPLC (226) 455

**SC-28538**
LC analysis of an antimicrobial 5-nitroimidazolyl-2-sulphide derivative in biological fluids (225) 488

**Selenium**
Improved method for selenium determination in biological samples by GC (225) 9

**Sep-Pak cartridges**
Evaluation of $C_{18}$ Sep-Pak cartridges for biological sample clean-up for tricyclic antidepressant assays (225) 189

**Serotonin**
Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

Determination of serotonin, its precursors, metabolites and [$^3$H]serotonin in lung by HPLC with fluorescence detection (225) 27

Pre-column derivatization and HPLC of biogenic amines in blood of normal and malignant hyperthermic pigs (228) 113

Simultaneous measurement of 5-hydroxytryptophan and L-dihydroxyphenylalanine by HPLC with electrochemical detection. Measurement of serotonin and catecholamine turnover in discrete brain regions (228) 123

Simultaneous determination of norepinephrine, dopamine and serotonin in rat brain regions by ion-pair LC on octyl silane columns and amperometric detection (228) 131

Determination of indoles in human and rat pineal (228) 155

**Sialic acids**
HPLC quantitation of N-acetylneuraminic acid in malignant melanoma and breast carcinoma (224) 381

**Sialidase**
Rapid and sensitive sialidase assay by HPLC and its application to detection of sialidase in human urine (230) 415

**Sialoglycoproteins**
Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

**Sisomicin**
Improved microdetermination of gentamicin and sisomicin in serum by HPLC with ultraviolet detection (222) 316

**Sodium flavodate**
Determination of sodium flavodate in body fluids by HPLC. Application to clinical pharmacokinetic studies (230) 168

**Somatostatin**
Rapid high-yield purification of neuropeptides from canine intestinal muscle (229) 57

**Spermidine**
Determination of monoacetyldiamines and -polyamines in urine by HPLC (222) 363

Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51

Determination of polyamines by ion-exchange TLC and video-densitometry (223) 198

HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

Reversed-phase ion-pair LC procedure for the simultaneous analysis of S-adenosylmethionine, its metabolites and the natural polyamines (227) 349

Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369

GC-MS analysis of amine metabolites in urine (228) 51

Single-step HPLC method for the analysis of acetylated polyamines (229) 47

MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142

**Spermidine synthase**
Rapid assay of spermidine synthase activity by HPLC (226) 208

**Spermine**
Measurements of polyamines and their acetylated derivatives in cell extracts and physiological fluids by use of an amino acid analyzer (223) 51

HPLC in the separation and measurement of di- and polyamines and their derivatives, and specific preparation of isomers of their monoacetyl derivatives (224) 371

Chromophoric determination of putrescine, spermidine and spermine with dabsyl chloride by HPLC and TLC (227) 369

MS identification of isoputreanine, a metabolite of spermidine and/or spermine, in human urine (230) 142

**Sphingomyelin**
HPLC determination of the lecithin/sphingomyelin ratio in amniotic fluid (223) 277

Improved TLC assay for monitoring lecithin/sphingomyelin ratios in amniotic fluid (226) 484

**Spirorenone**
Determination of plasma levels of spirorenone, a new aldosterone antagonist, and one of its metabolites by HPLC (230) 37

**Steroids**
Natural products chemistry. 81. Analysis of rat thymus steroids by liquid-gel chromatography and GC-MS (222) 475

Reliability of the estimation of serum cortisol by HPLC (222) 478

Steroid profiles of healthy individuals (223) 21

Studies on steroids. CLXV. Determination of isomeric catechol estrogens in pregnancy urine by HPLC with electrochemical detection (223) 33

The use of oestradiol-17β antiserum covalently coupled to Sepharose to extract oestradiol-17β from biological fluids (223) 193

Simple computer program for a low-cost desk-top calculator applied to the evaluation of GLC analyses of 17-ketosteroids and pregnanes (223) 253

LC assay of urinary estriol and electrochemical detection with a battery powered detector (224) 99

Enzymatic detection of urinary acidic 3α-hydroxysteroids on thin-layer chromatograms (224) 111

HPLC analysis of estradiol-17β and metabolites in biological media (224) 205

Analysis of prednisone, prednisolone and their 20β-hydroxylated metabolites by HPLC (224) 221

Quantitation of pregnenolone and 17-hydroxy-pregnenolone in human serum by automatic HPLC and subsequent radioimmunoassay (224) 229

Analysis of profiles of conjugated steroids in urine by ion-exchange separation and GC-MS (224) 355

Combined PC and reversed-phase HPLC method for the study of pregnenolone and progesterone metabolites (224) 399

Determination of minaxolone (CCI 12923) by GC with nitrogen-sensitive detection (224) 496

Determination of serum unconjugated estrone, estradiol-17β and estriol during pregnancy by selected ion monitoring (225) 275

Determination of 17-oxosteroids in serum and urine by fluorescence HPLC using dansyl hydrazine as a pre-labeling reagent (226) 1

Studies on steroids. CLXX. Separation and determination of bile acid 3-sulfates in human bile by HPLC (226) 13

Sensitive GC method for the determination of alphadolone in plasma (226) 213

Separation of some natural and synthetic corticosteroids in biological fluids and tissues by HPLC (227) 11

Determination of urinary dehydroepian-drosterone sulphate by combined HPLC and radioimmunoassay (227) 485

Estimation of free urinary aldosterone and 18-hydroxycorticosterone by a combination of automatic HPLC and radioimmunoassay (227) 492

Celite multi-column chromatography for the simultaneous separation of progesterone, deoxycorticosterone and 17α-hydroxyprogesterone from small plasma or tissue samples (228) 95

Enzymatic determination of urinary 17β-hydroxysteroids on thin-layer chromatograms (228) 317

Sample purification using a $C_{18}$-bonded reversed-phase cartridge for the quantitative analysis of corticosteroids in adrenal cell cultures by HPLC or GC-MS (229) 267

Simultaneous measurement of prednisone, prednisolone and 6β-hydroxyprednisolone in urine by HPLC coupled with a radioactivity detector (229) 283

Determination of methylprednisolone in central nervous tissue and plasma using normal-phase HPLC (230) 29

Determination of plasma levels of spirorenone, a new aldosterone antagonist, and one of its metabolites by HPLC (230) 37

Enzymatic detection of urinary conjugated steroids after gel chromatography (230) 263

**Sterols**
Simultaneous measurement of sterol and fatty acid composition in small samples (221) 367

**Substance P**
Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381

**Sufentanil**
Rapid and sensitive GC method for the determination of alfentanil and sufentanil in biological samples (224) 122

**Sugar alcohols**
Estimation of sugar alcohols by GLC using a modified acetylation procedure (226) 191

**Sugars**
see also carbohydrates

Determination of carbohydrates in urine by HPLC and optical activity detection (223) 321

Modification of the simultaneous determination of alditol acetates of neutral and aminosugars by GLC. Application to the fractionation of sialoglycoproteins from bone (225) 309

**Sulindac**
Measurement of sulindac and its metabolites in human plasma and urine by HPLC (225) 123

**Sulphadiazine**
Assay of trimethoprim, sulfadiazine and its $N^4$-acetyl metabolite in biological fluids by normal-phase HPLC (224) 59

Determination of 2,4-diamino-5-benzylpyrimi-dines in combination with sulphadiazine in biological fluids using HPLC (224) 301

**Sulphadoxine**
Contribution to the determination of four antimalaria drugs, chloroquine, quinine, pyrimethamine and sulfadoxine, alone or in mixtures in biological fluids. II. GC (in French) (224) 332

**Sulphamethoxazole**
Simultaneous determination of trimethoprim, sulphamethoxazole and $N^4$-acetylsulpha-methoxazole in serum and urine by HPLC (223) 139

**Sulphate**
Microassay of inorganic sulfate in biological fluids by controlled flow anion chromatography (225) 359

**Sulphated bile acids**
Studies on steroids. CLXX. Separation and determination of bile acid 3-sulfates in human bile by HPLC (226) 13

**Sulphathiazole**
HPLC determination of sulphathiazole in human plasma and urine (221) 419

**Sulphinpyrazone**
Simultaneous determination of sulfinpyrazone and four of its metabolites by HPLC (223) 460

HPLC analysis of the sulfide metabolite of sulfinpyrazone in plasma (227) 516

**Sulphonamides**
HPLC determination of sulphathiazole in human plasma and urine (221) 419

# T

**Taurine**
Separation of taurine and glycerophosphoryl ethanolamine on amino acid analyzers (223) 179

**Technetium-99m**
Quantitative determination of pertechnetate by HPLC with UV detection (227) 249

**Temazepam**
Determination of diazepam and its major metabolites in man and in the cat by HPLC (222) 95

Plasma concentrations of temazepam, a 3-hydroxy benzodiazepine, determined by electron-capture GLC (222) 125

Assay of flunitrazepam, temazepam and desalkylflurazepam in plasma by capillary GC with electron-capture detection (227) 199

**Teniposide**
Analysis of the anticancer drugs etoposide (VP 16-213) and teniposide (VM 26) by HPLC with fluorescence detection (224) 168

**Terbutaline**
A time-saving method for the determination of the $\beta_2$ sympathomimetics terbutaline, salbutamol and fenoterol. Preliminary results (229) 217

**Tetrabenazine**
Determination of therapeutic plasma concentrations of tetrabenazine and an active metabolite by HPLC (226) 175

**Tetracycline**
Method for rapid determination of urinary tetracycline by HPLC (222) 152

**Tetradecanoic acid octadecyl ester**
Identification of some abnormal metabolites in psoriatic nail using GC-MS (221) 199

**4,5,6,7-Tetrahydroisoxazolo[5,4-*c*]pyridine-3-ol**
Determination of the $\gamma$-aminobutyric acid agonist 4,5,6,7-tetrahydroisoxazolo[5,4-*c*]pyridine-3-ol in urine by HPLC (228) 279

**Tetroxoprim**
Determination of 2,4-diamino-5-benzylpyrimidines in combination with sulphadiazine in biological fluids using HPLC (224) 301

Quantitative TLC of trimethoprim and tetroxoprim using fluorescence densitometry (230) 212

**Thebaine**
Detection and measurement of opium alkaloids and metabolites in urine of opium eaters by methane chemical ionization mass fragmentography (230) 57

**Theobromine**
Reversed-phase HPLC determination of caffeine and its N-demethylated metabolites in dog plasma (226) 231

**Theophylline**
Simultaneous determination of theophylline and its major metabolites in urine by reversed-phase ion-pair HPLC (221) 85

Simultaneous determination of theophylline and caffeine after extractive alkylation in small volumes of plasma by GC-MS (221) 166

Simultaneous determination of dyphylline and theophylline in human plasma by HPLC (221) 170

Determination of plasma theophylline by straight-phase HPLC: elimination of interfering caffeine metabolites (222) 467

Simultaneous microdetermination of theophylline, caffeine and phenobarbital in blood collected on paper (224) 439

Measurement of plasma theophylline by GLC on the stationary phase SP-2510 (225) 208

Reversed-phase HPLC determination of caffeine and its N-demethylated metabolites in dog plasma (226) 231

Simultaneous determination of acetaminophen, theophylline and salicylate in serum by HPLC (229) 492

**Thiabendazole**
Determination of thiabendazole and 5-hydroxy-thiabendazole in human serum by fluorescence-detected HPLC (230) 79

**Thiadiazolecarboxaldoxime**
HPLC separation, isolation and identification of 1,2,3-thiadiazole-5-carboxaldoxime glucuronide in rabbit urine (226) 165

**Thiazides**
Determination of trichlormethiazide in human plasma and urine by HPLC (223) 466

**Thioguanine**
Reversed-phase HPLC analysis of 6-thioguanine applicable to pharmacologic studies in humans (227) 83

**Thioguanosine monophosphate**
Assay of an active metabolite of 6-thioguanine, 6-thioguanosine 5'-monophosphate, in human red blood cells (229) 121

**Thiols**
Determination of penicillamine and other thiols by combined HPLC and post-column reaction with Ellman's reagent: application to human urine (226) 498

Reversed-phase ion-pair HPLC of mercaptoacetate and N-acetylcysteine after derivatization with N-(1-pyrene)maleimide and N-(7-dimethylamino-4-methyl-3-coumarinyl)maleimide (229) 409

**Thiopental**
Simple, rapid and sensitive reversed-phase HPLC method for thiopental and pentobarbital determination in plasma and brain tissue (227) 207

**Thormählen-positive compounds**
Identification of two Thormählen-positive compounds from melanotic urine by GC-MS (222) 329

**Thromboxane**
Determination of prostaglandins and thromboxane as their pentafluorobenzyl-trimethylsilyl derivatives by electron-capture GC (230) 15

**Thymine**
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382

**Thyroid hormones**
Simultaneous determination of D- and L-thyroxine in human serum by LC with electrochemical detection (226) 383

**Thyronines**
Catalytic detection principle for HPLC: determination of enantiomeric iodinated thyronines in blood serum (224) 239

**Thyrotropin-releasing hormone**
Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381

**Thyroxine**
Catalytic detection principle for HPLC: determination of enantiomeric iodinated thyronines in blood serum (224) 239

Simultaneous determination of D- and L-thyroxine in human serum by LC with electrochemical detection (226) 383

**Ticrynafen**
Quantitative TLC determination of ticrynafen in plasma of the dog (222) 324

**Tienilic acid**
Determination of ethacrynic and tienilic acid in plasma by GLC-MS (227) 193

**Tiodazosin**
Determination of tiodazosin in plasma and whole blood by HPLC (230) 203

**Tobramycin**
Micro-determination of tobramycin in serum by HPLC with UV detection (225) 417

**Tocainide**
Simple method for the measurement of tocainide and lignocaine in blood plasma or serum using GLC with flame ionisation detection (225) 469

Electron-capture GLC determination of tocainide in biological fluids using fused silica capillary columns (229) 103

**Tocopherols**
Quantitation of individual tocopherols in plasma, platelets, lipids, and livers by HPLC (224) 389

Improved assay for $\alpha$-tocopherol in the picogram range, using GC-MS (225) 433

Quantitation of physiological $\alpha$-tocopherol, metabolites, and related compounds by reversed-phase HPLC (227) 174

**Tofisopam**
Determination of tofisopam in serum by HPLC (222) 303

**Triamcinolone acetonide**
Separation of some natural and synthetic corticosteroids in biological fluids and tissues by HPLC (227) 11

**Triazolam**
Assay of triazolam in plasma by capillary GC (223) 438

Electron-capture GC analysis of the triazolobenzodiazepines alprazolam and triazolam (225) 202

Simultaneous determination of triazolo-benzophenone [2',5-dichloro-2-(3-glycylamino-methyl-5-methyl-4H-1,2,4-triazol-4-yl)-benzophenone] and its major blood metabolite,

triazolam, in monkey plasma by electron-capture GLC (229) 355

**Triazolobenzodiazepines**
Assay of triazolam in plasma by capillary GC (223) 438

Electron-capture GC analysis of the triazolobenzodiazepines alprazolam and triazolam (225) 202

**Triazolobenzophenones**
Simultaneous determination of triazolo-benzophenone [2',5-dichloro-2-(3-glycylamino-methyl-5-methyl-4H-1,2,4-triazol-4-yl)-benzophenone] and its major blood metabolite, triazolam, in monkey plasma by electron-capture GLC (229) 355

**Trichlormethiazide**
Determination of trichlormethiazide in human plasma and urine by HPLC (223) 466

**Tricyclic antidepressant drugs**
Evaluation of $C_{18}$ Sep-Pak cartridges for biological sample clean-up for tricyclic antidepressant assays (225) 189

**Trifluoperazine**
Sensitive GC determination of trifluoperazine in human plasma (223) 95

Analysis of plasma trifluoperazine by GC and selected ion monitoring (228) 321

**Trifluoromethyldihydrocinchonidine**
Determination of the quinidine analog, 7'-trifluoromethyldihydrocinchonidine·2HCl in plasma and urine by HPLC (221) 97

**Triglycerides**
Studies of hyperlipidemia in drug-induced diabetic rats by HPLC (221) 19

Rapid determination of lipids in healthy human lymphocytes (227) 423

**Trimebutine**
Quantitative HPLC determination of antispasmodic trimebutine in human plasma. Pharmacokinetic studies after intravenous administration in humans (224) 149

**Trimetazidine**
Trace determination of trimetazidine in plasma by HPLC using fluorescence detection (224) 162

**Trimethobenzamide**
Analysis of trimethobenzamide in human saliva by GC-MS (224) 116

**Trimethoprim**
Simultaneous determination of trimethoprim, sulphamethoxazole and $N^4$-acetylsulpha-methoxazole in serum and urine by HPLC (223) 139

Assay of trimethoprim, sulfadiazine and its $N^4$-acetyl metabolite in biological fluids by normal-phase HPLC (224) 59

Quantitative TLC of trimethoprim and tetroxoprim using fluorescence densitometry (230) 212

**Trithiozine**
HPLC determination of trithiozine and its neutral metabolites in human plasma and urine (222) 257

**Tryptophan**
Analysis of some tryptophan and phenylalanine metabolites in urine by a straight-phase HPLC technique (222) 191

Determination of indoles and catechols in rat brain and pineal using LC with fluorometric and aperometric detection (223) 315

Determination of indoles in human and rat pineal (228) 155

**Tubocurarine**
Quantitation of D-tubocurarine in human plasma using HPLC (226) 255

**Tulobuterol**
Quantitative determination of tulobuterol and its metabolites in human urine by mass fragmentography (222) 53

Determination of tulobuterol in human serum by GLC (230) 148

**Tyramine**
Simultaneous extraction and separation of trace amines of biological interest (225) 301

GC-MS analysis of amine metabolites in urine (228) 51

**Tyramines**
Determination of noncatecholic phenylethyl-amines and monomethylated derivatives of phenylethylamine (224) 481

# U

**UK33274**
Determination of the vasodilator UK33274 by HPLC using fluorescence detection (221) 193

**Uracil**
Determination of 5-fluorouracil and pyrimidine bases in plasma by GC-chemical ionization-mass fragmentography (221) 382

**Uracil arabinoside**
Determination of cytarabine and uracil arabinoside in human plasma and cerebrospinal fluid by HPLC (225) 99

**Ureidopenicillins**
Improved procedure for the determination of the ureidopenicillins azlocillin and mezlocillin in plasma by HPLC (228) 409

**Uric acid**
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215

Simultaneous determination of allopurinol, oxipurinol and uric acid in human plasma by HPLC (228) 382

**Urine**
High-performance anion-exchange chromatography of human urine using perchlorate gradient elution systems (221) 215

**Urocanic acid**
HPLC determination of urocanic acid isomers in biological samples (230) 125

**Uronic acid**
Analysis of the monosaccharide compositions of total non-dialyzable urinary glycoconjugates by the dithioacetal method (226) 341

**Uroporphyrinogen decarboxylase**
The determination of uroporphyrinogen decarboxylase in tissues by HPLC coupled to fluorescence detection (229) 311

**UV detection**
LC determination of drugs in urine by direct injection on to a reversed-phase column. Fluorescence versus UV detection (225) 476

# V

**Valproic acid**
Performance evaluation of a reversed-phase, HPLC assay of valproic acid involving a "solvent demixing" extraction procedure and precolumn derivatisation (224) 289

Rapid GC determination of valproic acid in serum (225) 459

LC determination of valproic acid in plasma with spectrofluorimetric detection (in French) (225) 509

Valproic acid and several metabolites: quantitative determination in serum, urine, breast milk and tissues by GC-MS using selected ion monitoring (226) 69

Valproic acid analysis in saliva and serum using selected ion monitoring (electron ionization) of the *tert.* -butyldimethylsilyl derivatives (227) 433

**Vanillylmandelic acid**
Determination of vanillylmandelic acid in urine by pre-column dansylation using micro HPLC with fluorescence detection (223) 176

Simple high-performance liquid chromatographic method for the concurrent determination of the amine metabolites vanillylmandelic acid, 3-methoxy-4-hydroxyphenylglycol, 5-hydroxyindoleacetic acid, dihydroxyphenylacetic acid and homovanillic acid in urine using electrochemical detection (226) 361

Simple method for the determination of homovanillic acid and vanillylmandelic acid in urine by HPLC (227) 162

**Vasopressin**
Characterization of neuropeptides by reversed-phase, ion-pair LC with post-column detection by radioimmunoassay. Application to thyrotropin-releasing hormone, substance P, and vasopressin (222) 381

**Verapamil**
Simultaneous determination of verapamil and its seven metabolites by HPLC (222) 507

**Vidarabine**
Reversed-phase HPLC method to determine vidarabine and hypoxanthine arabinoside in biological fluids (229) 487

**Vioform**
GC determination of clioquinol (Vioform) in human plasma (226) 219

**Vitamin $B_2$**
Determination of the $B_2$ vitamer flavin-adenine dinucleotide in whole blood by HPLC (228) 311

**Vitamin $B_6$**
Separation of vitamin $B_6$ compounds by paired-ion HPLC (227) 181

**Vitamin C**
Ultramicromethod for the measurement of ascorbic acid in plasma and white blood cells by HPLC with electrochemical detection (224) 477

**Vitamin D**
Ternary solvent mixtures for improved resolution of hydroxylated metabolites of vitamin $D_2$ and vitamin $D_3$ during HPLC (221) 27

Silica Sep-Pak preparative chromatography for vitamin D and its metabolites (226) 198

High-performance liquid chromatography of 25-hydroxyvitamin $D_2$ and 25-hydroxyvitamin $D_3$ in human plasma. Use of isotachysterols and a comparison with GC-MS (226) 351

Separation of nanogram quantities of hydroxy metabolites of vitamin $D_3$ in plasma by TLC on silica gel (230) 319

**Vitamin E**
Quantitation of individual tocopherols in plasma, platelets, lipids, and livers by HPLC (224) 389

Improved assay for α-tocopherol in the picogram range, using GC-MS (225) 433

Quantitation of physiological α-tocopherol, metabolites, and related compounds by reversed-phase HPLC (227) 174

**Vitamins**
Determination of nicotinamide in human plasma and urine by ion-pair reversed-phase HPLC (221) 161

LC determination and time-concentration studies of riboflavin in hemodialysate from uremic patients (226) 471

Determination of nicotinamide and metabolic products in urine by HPLC (230) 309

**VM 26**
Analysis of the anticancer drugs etoposide (VP 16-213) and teniposide (VM 26) by HPLC with fluorescence detection (224) 168

**Volatile compounds**
Volatiles of exogenous origin from the human oral cavity (226) 315

**Volatile organic compounds**
A GC-MS study of profiles of volatile metabolites in hepatic encephalopathy (226) 291

**VP 16-213**
HPLC determination of 4'-demethylepipodophyllotoxin-9-(4,6-0-ethylidene β-D-glucopyranoside) (VP 16-213) in human plasma (222) 141

Analysis of the anticancer drugs etoposide (VP 16-213) and teniposide (VM 26) by HPLC with fluorescence detection (224) 168

# W

**Warfarin**
Rapid, reliable and sensitive assay for warfarin using normal-phase HPLC (228) 346

# X

**Xanthine**
Mass fragmentographic determination of xanthine and hypoxanthine in biological fluids (221) 9

Simultaneous determination of allopurinol and oxipurinol in biological fluids by mass fragmentography (229) 211

**Xylazine**
Determination of xylazine in plasma using HPLC (222) 308

# Y

**YM-09538**
Determination of the α,β-adrenoceptor blocker YM-09538 in plasma by HPLC with fluorescence detection (225) 115

# Z

**Zymograms**
Isoelectric focusing studies of human pancreatic secretion (221) 279

## PUBLICATION SCHEDULE FOR 1982

*Journal of Chromatography* (incorporating *Chromatographic Reviews*) and *Journal of Chromatography, Biomedical Applications*

| MONTH | J | F | M | A | M | J | J | A | S | O | N | D |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Journal of Chromatography | 234/1<br>234/2<br>235/1<br>235/2 | 236/1<br>236/2 | 237/1<br>237/2<br>237/3 | 238/1<br>238/2<br>239 | 240/1<br>240/2<br>241/1 | 241/2<br>242/1<br>242/2 | 243/1<br>243/2<br>244/1<br>244/2 | 245/1<br>245/2<br>245/3 | 246/1<br>246/2<br>247/1 | 247/2<br>248/1<br>248/2<br>248/3 | 249/1<br>249/2<br>250 | |
| Chromatographic Reviews | | 251/1 | | 251/2 | | | | | | | | 251/3 |
| Biomedical Applications | 227/1 | 227/2 | 228 | 229/1 | 229/2 | 230/1 | 230/2 | 231/1 | 231/2 | 232/1 | 232/2 | 233 |

## INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 244, No. 2, pp. 401–404. A free reprint can be obtained by application to the publisher.)

**Types of Contributions.** The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications:* Regular research papers (Full-length papers), Short communications and Notes. Short communications are preliminary announcements of important new developments and will, whenever possible, be published with maximum speed. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed four printed pages. For review articles, see page 2 of cover under Submission of Papers.

**Submission.** Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

**Manuscripts.** Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (*e.g.*, Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

**Introduction.** Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

**Summary.** Full-length papers and Review articles should have a summary of 50–100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Short communications and Notes are published without a summary.)

**Illustrations.** The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

**References.** References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the lay-out of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press", "submitted for publication", "in preparation" or "personal communication".

**Proofs.** One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

**Reprints.** Fifty reprints of Full-length papers, Short communications and Notes will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

**News.** News releases of new products and developments, and information leaflets of meetings should be addressed to: The Editor of the News Section, Journal of Chromatography/Journal of Chromatography, Biomedical Applications, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

**Advertisements.** Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

# A COMPLETE SYSTEM FOR THIN LAYER CHROMATOGRAPHY

**always comprises devices for the following duties:**

## SAMPLE APPLICATION

Example: **CAMAG LINOMAT III**

for comparatively large volume samples (2–99 µl); sprayed-on in the form of narrow bands; reproducibility 1%.

## CHROMATOGRAM DEVELOPMENT

Example: **CAMAG HPTLC Linear Developing Chamber**

accommodates 32 samples on a 10 x 10 cm HPTLC plate or even 72 samples on a plate 20 x 10 cm.

## CHROMATOGRAM EVALUATION

**CAMAG TLC/HPTLC SCANNER**

the most versatile photodensitometer of high precision and sensitivity, for research and routine;
automatic scanning of a complete chromatogram.

If you are planning to set up a new chromatography laboratory, or to complement your existing TLC equipment, ask the CAMAG specialist. He will help.



CAMAG TLC/HPTLC SCANNER ▶

▲ CAMAG LINOMAT III

▲ CAMAG HPTLC Linear Developing Chamber

CAMAG offers COMPLETE SYSTEMS for linear, circular and anticircular HPTLC and, of course, systems for conventional TLC.

CAMAG · Sonnenmattstr. 11 · CH-4132 Muttenz/Switzerland
Telephone (061) 61 34 34 · Telex 62649
CAMAG has agents in almost all countries.

CAMAG